



Airborne pollen in Sassari (NW-Sardinia): a 3-years survey, comparison between two pollen samplers.

Canu A.¹, Peana I.², Fadda M.L.³, Fois G.², Mannu G.², Masia P.¹, Pellizzaro G.¹

¹Institute of Biometeorology, National Research Council, CNR-IBIMET, Sassari


²Sardinia Environmental Protection Agency, ARPAS, Meteorological Department, Sassari

³Institute of Ecosystem Study, National Research Council, CNR-ISE, Sassari

The main aim of the present study was to analyze and compare the results of 3-years (2015-2017) aerobiological sampling in Sassari between two Hirst type volumetric spore traps located in two different areas of the city.

The study was carried out in North Western Sardinia, Sassari Italy (40° 43' 24" N, 8° 33' 13" E, 120 m s.l.m.). The first pollen sampler **SS6** (CNR) was located in the center of the city very close to a public garden. The second one **SS5** (ARPAS) was placed in the outskirts of the city.



The pollen concentration of the 20 taxa more abundant in the atmosphere of Sassari was considered. The following parameters were calculated for each pollen: start, end and duration of pollen season, date of peak pollen concentration, number of days from the beginning of the season to the peak, annual pollen index (API), percentage distribution of API and maximum daily concentration.



Spearman correlation analysis was performed between daily pollen concentration data collected by the two samplers. To study the effect of different urban location on pollen concentration of various taxa an analysis of variance with the GLM procedure of SAS (version 8.2; SAS Institute, Cary, NC) was performed considering the location and the interaction between location and year as fixed factors.

Average pollen season characteristics in both stations (SS5, SS6) Sassari 2015-2017

	Start DOY	End DOY	Duration (N. days)	Peak DOY	N. days Start-Peak	API
Anacardiaceae	92	116	25 (±4.6)	99	7.7	550.9 (±182.1)
SS5	85	135	50.7 (±25.6)	102	17.3	171.3 (±50.9)
SS6	98	160	124 (±20.1)	46	8.3	157.2 (±116.9)
Betulaceae	38	140	102.7 (±50.4)	67	28.3	174 (±117.5)
SS5	137	270	134 (±3.1)	150	12.7	881.2 (±115.2)
SS6	122	263	141.7 (±15.2)	150	27.7	442.7 (±87)
Compositae	114	278	164 (±8.2)	143	28.3	251.9 (±73.4)
SS5	115	287	172.7 (±42.5)	184	69.3	225.4 (±122.4)
SS6	34	132	98.3 (±25.2)	97	63	158.1 (±104.8)
Corylaceae	37	151	49 (±48.9)	109	71.7	178.6 (±130)
Cupressaceae/Taxaceae	24	191	167.3 (±132.9)	55	30.7	13972 (±6903)
SS5	33	279	247.3 (±96.44)	67	34.3	9186.8 (±442.3)
SS6	58	185	127.5 (±26.2)	118	60	585.4 (±45.3)
Ericaceae	61	208	148 (±9.9)	117	56	740 (±187.8)
Euphorbiaceae	43	280	238.3 (±131.6)	51	8.0	350.8 (±283.6)
SS5	19	298	280.3 (±94.3)	77	56.7	165.6 (±84.3)
SS6	109	182	74.7 (±12.4)	127	18.3	5796.2 (±1295.5)
Fagaceae	109	195	87.7 (±8.66)	120	11.3	5820.5 (±3067.3)
SS5	117	207	91.3 (±15.4)	136	19.3	1618.7 (±623.2)
SS6	114	236	123.3 (±18.1)	144	30.0	1249.2 (±775.7)
Oleaceae	130	161	31.3 (±8.7)	141	11.0	9746.4 (±4970)
SS5	133	168	35.7 (±5.1)	145	11.7	5402.3 (±490.6)
SS6	130	160	30.3 (±8.1)	141	11.0	5314.2 (±4829.7)
Olea	133	163	30.7 (±4.2)	145	11.7	4885.9 (±491.1)
SS5	27	195	169 (±142)	102	74.7	345.5 (±194)
SS6	30	334	305.3 (±56.8)	167	136.7	318.7 (±82.1)
Pinaceae	80	181	101.7 (±48.4)	126	45.3	644.2 (±106.9)
SS5	68	209	142 (±12.53)	125	57.0	672.6 (±177.1)
SS6	100	224	124.7 (±18.18)	126	26.3	330.8 (±124.2)
Plantaginaceae	95	241	147.3 (±21.1)	166	71.0	97.1 (±30.2)
SS5	85	116	32.7 (±8.7)	97	12.0	388.3 (±212.3)
SS6	82.67	112	30.7 (±19.1)	97	14.0	288.9 (±263.9)
Polygonaceae	77	159	83.3 (±31.1)	100	23.0	299.9 (±82.5)
SS5	59	190	132.3 (±28.7)	96	16.7	295.2 (±115.4)
SS6	56	115	60 (±13.7)	62	6.0	258.5 (±82.3)
Salicaceae	51	101	50.3 (±7.5)	68	16.7	190.7 (±30.2)
SS5	42	101	60.7 (±31.6)	78	36.4	96.2 (±59.1)
SS6	45	105	61 (±10.4)	67	22	163.2 (±57.7)
Ulmaceae	78	289	212 (±28.8)	99	21.3	9077.3 (±2512.3)
SS5	75	278	204.7 (±28.5)	113	38.3	7319.2 (±3626.8)
SS6						



Average percentage over the whole study period (2015-2017) for the 20 studied taxa

Over the three years of study Cupressaceae, Oleaceae, Urticaceae, and Fagaceae accounted of about 85% of the total pollen counts recorded in both SS5 and SS6 sites but the percentage contribution of each taxa is different.

Significance of Spearman correlation coefficients between daily pollen concentration of SS5 and SS6 samplers

	n	r	sign
Anacardiaceae	152	0.662	***
Betulaceae	379	0.440	***
Amaranthaceae	447	0.520	***
Compositae	441	0.453	***
Corylaceae	414	0.385	***
Cupressaceae/Taxaceae	769	0.817	***
Ericaceae	382	0.560	***
Euphorbiaceae	873	0.569	***
Fagaceae	275	0.763	***
Gramineae	375	0.800	***
Oleaceae	116	0.303	***
Olea	100	0.734	***
Fraxinus	937	0.641	***
Pinaceae	438	0.628	***
Plantaginaceae	455	0.310	***
Plantago	119	0.310	***
Polygonaceae	406	0.506	***
Salicaceae	200	0.410	***
Ulmaceae	213	0.206	***
Urticaceae	693	0.780	***

Significant probabilities: **, P<0.01; ***, P<0.001

Effect of different urban location on pollen concentration observed in the various taxa

	Effect P-value		
	2015	2016	2017
Anacardiaceae	***	NS	***
SS5/SS6	***	NS	*
Betulaceae	***	NS	***
SS5/SS6	***	NS	***
Compositae	NS	NS	NS
SS5/SS6	NS	NS	NS
Corylaceae	NS	NS	NS
SS5/SS6	NS	NS	NS
Cupressaceae/Taxaceae	NS	*	*
SS5/SS6	NS	*	*
Ericaceae	NS	NS	NS
SS5/SS6	NS	NS	NS
Euphorbiaceae	NS	NS	NS
SS5/SS6	NS	***	***
Fagaceae	NS	NS	NS
SS5/SS6	NS	NS	NS
Gramineae	NS	NS	NS
SS5/SS6	*	NS	NS
Oleaceae	NS	NS	***
SS5/SS6	NS	NS	NS
Pinaceae	NS	NS	NS
SS5/SS6	NS	NS	NS
Plantaginaceae	NS	NS	NS
SS5/SS6	***	***	***
Plantago	NS	NS	*
SS5/SS6	NS	NS	NS
Polygonaceae	NS	NS	NS
SS5/SS6	NS	NS	NS
Salicaceae	NS	NS	NS
SS5/SS6	NS	NS	NS
Ulmaceae	NS	NS	NS
SS6	***	*	NS
Urticaceae	NS	NS	***
SS5/SS6	NS	NS	***

*, P<0.05; ***, P<0.001; NS no significant

Spearman correlation analysis between daily pollen concentration recorded during the three years showed a positive correlation and a highly significant coefficient values for all taxa.

Some differences between sites were found when data were analyzed for each year separately. The analysis of variance showed highly significant differences in all three years for pollen of Plantaginaceae, and in two years for Anacardiaceae, Amaranthaceae and Euphorbiaceae. No difference among the sites was highlighted for some important taxa including Compositae, Corylaceae and Fagaceae.


Pollen of Plantaginaceae



Maximum pollen daily concentration



Annual Pollen Index



Some differences on maximum pollen daily concentration were observed between the two sites. However these differences were not found every year for each taxa. We in fact observed, for the same taxa, years in which the peak concentration was similar in both sites, and years in which it was really different (e.g. Oleaceae and Olea in 2016 and 2017 respectively; Fagaceae in 2017 and 2016 respectively).

Analysis of the pollen index for each pollen type and each study year revealed in some cases evident differences in the amount of pollen produced by the species present in the area surrounding the samplers. Clear differences in annual pollen index values were observed every year in Amaranthaceae and Plantaginaceae taxa. This could be probably due to a greater presence of these species in the neighborhood of the traps. For the others taxa we observed differences between sites only in some years.