

State of nature in the EU

Results from reporting under the nature directives 2013-2018

ISSN 1977-8449



State of nature in the EU

Results from reporting under the nature directives 2013-2018

Cover photo: © Daniel Danko, Environment & Me/EEA

Legal notice

The contents of this publication do not necessarily reflect the official opinions of the European Commission or other institutions of the European Union. Neither the European Environment Agency nor any person or company acting on behalf of the Agency is responsible for the use that may be made of the information contained in this report.

Brexit notice

The withdrawal of the United Kingdom from the European Union did not affect the production of this report. Data reported by the United Kingdom are included in all analyses and assessments contained herein, unless otherwise indicated.

Copyright notice

© European Environment Agency, 2020

Reproduction is authorised provided the source is acknowledged.

More information on the European Union is available on the Internet (<http://europa.eu>).

Luxembourg: Publications Office of the European Union, 2020

ISBN 978-92-9480-259-0

ISSN 1977-8449

doi:10.2800/088178

European Environment Agency
Kongens Nytorv 6
1050 Copenhagen K
Denmark

Tel.: +45 33 36 71 00

Web: eea.europa.eu

Enquiries: eea.europa.eu/enquiries

Contents

Authors and acknowledgements	4
Executive summary	5
The State of nature report in a nutshell	8
1 Introduction	9
2 Status of and trends in bird populations	15
2.1 Facts and figures	17
2.2 Population status and trends	18
2.3 Population status of species with species action plans.....	27
2.4 Improvement in trends.....	29
3 Status of and trends in habitats and species	35
3.1 Facts and figures	38
3.2 Conservation status	41
3.3 Trends in conservation status	53
3.4 Improvement in status and trends	62
4 Pressures and responses	71
4.1 Key pressures for species and habitats.....	71
4.2 Conservation measures.....	88
4.3 Restoration needs of habitats	100
5 Status of the Natura 2000 network	107
5.1 Facts and figures	108
5.2 Land use in Natura 2000 sites	112
5.3 Coverage of species and habitats by Natura 2000.....	115
5.4 Effectiveness of the Natura 2000 network.....	118
5.5 Actions needed to strengthen the Natura 2000 network	122
6 Meeting EU 2020 Biodiversity Strategy targets	127
6.1 Measuring progress towards target 1	127
6.2 Measuring progress towards target 3	129
7 Conclusion and outlook	136
Abbreviations	138
References	139

Authors and acknowledgements

This report was developed and written by Sandra Naumann, Rebecca Noebel, Zelmira Gaudillat, Ulf Stein, Lina Röschel, Sophie Ittner and McKenna Davis (European Topic Centre on Biological Diversity (ETC/BD)), Anna Staneva and Claire Rutherford (BirdLife International under contract of the European Commission) and Carlos Romão (European Environment Agency (EEA)).

Significant support in data processing and assessment, as well as guidance, was provided by Sabine Roscher, Nirmala Séon-Massin, Jérôme Bailly Maitre, Pablo Bolanos, Laura P. Gavilán, Thomas Ellmauer, Helmut Kudrnovsky, Dietmar Moser, Lubos Halada, Manuel Löhnertz, Mora Aronsson, Michelle Watson, Ruta Landgrebe, Katrina Abhold (ETC/BD), Eleni Tryfon (EEA), Eresha Fernando (BirdLife International), Ben Delbaere (NEEMO, monitoring of LIFE projects), and Angelika Rubin and Sofia Pachini (DG Environment).

The quality and breadth of the State of Nature report would not have been possible without the dedicated work of these individuals. We would also like to thank those who provided valuable feedback to the final draft of the report during the Eionet consultation as well as members of the Expert Group on Reporting under the Nature Directives and the Directorate-General for the Environment, European Commission, who also provided important input.

The assessments included in the report synthesise the work of the EEA's European Topic Centre on Biological Diversity. They are based on the information reported by EU Member States as part of the implementation of the Birds Directive (Article 12) and the Habitats Directive (Article 17) as well as a variety of other sources, such as the European Environment Information and Observation Network (Eionet), Corine Landcover and the Natura 2000 database.

Executive summary

Introduction

The EU nature directives — i.e. the Habitats and Birds Directives — coordinate conservation efforts for more than 2 000 species and habitats across the EU Member States with the aim of maintaining them at or restoring them to a favourable conservation status. This report describes the state of nature in the EU during the period from 2013 to 2018, based on Member States' reporting under these directives and on subsequent assessments at EU or EU biogeographical and marine levels. The data collected serve to identify successes and shortcomings in nature conservation, key pressures and threats, the status of current conservation measures and the restoration needed to further improve the conservation status of targeted habitats and species and the population status of birds.

The report also gives insights into the Natura 2000 network and its contribution to achieving favourable conservation statuses and to making progress towards targets 1 and 3 of the EU Biodiversity Strategy to 2020. Limitations due to poor data quality or gaps in data completeness still exist. A noteworthy portion of the reported information comes from expert opinion and partial surveys, indicating the absence or incomplete monitoring schemes in Member States. Nevertheless, the data presented are a milestone in assessing the state of nature in the EU and provide a strong basis for improving reporting, assessment and implementation to be able to achieve the aims of the EU Biodiversity Strategy for 2030.

Status of and trends in bird populations (Chapter 2)

This chapter provides an overview of the size of and trends in bird populations at the national level (reported under Article 12 of the Birds Directive) and of EU-level population statuses and trends for 463 bird species that naturally occur in the EU.

Around half of the species have a good EU status, which is slightly less (5 %) than during the last reporting period (2008-2012). The proportion of species with poor and bad status, however, has increased by 7 % in the last 6 years to reach a total of 39 %. This is partly

because of a mixture of changes in survey and analysis methodology and in data quality but also because of genuine deterioration. For breeding birds, increasing trends in populations are reported in the short and long term for around one quarter and one third of assessed taxa, respectively; one third of taxa have decreasing trends in population in the short and/or long term. The proportion of wintering birds with increasing short- and/or long-term population trends has dropped by 9 % compared with the previous reporting period. A quarter of Annex I taxa of the Birds Directive show a deterioration in their status, despite the implementation of dedicated conservation measures.

At the Member State level, around 50 % of improving population trends concern Annex I and Special Protection Area trigger species, most of which are wetland and marine birds. Breeding birds have the highest share of reports (about 80 %) showing improving population trends. Such improvements are due to the implementation of habitat protection or restoration measures and to improvements in knowledge due to research and better monitoring and awareness-raising activities. Farmland and forest birds, however, show few improving trends. Overall, the high percentage of unknown data across bird taxa highlights the need to establish or re-enforce appropriate, coordinated, state-supported monitoring schemes in all Member States.

Status of and trends in habitats and species (Chapter 3)

This chapter provides an overview of data and assessments reported by Member States under Article 17 of the Habitats Directive as well as assessments of conservation status and trends at the EU biogeographical and marine levels, covering 233 habitats and 1 389 species. Only 15 % of **habitat** assessments have a good conservation status, with 81 % having poor or bad conservation status at EU level and 4 % reported as unknown. Over 50 % of dune habitats and bog, mire and fen habitats have a bad conservation status. Compared with the previous reporting period, the share of habitats with bad conservation status has increased by 6 %. These

differences largely relate to changes in the methods used or variations in data quality.

Across the different terrestrial regions, the Atlantic and Steppic regions stand out with only 4 % and a notably high 72 % of habitat assessments with good conservation status, respectively. Marine regions have fewer habitats with good status than their terrestrial counterparts, with good conservation status reported only for the Marine Black Sea region and an overall high share of unknown assessments. Looking at habitat trends, only 9 % of all habitat assessments with poor or bad conservation status show improvement, while 36 % continue to deteriorate at the EU level. Grasslands, dunes, and bog, mire and fen habitats show strong deteriorating trends, while forests have the most improving trends. On average, 6 % of all habitat assessments — representing 105 habitat types — show improvements in conservation status and/or trends due to targeted management and restoration measures or expansions of the habitat area.

Around a quarter of **species** have a good conservation status at EU level, which is an increase of 4 % compared with the previous reporting period (presumably due to changes in the methods used or variations in data quality). Reptiles and vascular plants have the highest proportion of good conservation status (35 %). In contrast to habitats, the majority of terrestrial species assessments show higher shares of good than bad conservation statuses. Nevertheless, the two terrestrial regions with the lowest share of good conservation status assessments are the same as for habitats (i.e. Atlantic and Continental). Over half of the assessments report a poor or bad status.

The marine regions have many assessments with unknown conservation status, reflecting the general lack of marine species data. The Marine Baltic region has no unknown assessments but the highest percentage of species with bad conservation status (70 %). Looking at conservation status trends, only 6 % of species are improving (largely fish and mammals) and 33 % are unknown (largely marine mammals). Looking at improvements in status and/or trends since the last reporting period, these positive changes also mostly involve mammals and fish as well as plant species.

Pressures and responses (Chapter 4)

This chapter looks at the pressures and measures reported by Member States. Although the drivers of habitat degradation and species decline are diverse, agricultural activities such as abandoning extensive management and intensifying management practices are the most common pressure overall. Urbanisation is the second largest pressure, which especially affects habitats such as dunes and coastal and rocky habitats. Forestry

activities are the main pressure on species (e.g. arthropods, mammals and non-vascular plants). Pollution of air, water and soil from different origins affects most habitats, particularly in the Atlantic and Continental regions.

The majority of habitat groups are also vulnerable to natural processes (e.g. succession of semi-natural habitats). Wintering and passage birds are threatened by the exploitation of species (i.e. hunting and illegal killing). Further pressures include invasive alien species (affecting habitats more than species), modifications to hydrological flow (freshwaters), physical alterations to water bodies (freshwater fish), energy-related pressures (hydropower installations — freshwater fish; electricity and communication transmissions — mostly birds), pollution and climate change (droughts and decreases in precipitation — mostly amphibians). Pollution, particularly from agricultural activities and urbanisation, negatively affects a wide range of habitats and species.

The nature directives emphasise the need for conservation measures to respond to these pressures with mandatory requirements for Natura 2000 sites. For habitats, most of the measures applied aim to maintain forests, grasslands, freshwater and coastal habitats and include actions such as reducing agricultural pressures by reinstating appropriate grassland management or reducing diffuse pollution and the use of fertilisers. For non-bird species, the measures applied largely aim to maintain the status of vascular plants, arthropods, mammals and fish by, for example, preventing the loss of their natural habitats.

Approximately 40 % of the reports for birds indicate that measures have been taken, predominantly targeting breeding birds. For most habitats and species, there is a positive link between measures taken (especially proactive restoration measures to improve structure and functions) and a good conservation status. However, conservation status and trends have not significantly improved and deterioration is ongoing. These findings highlight the importance of implementing those measures that have not yet been taken or scaling up and monitoring those currently being implemented. In addition, at least 226 000 km² of Annex I habitats still need to be restored to ensure their long-term viability.

Status of the Natura 2000 network (Chapter 5)

Natura 2000 represents the largest coordinated network of nature conservation areas in the world, covering almost one fifth of the EU's terrestrial land area and approximately 10 % of its seas. The network consists of 'Special Protection Areas' under the Birds Directive and 'Special Areas of Conservation'/Sites of Community Importance' under the Habitats Directive, totalling almost 28 000 sites. Forests and

grasslands make up over 60 % of the network's terrestrial area. Land use changes within the network show a decrease in grasslands and forests (albeit less than outside the network) and an increase in arable land and permanent crops (despite their having decreased outside the network).

On average, species are better covered by the Natura 2000 network than habitats. While vascular plants and terrestrial reptiles are particularly well covered, fish and marine reptiles (except for their nesting sites) are much less well covered. Scrubs, heaths and dunes are the best covered habitats; forest coverage, however, falls rather short. Coverage of birds is generally low, especially for breeding birds with large breeding distribution areas and dispersed nesting sites (making it difficult to include all of these areas within the network).

Measuring the ecological effectiveness of a network of protected areas is difficult, as baseline data are scarce and the data have many data limitations, such as the lack of data enabling comparison of the conservation status of and trends in species and habitats inside and outside of the Natura 2000 network. Statistics for this assessment link the level of coverage of Annex II non-bird species and Annex I habitats and the proportion of good status assessments and reported changes in status and improvements in trends between reporting periods. The results show that a good conservation status and less deterioration/decline are more likely for species and habitats that are well covered by the network.

Although the results for birds are less conclusive, they indicate that decreasing population trends are sharper for species not targeted for protection under the Natura 2000 network (non-SPA species). Despite some positive impacts, however, the potential effectiveness of the network has not been achieved. Realising the network's potential requires an increase in the adaptive capacity of and functional connectivity between sites and the level of protection beyond sites; an increase in the effectiveness of management and monitoring inside and outside the network; addressing gaps in knowledge and communication; and streamlining biodiversity protection across key sectoral policies. These efforts should be supported by a reduction in the pressures that Natura 2000 sites face.

Meeting EU Biodiversity Strategy to 2020 targets (Chapter 6)

This chapter outlines the progress made towards targets 1 and 3 of the EU Biodiversity Strategy to 2020. Target 1 aims to achieve full implementation of the nature directives and, for a high percentage of European habitats and species, to maintain them at and restore

them to good conservation status by 2020. This target has not been reached: a 12 % gap needs to be closed to achieve the target for habitats; a 20 % gap remains for bird species; and a 2 % gap remains for non-bird species. In fact, high shares of assessments actually show further deterioration.

Target 3 aims to optimise agriculture and forestry's benefits for biodiversity. Target 3a strives to 'maximise areas under agriculture across grasslands, arable land and permanent crops that are covered by biodiversity-related measures under the Common Agricultural Policy'. No real progress has been made towards reaching this sub-target, and more than 45 % of Annex I assessments of agricultural habitats were assessed as bad. The status of and trends in agricultural habitats that are fully dependent on management measures are worse than those of partly dependent habitats such as natural grasslands. Furthermore, agricultural or farmland species are reported to have worse trends than forest species.


Target 3b aims to improve biodiversity and ecosystems affected by forestry. Assessments revealed that, while 31 % of forest habitat assessments have a bad conservation status, 32 % of forest bird species show an improving trend, 40 % a stable trend and only 16 % a deteriorating trend.

Conclusions and outlook

Despite significant efforts by Member States and some improvements, biodiversity in the EU continues to decline and faces deteriorating trends from changes in land and sea use, overexploitation and unsustainable management practices, as well as water regime modification, pollution, invasive alien species and climate change. Although some species and habitats show improvements, progress has not been sufficient to meet the objectives of the EU Biodiversity Strategy to 2020.

The new EU Biodiversity Strategy for 2030 strives to be more successful and ensure that ecosystems are healthy, resilient to climate change, rich in biodiversity and able to deliver essential ecosystem services. Ambitious targets address the main drivers of biodiversity loss, but they need to be accompanied by greater implementation action to be effective. Additional effort is also needed to improve Member States' monitoring capacities to fulfil the requirements of the nature directives, develop new indicators to evaluate the impact of the Natura 2000 network on biodiversity policy objectives (i.e. nature directives and EU Biodiversity Strategy for 2030) and ensure sufficient financial and human capacities to underline a robust governance and policy framework to fully implement the nature directives and relevant environmental regulations.

The State of nature report in a nutshell



1 Introduction

The EU nature directives — i.e. the Habitats and Birds Directives (EU, 1992, 2009) — coordinate conservation efforts for more than 2 000 species and habitats across all Member States in the EU. Their overarching aim is to restore or maintain these species and habitats to a favourable conservation status in the long term. At the centre of these policies is the Natura 2000 network, which covers almost 20 % of terrestrial land and 10 % of marine waters in the EU, making it the largest coordinated network of nature conservation areas in the world.

Every 6 years, EU Member States are required to report on the sizes of and trends in populations of birds (Article 12 of the Birds Directive) and on the conservation status of and trends in targeted habitats and species (Article 17 of the Habitats Directive) within their European territories⁽¹⁾. Reporting consists of three main steps: (1) collecting the data in reports for each Member State; (2) processing Member States reports; and (3) assessing the reported data. Details of each step are summarised in the graphic below (see Figure 1.1). The EEA and its European Topic Centre on Biological Diversity (ETC/BD) provide technical and scientific support to the European Commission and the EU Member States throughout all stages of the reporting process, and several working groups also provide assistance. In total, more than 220 000 people across Europe (almost 60 % of whom are volunteers) have contributed to this process.

The outcomes of Article 12 and Article 17 reporting form the basis of the *State of nature in the EU* report. Through streamlined monitoring processes based on a set of minimum standards, Member States reports provide a baseline for measuring the status of and trends in species and habitats over multiple monitoring periods (i.e. between 2000 and 2006, 2007 and 2012, and 2013 and 2018 for the Habitats Directive and 2008 and 2012, and 2013 and 2018 for the Birds Directive). The data collected serve to identify successes in nature conservation, shortcomings in current protection measures and additional actions and measures needed

to further improve the conservation status of targeted habitats and species and the population status of birds.


The EU population status of birds under the Birds Directive and the EU conservation status of habitats and species under the Habitats Directive (hereafter referred to as EU status) are evaluated using various categories and assessment methods. However, overarching categories such as 'good', 'poor', 'bad' and 'unknown' have been introduced in this report to ensure consistent communication of the EU conservation and population status under the Habitats and Birds Directives (see Table 1.1).

Article 12 and Article 17 reporting data also support other EU and global policies and contribute to wider conservation and management objectives. Data from nature directive reporting are used, for example, to measure progress towards meeting the targets of the EU strategies on biodiversity and towards meeting the global Aichi biodiversity targets. The findings from Member States' reporting also help to evaluate the implementation of the Sustainable Development Goals (SDGs) by identifying gaps and priority areas for action, e.g. SDG 13 (combat climate change and its impacts), SDG 14 (conserve the oceans, seas and marine resources) and SDG 15 (protect, restore and promote sustainable use of terrestrial ecosystems, sustainably manage forests, combat desertification, and halt and reverse land degradation and halt biodiversity loss).

This report, covering the period from 2013 to 2018, represents the most comprehensive overview of the state of nature in the EU. It also includes the first reporting of results from Croatia, the newest EU Member State. Unless specifically mentioned, the report includes data from the United Kingdom, as it was still an active EU Member State during the reporting period. Looking at the Member State and biogeographical and marine region levels as well as the EU level, the *State of nature in the EU* begins with the main results from the Article 12 and Article 17 reporting (Chapters 2 and 3, respectively), the analysis of the

(1) EU Overseas Countries and Territories and Outermost Regions are not covered by the nature directives.

Figure 1.1 The nature directives' reporting process



Source: Data on staff involved in data collection and assessment and report preparation under Article 12 of the Birds Directive and Article 17 of the Habitats Directive were gathered through a survey of Member States conducted in spring 2020.

Table 1.1 Categories of EU population status of birds and EU conservation status of habitats and species

EU Status	Good	Poor	Bad	Unknown
Categories of EU population status of birds	Secure	'Near threatened' (close to being at high risk of extinction in the near future according to IUCN Red List criteria), 'Declining' or 'Depleted'	'Threatened' (according to IUCN Red List criteria)	Unknown
Categories of EU biogeographical conservation status of habitats and species of Community interest	Favourable	Unfavourable- inadequate	Unfavourable-bad	Unknown

latter being also conducted at the biogeographical level. This includes the status of and trends in populations of birds and non-bird species and of habitats as well as needs for further improvement and restoration. Case studies are presented throughout the report, highlighting the assessments for which an EU status or trend has improved since the previous reporting period. Pressures and responses (i.e. in the form of conservation measures) are presented in Chapter 4 for habitats, non-bird species and birds. Furthermore, a dedicated section presents an in-depth analysis of additional restoration needs for habitats. Chapter 5 focuses on the Natura 2000 network and highlights its current status, changes in its coverage and its contribution to the EU status of birds, non-bird species and habitats. Finally, Chapter 6 outlines progress towards meeting targets 1 and 3 of the EU Biodiversity Strategy to 2020. On this basis, a series of conclusions are presented, reflecting on the information gathered and the actions needed with a view towards the next reporting period and the implementation of the EU Biodiversity Strategy for 2030.

The data collected by Member States on the status of Habitats Directive habitats, birds and non-bird species and on underlying pressures are a milestone in assessing the state of nature in the EU. Both the Habitats and Birds Directives require appropriate monitoring systems to be in place to provide the relevant information for producing such an all-encompassing assessment. Ideally, the data would

have been collected from robust and comprehensive surveys and using methods comparable across all Member States. In many cases, however, the reported information comes from partial surveys that were performed for different purposes. In other cases, suitable data do not exist and expert opinion has been sought. For Habitats Directive habitats and species, more than 40 % of the reported information comes from partial surveys where the estimates cannot be considered robust or representative of entire biogeographical regions. In fact, more than 20 % of the information reported by Member States is based only on expert judgement. The situation for bird data is similar, with more than 30 % of the information coming from partial surveys and more than 15 % being based on expert judgement. Data quality issues are further accentuated by numerous gaps in the information reported. Population trends over the last 12 years, for example, are an essential part of assessing the status at EU level for both Habitats Directive species and birds. However, trends were unknown or missing in 20 % of Member States' reports for breeding birds and 30 % for wintering birds as well as in 34 % of reports for Habitats Directive species.

It is worth mentioning that international multilateral agreements such as the Bern Convention and the Agreement on the Conservation of African-Eurasian Migratory Waterbirds have also established reporting processes. These efforts and their linkages to the nature directives' reporting are explained in Box 1.1.

Box 1.1 Bern Convention and Agreement on the Conservation of African-Eurasian Migratory Waterbirds supporting biodiversity reporting

In 2018, eight Contracting Parties to the Bern Convention reported for the first time under Resolution 8 (2012) on the conservation status and trends and the population trends of a limited number of habitats (nine) and species (12 birds and 25 non-bird species). This first reporting period covered the years 2013-2018 and the reporting format was fully harmonised and standardised with that of Article 12 and Article 17 reporting under the Birds and Habitats Directives, respectively. This first reporting round is clearly more important for building up experience and capacity, but nevertheless it also creates various opportunities to analyse, arrange and present data from all European countries together. Analyses at pan-European level are especially important for migratory and wide-ranging species, as for them isolated conservation efforts in some countries may not yield the desired results. In that sense, the contribution of non-EU Contracting Parties is particularly important for features for which significant proportions of their global resources occur outside the EU.

Moreover, the Agreement on the Conservation of African-Eurasian Migratory Waterbirds (AEWA) has fully streamlined its reporting procedures with those under Article 12, thereby avoiding Member States having to report twice.

Source: Roekaerts et al., 2020.

Moreover, reporting data were used and published in the latest IPBES ⁽²⁾ *Assessment report on biodiversity and ecosystem services for Europe and Central Asia*.

The State of nature in the EU report is accompanied by a methodological technical report, which describes the data and analyses in more detail; this is available as an ETC/BD report (Röschel et al., 2020). Further analyses, information and data are available from the following sources:

- State of nature in the EU online ⁽³⁾
- Biodiversity Information System for Europe ⁽⁴⁾
- webtools for Article 12 ⁽⁵⁾ and Article 17 ⁽⁶⁾.



DASHBOARD
with detailed information by Member State:
Article 12 data quality



DASHBOARD
with detailed information by Member State:
Article 17 data quality

⁽²⁾ Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services

⁽³⁾ <https://www.eea.europa.eu/themes/biodiversity/state-of-nature-in-the-eu>

⁽⁴⁾ <https://biodiversity.europa.eu>

⁽⁵⁾ <https://nature-art12.eionet.europa.eu/article12>


⁽⁶⁾ <https://nature-art17.eionet.europa.eu/article17>



Photo: © Josef Timar, REDISCOVER Nature/EEA



Figure 2.1 Summary of the Birds Directive reporting



2 Status of and trends in bird populations

The Birds Directive protects all wild bird species ⁽⁷⁾ that are naturally present in the European territory of the EU Member States, including over 460 species. It aims to protect those species from habitat destruction and pollution, deliberate killing and capture, significant disturbances, their sale and the destruction or taking of their eggs or nests. Common species are included, as well as rarer, threatened, near-threatened and other species requiring particular attention because of the specific nature of their habitat. Protection measures aim to cover the entire species lifecycle, including breeding, wintering and migration, and the areas where they occur.

Annexes I and II of the Birds Directive are important in terms of decision-making and policy, in particular when it comes to classifying protected areas or regulating the hunting of species in each Member State. The annexes contain a mix of taxonomic levels, depending on which of these was deemed appropriate to be included in each annex. Annex I of the Birds Directive lists 197 threatened taxa. Member States must establish targeted conservation measures, such as avoiding any significant disturbance, and designate Special Protection Areas (SPAs) for their survival. This also includes migratory bird species not covered by Annex I but that constitute a common heritage and require a broader international perspective to ensure their effective protection and conservation.

Annex II lists 86 taxa ⁽⁸⁾ that are permitted to be hunted, either in the entirety of their range within the EU or just in particular Member States. The Birds Directive protects these species from being hunted during their return migration to their breeding grounds and during their breeding season. Some overlaps exist between the annexes. For example, a species may have a subspecies that is considered to need the protection afforded by Annex I, although it can be hunted at the species level (and is hence also listed

in Annex II). Within the current report, all analyses relating to either Annex I or Annex II are based on the actual taxa found in the lists, and therefore they include both species and subspecies. Member States' reporting obligations under Article 12 of the Birds Directive serve to gather data on parameters such as the size of and trends in all regularly occurring wild bird populations, their breeding distribution and general information on the directive's implementation (e.g. main achievements, SPA classification, species action or management plans, and details on relevant introduced non-native bird species).

National Member State data are used to analyse population sizes and trends and to assess the population status of bird taxa at EU level. The exercise is based on a standardised methodology (Röschel et al., 2020). A total of 511 assessments were undertaken in this reporting period (see Box 2.1). Article 12 reporting also considers the seasonality in bird biology and distribution by distinguishing between breeding, wintering and passage (migratory) seasons. These are some of the main differences from the reporting under Article 17 of the Habitats Directive, where the conservation status and trends of (only a subset of) non-bird species and habitats is reported by the Member States themselves, and then assessed at biogeographical level within the country's territory, as a basis for the assessments conducted at EU biogeographical level (see Chapter 3).

This chapter thus provides the main results emerging from Member States reports and the EU-level assessments ⁽⁹⁾, representing the period from 2013 to 2018. In addition to highlighting the conservation needs of different taxa, several illustrative case studies are provided of bird species having shown improvements or deterioration in their population status and/or trends compared with the previous reporting period (2008-2012).

(7) In the context of this chapter, the word 'species' is used in the strict sense of bird populations considered at taxonomical species level only. Conversely the word 'taxa' (which is usually used as a general term to talk about a taxonomic group of any level) refers in this chapter (unless stated otherwise) to the selection of species, as well as subspecies and biogeographic populations, and their counterparts, that were assessed for this report.

(8) These include four non-native/feral species.

(9) Performed by BirdLife International in the context of the European Red List of Birds under a contract from the European Commission.

Box 2.1 EU assessments of the status of bird populations — what is included in this report

Based on the Article 12 Member States' reports, the EU population size and trend calculations and EU population status assessments of 511 taxa (all taxonomic levels included) were undertaken. These include:

- 463 species regularly occurring in the EU and breeding and/or wintering (with only one exception — *Phasianus colchicus* — whose long-standing introduced population was also assessed);
- 41 selected subspecies and their counterparts (including one subspecies that became extinct before the 2008-2012 reporting period, *Perdix perdix italiae*); please note that two other globally extinct taxa — *Pinguinus impennis* and *Haematopus meadewaldoi* — are not covered in this report, as Member States were not asked to report against these for the period 2013-2018;
- two biogeographical populations of *Anser brachyrhynchus*;
- three introduced or feral species (*Meleagris gallopavo*, *Branta canadensis* and feral populations of *Columba livia*), although Member States reported on a greater number of non-native species;
- two species that are native to the EU, but for which the data reported are too incomplete for an overall evaluation (*Alle alle* and *Larus ichthyaetus*);

It should be noted that there are more wintering and breeding taxa than reported by the Member States. Moreover, since the last publication of the *State of nature in the EU* report in 2015, 13 new species were split from other taxa and five previously split species were lumped into two species. Such taxonomical changes are considered in the analysis.

Most birds have seasonal behaviours, i.e. have a breeding, wintering and passage (migratory) season. Member States were requested to report on all regularly occurring breeding taxa and on a subset of wintering and passage taxa. EU-level population trends were calculated for 465 and 467 breeding taxa in the short and long term, respectively (two taxa became extinct in the EU before the current short-term trend period, thus only long-term trends were calculated for them), and for 91 wintering taxa in the short and long term. Passage data were used only as supporting information. The total number of taxa for which trends were calculated does not equal the total number of EU population status assessments produced, as the latter is only based on either the breeding or the wintering season, i.e. one unique value per taxa (for more information, see Röschel et al., 2020).


The majority of the national breeding data come from well-established monitoring schemes conducted every year during the breeding season, when birds are holding territories and are usually most active and easy to detect (e.g. Pan-European Common Bird Monitoring Scheme ^(a)). It should be underlined, however, that many less common or rare species are still not well monitored and the lack of reliable information makes the assessment of their status very difficult (see also Chapter 1).

The information about wintering bird populations usually comes from coordinated monitoring efforts conducted during the wintering season. Most birds are not holding territories during this season and are usually much harder to record. Exceptions to this difficulty are waterbirds, most of which gather in large flocks on water bodies and for which rigorous monitoring procedures are in place to avoid double counting (e.g. International Waterbird Census ^(b)). As monitoring schemes for other groups of wintering birds are rare, Member States report on a subset of wintering taxa, called key wintering species, most of which are waterbirds.

Migration is an equally fascinating and difficult phenomenon to study. Many bird species spend weeks or months flying between their wintering and breeding grounds. Passage data are useful to collect to complement both breeding and wintering analyses, when looking in more detail at the results and the possible reasons behind them, but can hardly be part of any analyses in themselves, as passage data are highly variable and sporadically monitored and reported across the EU. For this reason, no assessments were undertaken based on passage data for this report.

Notes: (a) <https://pecbms.info>
(b) <https://europe.wetlands.org/news/international-waterbird-census-2020>


Figure 2.2 Share of Member States reports of Annex I, Annex II and non-Annex species



Note: Total number of reports: 8 008.

Source: Article 12 Member States' reports and assessments.

Figure 2.3 Share of taxonomic groups covered by Article 12



Note: Total number of species: 463.

Source: Article 12 Member States' reports and assessments.

2.1 Facts and figures



Member States report on regularly occurring breeding and wintering species and on some migrating birds (Figure 2.2). There are, however, many more species present in the EU, including marginal and vagrant species. The Birds Directive covers a wide range of taxonomic groups, the largest of which is passerines, followed by the waders, gulls and auks group. Together, these groups represent over half of the species reported under Article 12 (Figure 2.3).

It is important to highlight that 30 species (not including subspecies) present in the EU do not exist elsewhere i.e. they are endemic to the EU. Other 'near-endemic' species can be found outside the EU, but the vast majority of their populations reside within the EU. There are 24 near-endemic species in the EU, totalling more than 50 species for which breeding and/or wintering populations occur entirely or almost entirely in the EU. Thus, the EU has a unique responsibility to protect such species, because, if they disappear from the EU, their risk of becoming globally extinct becomes very high. This responsibility is transferred to the Member States within which endemic and near-endemic species are present. Countries such as Spain⁽¹⁰⁾ have a comparatively high proportion of endemic species within their territories as well as large proportions of many EU species' populations.



Maps 2.1 and 2.2 show the number of taxa reported by Member States for the breeding and winter season, respectively. The size of the country shows species numbers in relation to the country's size, e.g. large countries with relatively small numbers of species look smaller than they actually are, while small countries with large numbers of species look bigger. The colour coding shows the 'absolute' numbers of taxa reported by each Member State. The number of taxa is correlated not only with the size of a country but also with other environmental parameters such as habitat type, availability, climate. In addition, the varying amount of effort put into the survey in each Member State, combined with the difficulty of detecting certain taxa and the variable climatic conditions, particularly in the wintering season, can potentially skew these numbers.

⁽¹⁰⁾ For more information, see <http://datazone.birdlife.org/species/search>.

Map 2.1 Number of reported breeding taxa per country



Map 2.2 Number of reported wintering taxa per country



Note: These maps are cartograms, which distort the geometry of regions to convey specific information by resizing. Here, the size of the country shows the number of reported species in relation to the country's size. The maps do not show all species appearing in a country. Thus, the maps show the result of reporting rather than the species diversity of a country. The boxes at the top right refer to the Macaronesian islands (Azores, Madeira and Canary islands). Romania did not report and is therefore not included in the maps.

Source: Article 12 Member States' reports and assessments.

2.2 Population status and trends


This section presents the results for the status of and trends in bird populations at EU level and the breeding and wintering population trends at Member State level. As indicated in Table 1.1, overall categories such as 'good', 'poor', 'bad' and 'unknown' have been used in this report to ensure consistent communication on conservation and population status in the EU across the Habitats and Birds Directives. The results are compared with those from the previous reporting period (2008-2012).

2.2.1 Overall status of bird populations at EU level

Population status was assessed for a total of 463 bird species (see Box 2.1). As shown in Figure 2.4, 47 % of bird species assessed have a good population status in the EU. This is 5 % less than in the period 2008-2012.

Conversely, the proportion of species with poor and bad status has increased by 7 % in the last 6 years and reached a total of 39 %. The population status of 14 % of the bird species in the EU is still unknown because of a lack of reliable data regarding their population sizes and trends.

Figure 2.4 EU population status of bird species



Note: The total number of species-level assessments is 463 (only one assessment per species is undertaken, regardless of the number of seasons it has been reported in). Categories: good (secure); poor (near threatened, declining, depleted); bad (threatened).

Source: Article 12 Member States' reports and EU assessments.

2.2.2 Trends in populations of breeding birds

Population trends for bird taxa were reported by Member States for a short-term period of 12 years (2007-2018) and a long-term period of 38 years (1980-2018). As bird monitoring schemes were introduced in many Member States much later than 1980, the long-term trend period reported is often shorter, while the short-term trend data are usually more complete and generally of higher quality because of more accurate monitoring methods.

In total, 465 short-term and 467 long-term breeding population trends were produced for birds at the EU level (see Box 2.1). The results shown in Figures 2.5 and 2.6, compared with the previous reporting period (2008-2012), indicate that:

- The number of taxa with an increasing population is 5 % lower (meaning that less than one quarter of all taxa assessed now have increasing short-term trends).
- The number of taxa with stable or fluctuating trends is 7 % higher.
- Almost one third of all taxa assessed have a decreasing short-term trend.
- In the long term, there is only a slight dominance of decreasing over increasing trends. The opposite was true in the period 2008-2012, meaning that, although the current numbers are quite similar to those of the previous reporting round, the overall picture is slightly more negative.
- One in five taxa has an unknown or uncertain short-term or long-term population trend. Nevertheless, in the case of the long-term trends, this means 10 % less unknown trends.

A closer look at species level suggests that, in the majority of cases, the changes listed above (whether positive or negative) are genuine, which means that they are not related to better knowledge or to a change in the methods of bird monitoring or calculating trends.

At Member State level, Article 12 reports indicate a large variation in breeding bird populations (see Figure 2.7):

- Apart from Croatia, all Member States reported more than one third of breeding populations as increasing or being stable.

- Almost half of all Member States reported decreasing short-term trends for a third of breeding populations. Examples include the Corncrake (*Crex crex*) and Grey Partridge (*Perdix perdix*) (see also Table 2.1).
- The degree of unknown population trends varies greatly among the Member States. The degree of unknown population trends is related to whether bird monitoring systems are in place and how effective and informative they are.

2.2.3 Trends in populations of wintering birds

In total, 91 short-term and long-term wintering population trends were produced for birds at the EU level (see Box 2.1).

The results of the analysis of short- and long-term wintering population trends at EU level (shown in Figures 2.8 and 2.9), compared with the previous reporting period (2008-2012), indicate that:

- The percentage of increasing and decreasing short-term trends is similar (46 % and 27 %, respectively).
- The percentage of taxa with stable trends has increased and that of taxa with fluctuating trends has decreased, which is mainly due to the introduction of a more precise definition for these two trends.
- The proportion of taxa with increasing long-term trends has dropped by 9 %, while that of those with decreasing trends has barely changed.
- Cases in which the short- or long-term trend is unknown or uncertain have increased by 4 % and 2 %, respectively. Although there appears to be a lower percentage of unknown trends for wintering birds than for breeding birds, it must be noted that Member States report on only a subset of wintering birds, the selection of which is heavily reliant on the monitoring data available for these taxa. Therefore, it is more likely that the wintering birds reported on will have known trends, as their initial selection for reporting will be biased towards these species.

At the Member State level, two thirds of all Member states reported more than 30 % of wintering populations as increasing or stable, while approximately half of the Member States indicate them as decreasing at a rate of above 15 %. Austria

and Finland have the highest trend in reported increasing wintering populations in the short term (Figure 2.10). Examples include the Great White

Egret (*Ardea alba*), Whooper Swan (*Cygnus cygnus*) and White-tailed Eagle (*Haliaeetus albicilla*) (see Table 2.1). However, short- and long-term trends were reported

Figure 2.5 Short-term breeding population trends at EU level






Figure 2.6 Long-term breeding population trends at EU level



Note: Statistics based on 465 short-term and 467 long-term EU breeding population trends. Includes trends for a limited number of subspecies and biogeographical populations.

Source: Article 12 Member States' reports and EU assessments.

Figure 2.7 Short-term trends in breeding bird populations by Member State



Note: The number of reports is indicated in parentheses. The total number of reports is 5 827. There are more breeding taxa than reported by the Member States.

Source: Article 12 Member States' reports and EU assessments.

as unknown for many wintering birds across Member States. Croatia, Latvia, Portugal and Belgium reported over 50 % of reported wintering

bird populations as being unknown for short-term trends, while seven other Member States reported over 50 % as being unknown for long-term trends.

Figure 2.8 Short-term wintering population trends at EU level






Figure 2.9 Long-term wintering population trends at EU level



Note: Statistics based on 91 EU wintering population trends. Includes trends for a limited number of subspecies and biogeographical populations.

Source: Article 12 Member States' reports and EU assessments.

Figure 2.10 Short-term trends in wintering bird populations by Member State



Note: The number of reports is indicated in parentheses. The total number of reports is 1 284. There are more wintering taxa than reported by the Member States.

Source: Article 12 Member States' reports and assessments.

2.2.4 Status of and trends in breeding bird populations by taxonomic group at EU level

A taxonomic group is a selection of taxa grouped together on the basis of their taxonomy. Usually taxa are grouped together at a higher taxonomic level (e.g. order, family). Figure 2.11 presents an overview of the EU population status of bird species grouped by taxonomic order, as orders are the taxonomic level that provide the most information.

Many taxonomic orders are represented in the EU by very few taxa, while some orders have a large number of taxa. It is therefore difficult to draw conclusions on the status of and trends in the breeding population of birds within very large or very small taxonomic orders.


For taxa in different orders, the following observations were made:

- Over half of the taxonomic orders presented have a good status for at least 50 % of the species assessed.
- Almost half of the orders presented have a bad status for at least 25 % of the species assessed.

Included in these are the orders containing high numbers of species, such as petrels, storm-petrels and shearwaters (see example in Box 4.1), ducks, geese and swans, as well as those containing waders, gulls and auks, the pheasants, partridges and grouse, the bustard, and the falcons (see Box 2.2)

- The taxonomic orders with a higher number of taxa with bad status also show a greater proportion of decreasing trends relative to the last reporting round. This applies to both short- and long-term trends (trends are not shown in Figure 2.11).
- The long-term trends in passerine breeding populations, the largest group of birds, have a large number of uncertain and unknown trends. This group alone accounts for almost half (45 %) of all uncertain and unknown long-term trends and almost half (47 %) of all unknown population statuses.
- Swifts and nightjars, as well as petrels, storm-petrels and shearwaters, also have a high proportion of uncertain and unknown short- and long-term trends within their respective groups. This could be explained by the overall difficulty in monitoring these particular groups of species.

Figure 2.11 Population status of EU bird species and subspecies, by taxonomic order



Note: The total number of all species is 463. The number of taxa concerned is shown in parentheses.

Source: Article 12 Member States' reports and EU assessments.

Box 2.2 Closer look: Birds of prey

Hen Harrier (*Circus cyaneus*) © Peter Harris, IUCN Red List

'Birds of prey' encompasses a diverse group of birds, such as vultures, eagles, kites, harriers, buzzards, honey buzzards, hawks and falcons. Many species within this group have been the target of conservation measures during the last two decades, particularly through the LIFE programme. While this has led to certain levels of recovery or stabilisation of some populations (e.g. vultures, some falcons and eagles), other groups, such as falcons and harriers, still face many outstanding threats.

Falcons and harriers: deteriorating populations

The most common threats to and pressures on falcons and harriers reported by Member States are land use change (various forms of agriculture) and different land use practices, electricity infrastructure, disturbance, use

of chemicals and illegal killing. Out of 10 **falcon species** covered by Member States' reporting, four are showing evidence of genuine deterioration in both their status and their trend (*Falco biarmicus*, *F. columbarius*, *F. cherrug*, *F. rusticolus*). Five of the remaining species have not had a change in status, but their trends indicate a worrying outlook for the future: two have decreasing short- and long-term trends (which means that they may have a deteriorating status in the near future) and another two have been recovering but are currently stable (which indicates that the positive improvements have ceased). Interestingly, *F. eleonora* was assessed for first time, and Spain is the only country reporting increases and happens to be where the entire species population is within the Natura 2000 network. Finally, *F. peregrinus* is the only species with an overall increasing long- and short-term trend. **Harriers** are in a similar situation, with two of four species having genuinely decreased (i.e. *Circus cyaneus* and *C. pygargus*).

Vultures: improving and stabilising populations

Unlike falcons and harriers, vulture populations are improving and stabilising thanks to species action plans, LIFE projects and targeted large-scale conservation efforts. Three of four species reported on by Member States (*Aegypius monachus*, *Gyps fulvus* and *Gypaetus barbatus*) have increasing short- and long-term trends and one is stable (*Neophron percnopterus*). Despite these positive improvements, two of these species are still threatened because of steep historical declines. All of these four species are still in a relatively fragile situation, as they will depend on conservation measures until the ongoing pressures they still face, such as electricity infrastructure, illegal killing and harvesting and the poisoning of animals on which they feed, are addressed and managed.

Need for improvement and restoration of habitats

Further measures are needed to curb deteriorating population trends, improve the conservation statuses of birds of prey and ensure that their populations will stabilise or continue to improve in the future. Member States report that improving and restoring habitats as well as appropriate land use practices are the main conservation measures needed for these species. There are examples of beneficial agri-environmental practices that have been developed in projects such as the 'Locally led scheme for the conservation of the hen harrier in Ireland' ^(e), funded under the EIP-Agri (European Innovation Partnership 'Agricultural productivity and sustainability') scheme. In Groningen, the Netherlands, the introduction of set-aside land in the late 1980s and the implementation of dedicated agri-environment schemes in the 1990s (including field margins of between 9 m and 12 m as feeding areas) have been fundamental to the harriers' success. The Montagu's harrier population in Groningen now exceeds 60 breeding pairs (BirdLife International, 2011).

Note: ^(e) <https://ec.europa.eu/eip/agriculture/en/find-connect/projects/locally-led-scheme-conservation-hen-harrier>

2.2.5 Status of and trends in populations of birds listed in Annexes I and II of the Birds Directive at EU level

Figure 2.12 provides an overview of the population status of bird species listed in Annex I (species for which SPAs must be designated) and Annex II (huntable species) of the Birds Directive and also of species not listed in the annexes. The results, compared with those for the previous reporting period (2008-2012), indicate that:

- The proportion of taxa with good status listed in Annex I has decreased by 8 %, while those with poor and bad status have increased by 6 %. A closer look at these changes reveals that several species have undergone a genuine deterioration in their status despite the special conservation measures provided by Annex I of the Birds Directive.
- 16 % of Annex I taxa still have an unknown status. This is worrying, as a species should be well monitored and its status known so that it can be effectively protected. However, for some Member States, this could also be due to a mismatch between the timings of reporting under the Birds and Habitats Directives and the cycles of national monitoring schemes.
- The situation is similar with Annex II, in that the proportion of taxa with a good status is 9 % lower


and that of poor and bad taxa is 9 % higher. This also means that almost half of all species listed under Annex II now have a poor or bad status.

- When comparing species listed in the Birds Directive annexes and those not covered by special conservation measures, the results show that Annex I has a much higher proportion of taxa with unknown status (16 %). This can be partly explained by the fact that Annex I species are often rare and difficult to monitor. Nevertheless, as the relative share of unknown status has increased compared with the previous reporting period, it can be concluded that there has been a genuine increase in the knowledge gap for Annex I taxa and that greater effort should be put into studying and monitoring them to inform appropriate decision-making.

Annex II has a relatively small proportion of taxa with unknown status. This can be explained by the fact that huntable species are often common and generally better monitored and because many countries collect data on hunting bags. Standardised and regular monitoring should nevertheless be established across all Member States for Annex II species.

Table 2.1 below presents examples of breeding and wintering birds with an increasing or decreasing short-term trend in different EU Member States and their long-term trend at EU level.










Figure 2.12 EU population status of bird taxa in Annexes I and II of the Birds Directive, birds not in Annexes I or II, and all birds grouped by Annex I and II, non-annex and all taxa







Note: The total number of assessments is 505.

Source: Article 12 Member States' reports and EU assessments.

Table 2.1 Bird groups with increasing or decreasing short-term trends in EU Member States (MSs) and their long-term EU trend

Species	Group	Directive status	List of MS with an increasing or decreasing short-term trend	Long-term trend at EU level
Breeding bird populations showing increasing short-term trends in almost 50 % of all MSs				
 Gadwall (<i>Mareca strepera</i>) © Andreas Trepte	Waterfowl	Annex II, part A	BE, CZ, DE, EE, ES, FI, IE, IT, LT, LV, NL, PL, PT, SE, SK, UK	↗
 Greylag Goose (<i>Anser anser</i>) © Public domain	Waterfowl, Marine	Annex II, part A	AT, BE, CZ, DE, DK, FR, HU, IT, LT, LV, NL, PL, SI, SK, UK	↗
 Goosander (<i>Mergus merganser</i>) © Philippe Moret	Waterfowl, Marine	Annex II, part B	AT, BG, CZ, DE, DK, FI, FR, HR, HU, IT, LT, SI, SK, UK	→
 Eurasian Nuthatch (<i>Sitta europaea</i>) © Chme82	Forest		CZ, DE, DK, ES, FI, IT, NL, PL, PT, SE, SI, SK, UK	↗
Breeding bird populations showing decreasing short-term trends in almost 50 % of all MSs				
 Northern Lapwing (<i>Vanellus vanellus</i>) © Charles J Sharp	Farmland, Waterfowl, Marine	Annex II, part B	BE, DE, DK, HU, IE, LT, LU, NL, PL, SE, SI, SK, UK	↘
 Grey Partridge (<i>Perdix perdix</i> , except subspecies <i>hispaniensis</i> and <i>italica</i>) © Roger Culos	Farmland	Annex II, part A	AT, BE, DE, EE, FR, HU, IT, LU, NL, PL, SI, SK, UK	↘
 Red-backed Shrike (<i>Lanius collurio</i>) © Charles J Sharp	Farmland	Annex I	BG, CZ, EE, ES, FI, FR, GR, IT, LT, LU, LV, SI, SK	?
 Corncrake (<i>Crex crex</i>) © Ron Knight	Waterfowl	Annex I	BE, BG, DE, EE, FR, HR, IT, LT, LU, LV, NL, PL, SK	→
Wintering bird populations showing increasing short-term trends in 10 or more MSs				
 Great White Egret (<i>Ardea alba</i>) © JJ Harrison	Waterfowl, Marine	Annex I	BE, BG, CZ, DE, ES, FR, GR, LU, NL, PL, PT, SK	↗
 Whooper Swan (<i>Cygnus cygnus</i>) © Thermos	Waterfowl, Marine	Annex I	BG, DE, DK, FI, FR, GR, IE, PL, SE, UK	↗
 White-tailed Eagle (<i>Haliaeetus albicilla</i>) © Yathin S Krishnappa	Marine	Annex I	AT, CZ, FI, FR, HU, LT, NL, PL, SE, SK	↗
 Northern Shoveler (<i>Spatula clypeata</i>) © Rudolphous	Waterfowl, Marine	Annex II, part A	AT, BE, CY, DE, ES, FR, NL, PT, SI, UK	↘

Wintering bird populations showing decreasing short-term trends in 10 or more MSs					
	Mallard (<i>Anas platyrhynchos</i>) © H. Zell	Waterfowl, Marine	Annex II, part A	AT, BE, CZ, DE, FR, IE, IT, MT, NL, PT, SI, UK	→
	Common Coot (<i>Fulica atra</i>) © Philippe Amelant	Waterfowl, Marine	Annex II, part A	CY, DE, DK, FI, HU, IE, IT, SI, SK, UK	↘
	Common Goldeneye (<i>Bucephala clangula</i>) © Maga-chan	Waterfowl, Marine	Annex II, part B	AT, BE, DE, FR, IE, IT, LU, NL, SK, UK	→
	Red-breasted Merganser (<i>Mergus serrator</i>) © Andreas Trepte	Waterfowl, Marine	Annex II, part B	BE, BG, CY, ES, FR, IE, MT, SE, SI, UK,	→

Notes: Waterfowl: African-Eurasian Waterbird Agreement; Marine: Marine Strategy Framework Directive; Forest: Common Forest Bird Indicator; Farmland: Common Farmland Bird Indicator.

↗ Increasing ↘ Decreasing → Stable ? Unknown

Key messages

- Population status was assessed for a total of 463 bird species: 47 % have a good population status and 39 % have a poor or bad status.
- 23 % of breeding birds have increasing short-term trends and 30 % have decreasing short-term trends; there are more decreasing than increasing long-term trends.
- Short-term trends for wintering birds are consistent with the results for the years 2008-2012, but the proportion of taxa with increasing long-term trends has dropped by 9 %.
- The proportion of taxa listed in Annex I with good status has decreased by 8 % since the last reporting period, while taxa with poor and bad status have increased by 6 %.
- The proportion of Annex II species with a good status is 9 % lower than in the 2008-2012 reporting period, while taxa with poor and bad status are 9 % higher.
- A good status for at least 50 % of the assessed species is reported for half of the taxonomic orders, but a high proportion of the remaining taxa within these orders are still assessed as 'bad'.
- The high percentage of unknown data suggests a need to establish or re-enforce appropriate coordinated and state-supported monitoring schemes within and between all Member States.



DASHBOARD
with detailed information by Member State:
breeding birds



DASHBOARD
with detailed information by Member State:
wintering birds

2.3 Population status of species with species action plans

To achieve the conditions set out in Annex I of the Birds Directive, species action plans (SAPs) are sometimes used at national or international level to help organise and maximise the effectiveness of conservation efforts within a country or across several countries. SAPs are, however, not just developed for Annex I species. Annex II species may also have management plans to help guide the balanced, sustainable and ecologically sound hunting of a species.

In total, 75 bird species and subspecies have an international SAP. These plans were either produced by the EU itself or were produced under international agreements (such as the Agreement on the Conservation of African-Eurasian Migratory Waterbirds or the Bonn Convention for Migratory Species) and endorsed by the EU. Of all bird species with a SAP, 63 are listed in Annex I, and 11 are listed in Annex II (two are listed in both annexes). Figure 2.13 provides an overview of the EU population status of species with and without SAPs.

For taxa with a SAP, the following observations can be made:


- A higher proportion of taxa with a SAP have a bad status (52 %) than those without a SAP (14 %). This is not surprising, as SAPs target species that were assessed as having a bad status (threatened) at the time of their development.
- Only 23 % of species with a SAP have a good status compared with 50 % of species without a SAP.

Box 2.3 provides two examples of birds for which a SAP has been developed but which have different statuses and trends.

In total, 13 bird taxa have an EU management plan. Management plans were produced for huntable species (Annex II) considered to have a bad status in the EU. Almost half of the taxa with management plans have a bad status, and almost another quarter have a poor status. This means that over two thirds of the taxa that have management plans are in a poor or bad status category (as shown in Figure 2.14).

Several management plans were prepared between 1997 and 2000, and all of the taxa with plans that have been assessed as threatened (a total of six) have decreasing short-term trends. This leads to the possible conclusion that the plans have not delivered the expected conservation results.


Figure 2.13 EU population status of taxa for which a SAP has been developed versus those taxa which do not have a SAP



Note: The total number of taxa with SAPs is 75 and of taxa without SAPs is 430.

Source: Article 12 Member States' reports and EU assessments.

Figure 2.14 Status of taxa with and without a management plan



Note: The total number of taxa with a management plan is 13 and without a plan is 498.

Source: Article 12 Member States' reports and EU assessments.

Box 2.3 Closer look: Birds with species action plans

Photo: Lesser White-fronted Goose © Vasily Sokolov, IUCN Red List

The **Lesser White-fronted Goose** (*Anser erythropus*) is a goose species breeding in the tundra zone from northern Fennoscandia to eastern Siberia. While migrating, the species prefers low-growing salt marshes/coastal meadows and natural steppes but, because of the transformation of its original staging and wintering habitats in the southern parts of the range, the main feeding habitat in many areas today is agricultural land. In 2008, an International Species Action Plan for the Conservation of the Lesser White-fronted Goose ^(*) was adopted. Moreover, LIFE projects such as Anser-Eur (LIFE05 NAT/FIN/000105) and Safeguard LWfG (LIFE10 NAT/ GR/000638) targeted the species with the objective of improving and monitoring the conservation status of the species at the most important breeding, staging and wintering sites along the European flyway. This

was to be quantified by halting or slowing down the negative trend in the population size at the monitoring sites, increased survival during the wintering period, an increase in the time that the geese spend in safe, managed project sites, and increased awareness of hunters, farmers and landowners of the species and its conservation. The projects have improved cooperation between five countries along the species' flyway (Estonia, Finland, Greece, Hungary and Norway), including the preparation of coordinated national species action plans (approved by the end of the project in Estonia, Finland and Norway) and a joint contribution to the new International Single Species Action Plan. This coordinated effort has led to a positive trend in the breeding populations in Finland.

The **waders of wet grassland habitats** include eight species that are covered by an international multi-species action plan (MSAP). Five of these species have a bad EU status: Baltic Dunlin, Eurasian Oystercatcher, Black-tailed Godwit, Common Redshank and Northern Lapwing. Three have a poor EU status: Ruff, Eurasian Curlew and Common Snipe. The European populations of all eight species have declined considerably in recent decades, and these declines are still ongoing, except for the Eurasian curlew whose short-term trend is now stable. In fact, between 60 % and 90 % of Member States have reported that the species are deteriorating despite the existence of the MSAP. Their status is largely linked to the quality and extent of sympathetically managed wet grassland habitats and the wider landscapes within which they are found. In the past, ideal conditions for these species were intrinsically created and maintained by more traditional farming practices. However, the policies and incentives of pillar 1 of the common agricultural policy have primarily focused on increasing agricultural production. In fact, the population declines of wet grassland breeding waders have been more pronounced within the EU than in non-Member States. While the MSAP (2018) partially targets the restoration of wet grasslands, it has not been in place for long enough to have an effect on these populations, as it takes time for the conservation measures to be implemented and yield positive results. Efforts thus need to be put in place to implement the conservation measures identified in the MSAP in the future.

Note: ^(*) <https://www.unep-aewa.org/en/publication/international-single-species-action-plan-conservation-lesser-white-fronted-geese-western>

Key messages

- 75 bird species and subspecies have an international species action plan (63 from Annex I and 11 from Annex II).
- Only 23 % of species with a species action plan have a good status compared with 50 % of species without an action plan.
- 13 bird taxa have an EU management plan; over two thirds of these have a poor or bad status.



DASHBOARD
with detailed information by Member State:
International SAPs

2.4 Improvement in trends

Within Article 12 reports, Member States provide information on the population size and breeding distribution of birds and their trends. In around 15 % of all Member States' reports, both short- and long-term population trends are increasing. Bird taxa identified in this chapter as having an improving population trend are based on Member States' Article 12 reports, which apply the same method used to identify taxa with non-secure improved status under the evaluation of progress towards target 1 of the EU Biodiversity Strategy to 2020 (see Chapter 6). Bird reports were thus selected which meet one of the following two criteria:

1. showed an increasing population trend in the last 12 years (2007-2018), regardless of the 38-year long-term trend (1980-2018); or
2. showed a stable or fluctuating short-term population trend in the last 12 years, after a decreasing long-term trend.

This group of species will be referred to as '**improving bird taxa**' in the following section. In total, there are 2 148 Member States' reports recording the

improving state of populations according to the criteria above; globally this corresponds to 397 bird taxa for which there is an improvement in at least one Member State and for at least one of the reported seasons. Under the first criterion fall 1 715 reports, or 367 taxa. The second criterion is met by 433 reports, or 214 taxa.

Overall, approximately 49 % of the reported improvements relate to Annex I and SPA trigger species (see Table 2.2).

The majority of all taxa that show improving trends are recorded for more than one Member State. Within the group of Annex I and SPA trigger species, the following species are the most frequently mentioned across Member States: White-tailed Eagle (*Haliaeetus albicilla*) ($n = 30$), Great White Egret (*Ardea alba*) ($n = 25$), Crane (*Grus grus*) ($n = 24$), Whooper Swan (*Cygnus Cygnus*) ($n = 23$) and Red Kite (*Milvus milvus*) ($n = 21$).

Figure 2.15 presents the number of Member States' reports showing improving trends for each species group. In addition, the proportion of each of the improvement categories is compared with the overall number of Member State reports in each

Table 2.2 Bird taxa reports showing improving trends

Improvements in the reporting	Reports of improving bird taxa	Reports of improving bird taxa listed as Annex I or SPA trigger species
Increasing → Increasing	1 213	685
Stable → Increasing	146	38
Fluctuating → Increasing	15	10
Decreasing → Increasing	123	50
Unknown → Increasing	15	9
Unknown → Increasing	196	65
No data → Increasing	7	5
Decreasing → Stable	379	147
Decreasing → Fluctuating	54	42
Total	2 148	1 051

Note: Population trend long (1980-2018) → Population trend short (2007-2018).

Source: Article 12 Member States' reports and EU assessments.

species group. These results reveal that the share of improving trends is very similar across species groups, averaging 26 %. However, uncertainties remain as many bird species are not categorised.

Around 44 % of all bird species with improving trends are covered by the Agreement on the Conservation of African-Eurasian Migratory Waterbirds (AEWA ⁽¹⁾). Many of these species are also categorised as marine species according to the EU Marine Strategy Framework Directive (EU, 2008) and make up 33 % of all improvements. There are also some farmland and forest bird species that show improving trends, but these comprise only around 9 % of all improving trends. To interpret these percentages it has to be taken into account that approximately 60 % of the improving wetland bird species are also classified as marine birds. Overlaps also exist between some farmland/forest birds and wetland birds.


The shares of improvements for SPA and Annex I marine and wetlands bird species are higher but lower than those for farmland and forest bird species compared with other (non-Annex I and SPA) improving bird species. This may suggest that either the SPA coverage for forest and farmland birds is currently insufficient or the habitats within SPAs are not sufficiently improved for these species. Table 2.3 lists examples of Annex I/SPA trigger

bird species showing improving trends within each taxonomic group.

Around 78 % of Member States' reports show improving population trends attributed to breeding birds, 18 % to wintering taxa and 4 % to passage taxa. Looking at Annex I and SPA trigger species, their share of improvements is highest for passage birds (98 %), followed by wintering birds (73 %) and breeding birds (approximately 40 %).

Improvements can be driven by various actions and factors. Improvements resulting from conservation measures, such as habitat protection or restoration and research, monitoring and awareness-raising activities (see Box 2.4), lead to real positive changes in species' populations at Member State level and are considered genuine. Other improvements can be caused by changes in methodology, better knowledge, natural population dynamics (e.g. typical for wintering birds) or large-scale landscape changes. For example, agricultural intensification may create artificial feeding habitats for certain migrating and/or wintering birds (e.g. Greylag Goose and Barnacle Goose) in parts of their range, causing larger populations or improving trends to be observed in some Member States (and potentially decreases elsewhere).

Figure 2.15 Number of Member States' reports showing improving trends by species group



Note: Percentage of all reports in the groups; taxa could be listed under more than one category. Total number of reports showing improvement: 2 148. Number of Annex I and SPA trigger species showing improvement: 1 051.

Source: Article 12 Member States' reports and assessments.

⁽¹⁾ <https://www.unep-aewa.org>

Table 2.3 Examples of Annex I/SPA trigger taxa showing improving trends by taxonomic order

Taxonomic group	Total reports (n)	Example		
		Species name	MS	Changes in population trend between long and short period
Passeriformes	711	Red-backed Shrike (<i>Lanius collurio</i>)	UK	Dec → Inc
Anseriformes	390	Ruddy Shelduck (<i>Tadorna ferruginea</i>)	BG	Dec → Inc
Charadriiformes	302	Black Guillemot (<i>Cepphus grille</i>)	SE	Dec → Inc
Accipitriformes	176	Red Kite (<i>Milvus milvus</i>)	ES	Dec → Inc
Pelecaniformes	111	Bittern (<i>Botaurus stellaris</i>)	BE	Dec → Inc
Gruiformes	63	Common Coot (<i>Fulica atra</i>)	PL	Dec → Inc
Columbiformes	57	Stock Dove (<i>Columba oenas</i>)	PL	Unk → Inc
Piciformes	51	Wryneck (<i>Jynx torquilla</i>)	NL	Dec → Inc
Strigiformes	38	Great Grey Owl (<i>Strix nebulosa</i>)	FI	Sta → Inc
Suliformes	37	Shag (<i>Phalacrocorax aristotelis aristotelis</i>)	IE	Sta → Inc
Falconiformes	36	Red-footed Falcon (<i>Falco vespertinus</i>)	HU	Dec → Inc
Podicipediformes	34	Great Crested Grebe (<i>Podiceps cristatus</i>)	PL	Unk → Inc
Galliformes	26	Capercaillie (<i>Tetrao urogallus</i>)	PL	Dec → Inc
Coraciiformes	24	Kingfisher (<i>Alcedo atthis</i>)	AT	Unk → Inc
Caprimulgiformes	22	Nightjar (<i>Caprimulgus europaeus</i>)	LT	Sta → Inc
Ciconiiformes	22	Black Stork (<i>Ciconia nigra</i>)	GR	Unk → Inc
Gaviiformes	12	Red-throated Diver (<i>Gavia stellate</i>)	DE	Unk → Inc
Otidiformes	12	Little Bustard (<i>Tetrax tetrax</i>)	FR	Dec → Inc
Phoenicopteriformes	7	Greater Flamingo (<i>Phoenicopterus roseus</i>)	ES	Inc → Inc
Procellariiformes	7	Storm-petrel (<i>Hydrobates pelagicus</i>)	FR	Unk → Inc
Bucerotiformes	5	Hoopoe (<i>Upupa epops</i>)	ES	Sta → Inc
Cuculiformes	4	Cuckoo (<i>Cuculus canorus</i>)	NL	Dec → Inc
Pterocliiformes	1	Pin-tailed Sandgrouse (<i>Pterocles alchata</i>)	FR	Dec → Sta

Note: Population trend (long → short): Dec, decreasing; Flu, fluctuating; Inc, increasing; Sta, stable; Unc, uncertain; Unk, unknown; MS, Member State.

Source: Article 12 Member States' reports and assessments.

Box 2.4 Closer Look: Recovery of the Eurasian Spoonbill in France

Photo: Eurasian Spoonbill © Martin Mecnarowski, IUCN Red List

The Eurasian Spoonbill (*Platalea leucorodia leucorodia*) disappeared from France as a breeding bird in the 16th century but started reappearing sporadically in the country between 1973 and 1981 and regularly in the Grand-Lieu Lake (Loire Atlantique). In the mid-20th century, the species was once again endangered because of pollution, hunting and loss of breeding sites as a result of agricultural and hydroelectric developments. The aquatic ecosystem has been disturbed by the draining of feeding areas and transforming them into intensive agricultural zones as well as by water contamination from agricultural activities.

Since the 1990s, national and regional governments have been implementing conservation measures for the Eurasian Spoonbill. Recovery has been facilitated by

the development of a major international conservation plan, including measures such as targeted control of water levels, protecting flood plains and monitoring the development of vegetation and siltation. In France, conservation measures have included wetland protection, restoration and management, which has been supported by research, monitoring and awareness-raising activities.

Of particular importance has been the Grand Lieu LIFE project (LIFE NAT/F/000841), which is implemented at a key site for the spoonbill. Specific measures in the project included acquiring adjacent land to carry out restoration operations such as desilting and dredging to remove a silt bank that had built up at the lake's outlet. The project also included a series of scientific studies to monitor multiple parameters (oxygen levels, sedimentation rates in the lake, suspended sediments and pollutants, and vegetation productivity). To raise public awareness, the former hunting lodge of the Guerlain family was refurbished as a visitor centre hosting educational exhibits on the fauna and flora of the lake. The project has improved the habitat for waterbirds, including the spoonbill, resulting in its breeding population increasing from around 20 to almost 200 pairs. As a result of such diverse measures in France and elsewhere in Europe, the majority of Eurasian spoonbill populations are recovering (especially in north-west Europe, Hungary and the Netherlands), but some populations remain vulnerable.


Source: Tucker et al. (2019).

Key messages

- In around 28 % of all Member States' reports, both short- and long-term population trends are increasing.
- Approximately 50 % of the improvements reported relate to Annex I and SPA trigger species.
- The majority of all bird species with improving trends are wetland and marine birds.
- The share of improvements in SPA trigger and Annex I marine and wetlands bird species are higher than those for farmland and forest bird species.
- Breeding birds have the highest share of improving population trends (approximately 80 %).



Figure 3.1 Summary of the Habitats Directive reporting



3 Status of and trends in habitats and species

The Habitats Directive protects animal and plant species of Community interest, including endangered, vulnerable, rare and endemic species or those requiring particular attention, as well as a number of natural and semi-natural habitat types (called habitats of Community interest), which are in danger of disappearing, have a small range or are outstanding examples of biogeographical regions. The directive outlines requirements for their protection and sustainable use. The 233 protected habitats included in Annex I and over 1 000 species included in Annex II are targeted by designating Sites of Community Importance (SCIs), which are, together with Birds Directive sites, one of the components of the Natura 2000 network. Other species benefit from strict protection (Annex IV) or regulated use (Annex V).

The implementation of several specific measures of the directive (including reporting on the status and trends of habitats and species) is structured along the nine biogeographical regions listed in the Habitats Directive and five additional marine regions based on those from the Marine Strategy Framework Directive (see Map 3.1). Reports from the Member States bring together information on the conservation status of and trends in the species and habitats within each biogeographical and marine region, relevant pressures and threats, conservation measures and the importance of Natura 2000. A separate report is prepared for each region where the habitat or species is present. In addition, Member States report on the directive's implementation (e.g. status of Natura 2000


and its management, information on Annexes IV and V species).

On the basis of Member States' assessments for each habitat and species present in each biogeographical/marine region, separate EU regional assessments are conducted for the biogeographical regions and marine regions as a whole. They were carried out by experts from the EEA and the European Topic Centre on Biological Diversity (ETC/BD) between October 2019 and February 2020, followed by a public consultation in February and March 2020.


This chapter provides the main results from these reports, representing the period from 2013 to 2018. For this report, conservation status (see Box 3.1) and trends at Member State and EU biogeographical level are dealt with jointly in the following sections. In addition to highlighting the need for improvement, examples are provided of species and habitats having exhibited genuine improvements in conservation status or trend and positive trends since the previous reporting period.

To interpret the results further, it has to be noted that they do not express the actual proportion of the habitat area or the proportion of species populations with favourable or unfavourable status. Due to the nature of the data and methodology, the analysis shows the number of reports for each conservation status and does not reflect the shares of habitat area or species population that each Member State actually inherits.

Map 3.1 Biogeographical and marine regions in the EU



Reference data: ©ESRI



Note: For the marine regions, the boundaries are indicative only and do not imply any legal status.

Source: EEA and ETC/BD, May 2020.

Box 3.1 Approach to assessing conservation status

The conservation status of a species or habitat is derived using four parameters. The agreed method for the evaluation of conservation status separately assesses each of the parameters of conservation status, with the aid of an evaluation matrix, which is a part of the report format approved by the Member States, and then combines these assessments to give an overall assessment of conservation status. The aggregation matrix follows the one-out, all-out rule: if one parameter is assessed as bad, the overall conservation status is bad.

Parameters for assessing the conservation status of habitat types	Parameters for assessing the conservation status of species
Range	Range
Area covered by the habitat within its range	Population
Structure and functions, including the status of typical species	Habitat for the species
Future prospects	Future prospects

The status of a habitat is derived by individually assessing four parameters: natural range, area, structure and functions, and future prospects. The status of the range broadly reflects past and present changes in the habitat's natural distribution, while the status of the area reflects past and present declines in its surface area. Structure and functions are based on the condition of specific structural features and habitat functions. Habitat structure is considered to be the physical components of a habitat, which will more than likely be formed by species both living and dead (e.g. dead trees in a forest) but can also include abiotic features. It provides information on the proportion of the habitat area by condition, its trends, and typical species. The condition is reported as a minimum or maximum range in 'good', 'not-good' or 'unknown' condition. This parameter is further used in the restoration analysis (see Section 4.3).

As for habitats, status is derived for species by individually assessing the status for four parameters: natural range, suitable habitat, population, and future prospects. The status of the population reflects past and present changes in population size. Suitable habitat provides information on its quality, availability and trends.

A more detailed description of the methodology behind the Article 17 conservation status assessment can be found in Röschel et al. (2020).

3.1 Facts and figures

This section presents a short overview of the different types of habitats and species included in the Habitats Directive annexes. Annex I of the Habitats Directive comprises 233 different habitat types, 224 of which are terrestrial and nine marine. The reported terrestrial habitats cover almost one third of the terrestrial area of the 28 Member States (EU-28; 1 July 2013 to 31 January 2020), equivalent to 1.3 million km². Marine habitats cover an additional 0.4 million km² of EU waters. Annexes II, IV and V of the Habitats Directive list 1 389 species of European interest ⁽¹²⁾.

3.1.1 Share of different types of species and habitats

The Habitats Directive covers a wide range of habitat types and species. Over one third of the total number of habitat types covered by the directive are forests (Figure 3.2), making up nearly 0.5 million km² in area (Table 3.1). In the case of forests, this makes up approximately 27 % of the total forested area in the EU. Another 14 % of all habitats correspond to natural and semi-natural grasslands, which make up around 0.23 million km². As regards the number of habitat types, temperate heath and scrub, sclerophyllous scrub, bogs, mires and fens and rocky habitats each comprise only about 5 % of the habitats listed. For species, almost half are vascular plants, while molluscs and non-vascular plants represent 3 % each (Figure 3.3).

The Habitats Directive makes a fundamental contribution to the conservation of Europe's biodiversity. The habitat types protected by the directive overlap to a certain degree with more than half of the natural and semi-natural terrestrial, freshwater and marine habitat types in the European Nature Information System (EUNIS) habitat classification. The overlap is at EUNIS class level 3, which is a level broadly corresponding to the directive's Annex I habitat types and to the European Red List of habitats. The Habitats Directive includes about 5 % of the total number of European species referenced in the publications of the International Union for the


Conservation of Nature (IUCN) European Red Lists. The Red Lists rate the extinction risk of subsets of European species. The Habitats Directive's coverage of species is lower for species-rich groups, such as molluscs, arthropods and vascular plants and higher for amphibians (85 %), reptiles (70 %), mammals (64 %) and freshwater fish (39 %). Many species not included in the annexes of the directive benefit indirectly, as do their habitats, from the directive's protection regime.

3.1.2 Spatial distribution of species and habitats covered by the Habitats Directive

Habitats and species covered by the Habitats Directive are not equally distributed across the EU, and some regions have a higher diversity than others (Maps 3.2 and 3.3). A higher diversity of Annex I habitats can be found within mountainous regions and in other sparsely populated areas, e.g. Finland and the diverse Mediterranean southern and south-western Europe. Regions with a low diversity of habitats tend to be areas of more intensive land use. Examples include the lowland parts of the Atlantic region, the artificial plains of large European rivers, such as the Po plain in northern Italy and the Danube basin in eastern and south-eastern Europe, or the flat regions of Spain such as Castile and León.


For species, their occurrence across the EU shows fewer distinct patterns, partly reflecting the fact that the directive protects species with very diverse ranges: from endemic species with restricted ranges to common and more widely distributed species. In addition, the comprehensiveness of inventories and mapping vary among the Member States and some species pose methodological challenges, e.g. mapping of highly mobile and common species. Nevertheless, there is a greater diversity of listed species in mountainous regions and in some Mediterranean countries, such as Greece and Cyprus, and in the Macaronesian region, which has high numbers of endemic species. The diversity of species covered by the Habitats Directive is generally lower in the northern Atlantic and Boreal regions of Europe and higher in central and southern Europe.

⁽¹²⁾ Annex II: species requiring designation of Special Areas of Conservation (Natura 2000). Annex IV: strictly protected species. Annex V: species for which exploitation may require specific regulation and management.

Figure 3.2 Share of habitats protected by the Habitats Directive (Annex I)

Note: Based on total number of habitats covered by the Habitats Directive.

Source: Article 17 Member States' reports and assessments.

Figure 3.3 Share of species protected by the Habitats Directive (Annexes II, IV and V)

Note: Based on total number of species covered by the Habitats Directive.

Source: Article 17 Member States' reports and assessments.


Table 3.1 Area of Annex I habitat types per group (marine and terrestrial)

Habitat group	Average area (km ²)
Bogs, mires and fens	137 700
Coastal habitats	406 000
Dune habitats	9 200
Forests	491 900
Freshwater habitats	127 800
Grasslands	234 300
Heaths and scrub	88 300
Rocky habitats	121 800
Sclerophyllous scrub	35 100
Total Annex I	1 652 200


Note: Surface areas as reported by Member States and rounded to the nearest 100 km². The area reported for Romania is clearly overestimated: the reported area correspond to 127 % of the terrestrial area of this Member State.

Source: Article 17 Member States' reports and assessments.


Map 3.2 Spatial diversity of habitats covered by the Habitats Directive (marine habitats are excluded)




Reference data: ©ESRI



Map 3.3 Spatial diversity of species covered by the Habitats Directive (marine species are excluded)



Reference data: ©ESRI



Source: Prepared by ETC/BD based on the number of reported habitats/species per 10 km × 10 km grid cell: EEA; country boundaries. © OpenStreetMap based on Article 17 Member States' reports and assessments.

3.2 Conservation status

This section presents insights into the conservation status of habitats and species at the EU regional and Member State levels. It is based on the definition of 'favourable conservation status' provided in Article 1 of the Habitats Directive. The 'unfavourable' status is divided into two categories, reflecting its distance from 'favourable'.


In this report, to simplify readability, the species and habitat conservation status categories are labelled 'good' for a favourable conservation status and 'poor' or 'bad' for an unfavourable-inadequate or unfavourable-bad conservation status (see Table 1.1). Because of the nature of data acquisition for each biogeographical and marine region, the conservation status is presented for habitat and species assessments rather than for single habitats and species. The number of 'assessments' refers to the assessment made by a Member State in one biogeographical region. Consequently, a species or habitat type that occurs in more than one biogeographical region within a single Member State can have more than one assessment. More information on the methodology underlying the conservation status assessment can be found in the separate methodological technical report (Röschel et al., 2020).

3.2.1 Habitat assessments

The results of the 818⁽¹³⁾ conservation status assessments of habitats at the EU regional level show that, while 15 % of habitat assessments show a good conservation status, the vast majority show an unfavourable conservation status (45 % poor and 36 % bad) (Figures 3.4 and 3.5). Compared with the last reporting period (2008-2012), bad conservation status for habitats has increased by 6 %. These differences are generally related to changes in the methods applied at the EU or Member State levels or are due to variations in data quality.

Of the habitat groups that were reported on, coastal habitats have the lowest proportion of assessments showing good conservation status. Dune habitats and bogs, mires and fens are most frequently assessed as having a bad conservation status (over 50 %). As one of the most species-rich habitats in Europe, grasslands are also among the habitats with the highest share of assessments showing a bad conservation status (49 %); managed grasslands are in a particularly bad state (see Section 6.2.1). Ranging from almost desert-like grasslands in Spain to humid meadows in central and northern Europe, grasslands are home to a great diversity of insects


Figure 3.4 Conservation status of habitats at EU level



Note: Statistics are based on the number of EU habitat assessments (818).

Source: Article 17 Member States' reports and EU assessments.

Figure 3.5 Conservation status per habitat group at the EU level



Note: The number of assessments per group is indicated in parentheses. The total number of assessments is 818.

Source: Article 17 Member States' reports and EU assessments.

⁽¹³⁾ In total, 828 assessments were conducted for habitats. Because of data inconsistencies, only 818 were used for the statistical assessment.


and other arthropods, including many pollinator species and other grassland-adapted species. An in-depth analysis shows that grasslands (and other habitats) with particular relevance for pollinator species have a higher proportion of bad conservation status compared with the rest of the habitat group (see the analysis in Box 6.2). For grasslands, as well as bogs and mires, the majority of the total habitat area has been lost over the last century due to intense human intervention, such as agricultural intensification, draining peatlands and land take/urbanisation (see Box 4.8 and Section 4.3.1 for more information on the dominant pressures affecting habitats). Rocky habitats (found mostly on mountain slopes and in more remote areas) provide a more uplifting picture, standing out with a comparatively high proportion of habitat assessments that show a good conservation status.

The overall status of a habitat is determined by a combination of status assessments for the parameters (see Box 3.1). A poor or bad overall conservation status is most frequently due to a poor or bad status of the habitat's structure and functions or future prospects. For many habitats, the area or — even more frequently — range have a good status (see Figure 3.6).

Across the EU's biogeographical and marine regions, conservation status assessments vary greatly (see Figure 3.7). Some of the observations are as follows:

- For the terrestrial regions, the Steppic⁽¹⁴⁾ region stands out with 72 % of habitat assessments showing a good conservation status; this region is exclusively confined to Romania, which reported the highest overall proportion of habitats with good status among the Member States (see Figure 3.8). For all other regions, not good (poor and bad) conservation statuses have been reported for more than 60 % of the habitat assessments.
- The Atlantic region has the smallest portion of habitats with good conservation status (4 %), with grasslands, forests and coastal habitat assessments being particularly degraded. It is also the region with the highest share of bad conservation status (over 50 %). Bad conservation status is also high for the Boreal, Pannonian, Continental and Mediterranean regions.
- Marine regions (with only nine habitat types⁽¹⁵⁾) have good statuses reported less frequently


Figure 3.6 Status of parameters for habitats at EU level



Note: The total number of EU assessments is 818.

Source: Article 17 Member States' reports and EU assessments.

Figure 3.7 Conservation status of habitats for each biogeographical and marine region at EU level



Note: Statistics are based on the number of EU habitat assessments. The number of assessments per region is indicated in parentheses. The total number of assessments is 818.

Source: Article 17 Member States' reports and EU assessments.

⁽¹⁴⁾ Following the methods for assessing EU conservation status, the regional status for habitats that occur in only one Member State within a biogeographical region is the same as in that Member State.

⁽¹⁵⁾ Sandbanks (1 110), Posidonia beds (1 120), estuaries (1 130), mudflats and sandflats (1 140), large shallow inlets and bays (1 160), reefs (1 170), submarine structures (1 180), Boreal Baltic narrow inlets (1 650) and sea caves (8 330).

on average than terrestrial regions, with good conservation status reported for only one marine region (i.e. the Marine Black Sea, with 14 %).

- The Marine Baltic and Marine Atlantic regions show a particularly high share of bad status

assessments, with 71 % and 57 %, respectively. However, there are many marine habitats where conservation status was assessed as unknown because of the lack of data from Member States, e.g. for the Marine Macaronesian region, where all assessments were unknown.

Box 3.2 Closer look: Conservation measures to restore dunes and reefs



Photo: Dunes in the Netherlands © NEEMO

Bringing back dynamics in the Dutch dunes

Shifting dunes (2120) are present in most biogeographical regions, but the vast majority of this rare habitat's surface area can be found along the Atlantic coasts. They have the ability to provide a natural buffer against some of the negative effects of climate change but are highly dependent on natural processes such as wind erosion. A key threat to the habitat is human-induced fixation as a coastal defence, realised through the creation of solid dykes. The dunes are also threatened by the planting or encroachment of — often invasive alien — scrub and trees. A number of LIFE projects — especially in the Netherlands — have focused on restoring the natural dynamics, aiming to set back succession to an earlier stage. For example, ambitious measures were taken to create openings at five locations in

a fixed dune row at the Natura 2000 site Zuid-Kennemerland. These openings measured up to 15 m high and 50 m wide. This had extended the surface area of white dunes in the site from 154 ha to 175 ha by the end of the project. Wind patterns now drive mobile dunes, which are gradually 'walking' over the area. This supports many pioneer species that are typical of these habitats but which are endangered in the Netherlands. The project created corridors between white and grey dunes and thereby increased their resilience. Succession will now allow vegetation patches to move around, benefiting all of the target habitats and their typical species. A gradual improvement in this habitat's status has been recorded in the Atlantic region (from bad in 2006 to poor in 2018), with the strongest improvements in the Netherlands (poor in 2012 to good in 2018).


Restoring reefs in Kattegat

Reefs are particularly difficult habitats to restore, as evidenced by the project BLUEREEF (LIFE06 NAT/DK/000159). This LIFE project set out to rebuild offshore cavernous boulder reefs in the Kattegat which had been exploited. The idea was to build sea defences by restoring the structure and function of the cavernous element of the shallow offshore boulder reefs and by stabilising the top of the existing boulder reef. The project restored the target area using natural stones from a quarry in the southern part of Norway at significant cost. The project increased marine life, including the restoration of 6 tonnes of macroalgal vegetation and 3 tonnes of bottom-living fauna, and evoked a three- to six-fold increase in cod in the reef area. The project also increased awareness among environmental managers, policymakers and the broader public of marine nature restoration, conservation and management issues. A 'code of conduct' (BLUEREEF, 2013), containing valuable experiences and recommendations for carrying out restoration projects, was produced to potentially inspire other areas in northern Europe to restore natural stone reefs. For Danish initiatives, the code also gives guidance on obtaining the necessary permission from different marine authorities. Many of its recommendations are also relevant for the restoration of other marine nature types, such as biogenic reefs. This was a significant input at the local scale but, because the categorisation of the habitat type 'reefs' is so broad, this type of project is unlikely to have an impact on the overall conservation status of the habitat. The conservation status of reefs in Denmark's Marine Atlantic region remains bad.

Looking at the spatial distribution of conservation status, Member States' assessments show highly diverse results across the EU territory. Consistent with the analysis of biogeographical and marine regions, Map 3.4 illustrates high levels of variation among regions and a finer pattern inside the biogeographical regions. In line with the overall statistics on biogeographical regions, the areas with prevailing good conservation status are situated within the Alpine region and in a few Member States (mainly Croatia, Greece, Portugal and Romania). The Baltic and northern Atlantic regions, however, exhibit overwhelmingly bad conservation status. The highest concentration of distribution grids with predominantly bad status can be seen in the United Kingdom (strongly affected by atmospheric pollution) and the Marine Baltic region. In general, higher concentrations of distribution grids with predominantly bad status are seen in lowland and flat areas (e.g. Po basin, Danube basin, southern Sweden, Poland). This map also raises some data issues that are important to keep in mind when interpreting the results: the map is based on the conservation status assessed for each Member State's biogeographical/marine region projected into the 10 km × 10 km distribution grid and is affected by the quality of Member State distribution data. Conservation status assessment at Member State level includes a significant degree of expert input, which may lead to different results among Member States. Moreover, the number of habitats assessed varies greatly, as is the case for the United Kingdom and Marine Baltic region because they have less habitat variety than other countries or regions.

Member States' reporting shows significant variations in terms of the conservation status of habitats within their territories (Figure 3.8). With the exception of Romania, Estonia, Greece and Cyprus, Member States report a good conservation status for less than 40 % of their habitat assessments. Belgium and Denmark have, on the other hand, the lowest share of habitats with a good conservation status and — together with the United Kingdom — report more than 70 % of their habitats as having a bad conservation status. Some of the habitats with the poorest conditions across these countries are grassland habitats, such as *Molinia* meadows, as well as bogs, mires and fens, such as calcareous and alkaline fens. Spain and Croatia have the highest shares of unknown assessments among Member States (each exceeding 10 %).


Figure 3.8 Conservation status of habitats at Member State level




Note: Statistics are based on the number of habitat assessments at Member States' biogeographical/marine level. The number of assessments per Member State is indicated in parentheses. The total number of assessments is 3 246.

Source: Article 17 Member States' reports and assessments.

Map 3.4 Spatial distribution of habitats' conservation status at Member State level represented in a 10 km × 10 km grid



Spatial distribution of habitats conservation status at Member State level represented in a 10 x 10 km grid



Note: Details on the methodology available from Röschel et al., 2020.


Source: Article 17 Member States' reports and assessments.

3.2.2 Species assessments

Looking at the 2 825 ⁽¹⁶⁾ EU regional assessments undertaken for animal and plant species covered by the Habitats Directive (see Figure 3.9), more than one quarter (27 %) have a good conservation status. Compared with the previous reporting period, this indicates an increase of 4 % ⁽¹⁷⁾. These differences are typically related to changes in methods applied at EU or Member State level or to variations in data quality. Nonetheless, over 60 % of the assessments report a poor or bad status (42 % and 21 %, respectively). Although the number of regional assessments that are classified as 'unknown' has decreased since the last reporting period (from 17 % to 10 %), it remains significantly higher than for the habitat assessments (4 %). The species with unknown status are often difficult to observe because of their cryptic lifestyle or often need for destructive methods for their inventories (e.g. *Vertigo* snails). There are also knowledge gaps for common or dispersed species requiring extensive surveys to gather information (e.g. many amphibians and reptiles or bats). Marine mammals are also among those species with the highest proportion of unknown assessments (over 78 %). Generally, the lack of appropriate monitoring is an important factor, in particular for cetaceans, for which, apart from two Annex II species, the range, population size and suitable habitat area are unknown in the majority of Member States.

The species groups with the highest proportion of good conservation status are reptiles and vascular plant species, with 36 % and 40 %, respectively (see Figure 3.10). However, a number of species remain in bad conservation status, such as many Macaronesian endemic reptiles, or mammals such as the European Mink (*Mustela lutreola*). More effective conservation efforts are needed, especially for fish and molluscs, which each have around 30 % of species with a bad conservation status.


Figure 3.9 Conservation status of species at EU level



Note: Statistics are based on the number of EU species assessments.

Source: Article 17 Member States' reports and EU assessments.

Figure 3.10 Conservation status per species group at the EU level



Note: The number of assessments per group is indicated in parentheses. The total number of assessments is 2 825.

Source: Article 17 Member States' reports and EU assessments.

⁽¹⁶⁾ In total, 2 925 assessments were conducted for species, but 100 of these assessments corresponding to occasional or marginal occurrences of species within a biogeographical region were excluded.

⁽¹⁷⁾ Further information on the improvement of conservation status for species is provided in Section 3.4.3 in the analysis of genuine changes in conservation status assessments.

Box 3.3 Closer Look: Conservation action for species**Photo:** Inagua project © NEEMO

stand-alone drip irrigation system was installed, leading local seepage water to the places where the species were planted. This has served to rescue the species, while strong stakeholder engagement with regional and local authorities, a local company, cattle breeders, land managers and scientists ensures lasting support and a high chance that the species will survive.

Endemic plant species in the Canary Islands

The Canary Islands are rich in endemic species, i.e. species that only live in that part of the world. Populations of such plants and animals are often small and highly vulnerable. One of these endemic species, *Teline rosmarinifolius*, has experienced a decline from good status in 2006 to poor in 2012 due to overgrazing and devastating forest fires. Thanks to the LIFE project Inagua (LIFE07 NAT/E/000759), a genuine improvement to a favourable status was reported in 2018. A key component of the project was to take the opportunity of the forest fires to control overgrazing by goats and to develop a fire management plan. Some of the plant species were grown in nurseries and then planted and fenced to protect them from grazing. At the same time, a

**Photo:** Allis Shad © NEEMO**Allis Shad in Germany**

Allis Shad (*Alosa alosa*) is an anadromous fish species whose distribution is mainly in the north-east Atlantic and the rivers that drain into it. The distribution range has decreased dramatically. Only 150 years ago, several hundred thousand shad were caught annually in the Rhine system and formed an important part of the local economy in the surrounding communities. Across the EU, its conservation status is bad and mostly still declining. The main reasons for the Allis shad's decline are overfishing, increasing river pollution, destruction of spawning grounds and the construction of river obstacles such as dams and weirs. Two projects in Germany have helped to restore this fish species in the Rhine. The first (Projekt Maifisch, LIFE06 NAT/D/000005) introduced 4.8 million Allis shad larvae into

the Rhine, thereby reintroducing a species that had become extinct in the Rhine around 100 years ago. A follow-up project (*Alosa alosa*, LIFE09 NAT/DE/000008) released a further 6.2 million shad larvae into the river. It was reported that 250 adults were observed returning to the Rhine from the North Sea, which suggests that several thousand adult shad are returning annually to the Rhine system. Despite there being no improvement in conservation status, the genuine improvement in trend is entirely due to the project's activities. Nevertheless, further improvements may rely on actions relating to removing river barriers rather than introducing additional larvae.


The overall status of a species is determined by combining status assessments for the four parameters (see Box 3.1). Future prospects are most likely to impede achieving good status, and 49 % are reported as poor or bad (Figure 3.11). Bad overall conservation status is more frequently due to bad population status than to the status of the suitable habitat, the highest proportion being for fish. Assessments indicate that species range has the highest proportion of good status (over 56 %). Data gaps in Member States' reports remain one of the largest challenges for assessing the status of species, with around one fifth of assessments missing the statuses of population, habitat and future prospects.

The conservation status of species assessments varies among the EU's biogeographical and marine regions (see Figure 3.12).

- In contrast to the results for habitats, the majority of terrestrial species assessments show higher shares of good than bad conservation status.
- The two terrestrial regions with the lowest share of good conservation status assessments are the same as for habitats (Atlantic and Continental, both with around 20 %).
- All marine regions, except the Marine Baltic region, have high percentages of unknown assessments, reflecting the general lack of marine population data (see examples in Table 3.3); the Marine Baltic region has over 75 % of its species assessments in a bad conservation status; however, this only concerns seven species.

Member States' assessments show highly diverse results in conservation statuses across the EU territory (see Map 3.5). The map reveals that areas with predominantly bad status are very scarce and rather localised. Overall, the proportion of areas with largely good conservation status is much higher than that for habitats, while the proportion of bad status is much lower. Poor conservation status is primarily found in central and western European countries. In general, the widely distributed species (occurring in more than 10 000 grids in a Member State's region) are more likely to have good status (45 %), while widely distributed habitats more frequently have poor or bad status (80 %). The limitations arising because of data quality are discussed above (see Map 3.5).


Figure 3.11 Status of parameters for species at EU level



Note: The total number of EU assessments is 2 825.

Source: Article 17 Member States' reports and EU assessments.


Figure 3.12 Conservation status of species for each biogeographical and marine region at EU level




Note: Statistics are based on number of EU species assessments. The number of assessments per region is indicated in parentheses. The total number of assessments is 2 825.

Source: Article 17 Member States' reports and EU assessments.

Map 3.5 Spatial distribution of species' conservation status at Member State level presented in a 10 km × 10 km grid



Spatial distribution of species conservation status at Member State level represented in a 10 x 10 km grid




Notes: Details on the methodology available from Röschel et al., 2020. In some exceptional cases, such as widely ranging but poorly known cetaceans, Member States submitted maps based on a 50 km × 50 km grid.

Source: Article 17 Member States' reports and assessments.

Looking at individual Member State assessments (Figure 3.13), the biggest proportion of species assessments showing a good conservation status were reported by Cyprus, Ireland, Estonia and Malta (over 50 %). The Common Pipistrelle (*Pipistrellus pipistrellus*), for example, is reported as having a good status in all four countries. In seven Member States, the species assessments showing a bad conservation status exceed 30 % (i.e. Austria, Luxembourg, Netherlands, Sweden, Denmark, Belgium and Germany). Generally, Member States reported a higher proportion of unknown assessments for species than for habitats, with the newest EU Member State, Croatia, having the biggest knowledge gaps (unknown conservation status for 47 % of the assessments). Data on marine mammals are particularly lacking.

Figure 3.13 Conservation status of species at Member State level



Note: Statistics are based on the number of species assessments. The number of assessments per Member State is indicated in parentheses. The total number of assessments is 7 589.

Source: Article 17 Member States' reports and assessments.

Box 3.4 Closer Look: Status of the Atlantic biogeographical region

Photo: French Atlantic coast © Pixabay

The Atlantic region includes over half of Europe's coastline. Altogether, 117 habitat types and 52 plant and 81 animal species listed in the Habitats Directive are found in the Atlantic region. It has specific, regional features: low, flat land, a very varied and dynamic coastline rich in habitats and species and an oceanic climate.

The Atlantic region is one of the most heavily populated (by almost one third of the EU population) and intensely managed areas in Europe, putting massive pressure on the natural environment. Therefore, many natural and semi-natural habitats only exist as isolated, fragmented patches in a largely artificial landscape. This badly affects the conservation status of habitats and species in this region.

Around half of the coastal habitats in the directive are present in this region, as are 17 of the 21 coastal and inland dunes. Specifically, the wide range of sand dunes is one of the typical characteristics of this region that also faces particular distress: shifting 'white dunes' (2120), for example, have a poor or bad conservation status in eight of the nine countries with this habitat. The sandy coastal grasslands called machair (21A0) are one of the few habitats restricted to the Atlantic region and currently exhibit a good (United Kingdom) and a poor (Ireland) conservation status. Endemic plant species that rely on these habitats, such as the Shore Dock (*Rumex rupestris*) are consistently reported with a poor conservation status.

Further inland, much of the conservation interest lies in those habitats that were originally formed by low-key management practices, such as the natural and semi-natural grasslands. Overall, grasslands cover around 30 % of the Atlantic region, and, although most have been transformed or impoverished by intensive agriculture, important vestiges of species-rich habitats remain. Seventeen different types listed in the Habitats Directive are found here, including various forms of calcareous grasslands. The status of these habitats is nonetheless critical: seven of the eight assessments on calcareous grasslands (6110 and 6120) report a bad conservation status.

The first priority for conservation is to protect the remaining areas from further development. However, this requires determined efforts across the broader countryside to help reconnect the otherwise isolated habitats.

Sources: Article 17 reports and assessments, and European Commission (2020a).

Key messages

- Only 15 % of **habitat** assessments at EU level show a good conservation status, while the majority continues to show poor (45 %) or bad (36 %) status
- Dune habitats and bogs, mires and fens are most frequently identified as having a bad conservation status (over 50 %).
- Around one quarter of **species** have a good conservation status at EU level. However, over 60 % of the assessments report a poor or bad status.
- Reptiles and vascular plants are the species with the highest proportion of good conservation status (more than 35 %), while fish have the highest proportion of bad conservation status (38 %).
- Marine mammals (cetaceans) are among the species with the highest proportion of unknown assessments (over 78 %).
- The Atlantic and Continental regions contain highest share of poor and bad conservation statuses among all biogeographical regions for both habitats and species.
- A high percentage of unknown data indicates a need to establish or reinforce appropriate and ideally coordinated and state-supported monitoring schemes in all Member States.



DASHBOARD
with detailed information by Member State:
conservation status and trends



3.3 Trends in conservation status

Looking at trends is an essential part of conservation status assessment, as they inform and enable reflection on how statuses are evolving within the reporting period. Reported trends are identified as improving, stable or deteriorating. The category 'unknown' includes both assessments of unknown conservation status and assessments of unfavourable status with an unknown trend; habitats and species with a favourable status are not included in this analysis, although they were reported by Member States. More information on the methodology can be found in the separate methodological technical report (Röschel et al., 2020). The conservation status trends are also available for assessments with a good status. Assessments showing a good conservation status mainly exhibit stable and improving trends; this is the case for 87 % and 12 % of the assessments for habitats and 77 % and 15 % for species at an EU regional level. The trend analysis below thus only focuses on the habitats and species with unfavourable status (poor or bad conservation status). This allows the recognition of subtle changes in both categories and guides the analysis of where more conservation efforts need to be taken.

The following section presents the results for the trends in conservation status of habitats and species at the EU regional and Member State regional levels.


3.3.1 Habitat trends

As presented in Section 3.2.1, 81 % of the habitat assessments show an unfavourable conservation status (poor or bad). The overall results from the EU regional habitat assessment show that the proportion of improvement across these assessments is quite low. Only 9 % show improving trends, while 36 % continue to deteriorate at the EU scale. The share of unknowns remains high (over 20 %) (see Figure 3.14).

Trends in conservation status differ among the various groups of habitats (see Figure 3.15):

- Deteriorating trends are observed for at least 25 % of all assessments across habitat groups, except for rocky habitats (15 %).
- Forest habitats exhibit the highest proportion of improving trends among the assessments (13 %).


Figure 3.14 Conservation status trends for habitats not in good status at EU level



Note: Conservation status trends are based on EU habitat assessments.

Source: Article 17 Member States' reports and EU assessments.

Figure 3.15 Conservation status trends for habitats not in good status per habitat group at EU level



Note: Conservation status trends are based on EU habitat assessments. The number of assessments is indicated in parentheses. The total number of assessments is 698.

Source: Article 17 Member States' reports and assessments.

Box 3.5 Closer look: Increase in Macaronesian laurel forests in the Azores

Photo: Laurel forest on São Miguel island © Neemo

In the east of São Miguel island (Azores), the survival of the rare endemic Azores Bullfinch (*Pyrrhula murina*) depends on the existence and quality of the Macaronesian laurel forest (*Laurus*, Ocotea 9360). The seeds, flower buds and fleshy fruit of the once-thriving laurel forests provide food for the critically endangered bird, of which there were just 100 remaining pairs in 2003. However, invading alien plant species brought to the archipelago by colonisers are threatening the laurel forests and creating a shortage of food for the birds.

Three consecutive LIFE projects have taken on the main task of saving the forest and the species for future generations — and it seems that they have succeeded.

The conservation status of Macaronesian laurel forests improved from poor in 2012 to good in 2018 and stabilised at the favourable level. The Azores bullfinch population stabilised at between 627 and 1 996 specimens and increased in area up to 160 km². This was accomplished by, inter alia, enlarging the Pico da Vara/Ribeira do Guilherme Special Protection Area by almost three times, covering the whole species range, and removing invasive species, such as the exotic *Cryptomeria* and *Hedychium* stands, replacing them with more than 300 000 saplings of diverse native plants cultivated in local nurseries, and creating a special nursery dedicated to the production of native plants (PRIOLO LIV03 NAT/P/000013, LAURISSILVA SUSTENTAVEL LIFE07 NAT/P/000630, Life Terras do Priolo LIFE12 NAT/PT/000527).

Bogs, mires and fens, grasslands and dune habitats have the highest proportion of deteriorating trends, each with over 50 %.

- Among grasslands, mainly hay meadows ⁽¹⁸⁾, Molinia meadows ⁽¹⁹⁾ and several types of semi-natural dry grasslands ⁽²⁰⁾ show a deteriorating conservation status trend. The main reasons for these trends are their dependency on particular sustainable management measures (see Section 4.2 for more detail).

Trends in the conservation status of habitats are spatially distinct among EU countries and regions. Most regions show predominantly stable or deteriorating trends, especially in northern Italy, Germany, Ireland and Sweden as well as coastal parts of Portugal (see Map 3.6). The predominance of unknown trends in marine regions in the Atlantic is also noticeable. The map illustrates the trends

as percentages of trend distributions within a 10 km × 10 km grid.


Looking at the habitat assessments at individual Member State level (Figure 3.16), five countries report an improving trend in status for over 20 % of their habitats (Greece, Netherlands, Belgium, Bulgaria and United Kingdom). Bog woodlands, for example, show positive trends in conservation status in three of these Member States. Moreover, Belgium, Bulgaria and Greece have an improving trend that outweighs their respective deteriorating trends in the overall assessment. While seven countries did not report any improvement for habitats with unfavourable status, deteriorating trends were reported by all Member States. Portugal, Hungary, Germany, Ireland and Slovenia reported a deteriorating trend for more than half of their habitat assessments. Overall, the degree of unknown trends in conservation status varies greatly among the Member States, ranging from 0 % (in Hungary) to over 80 % (in Lithuania).

⁽¹⁸⁾ Habitat types of this group include lowland hay meadows (6510) and mountain hay meadows (6520).


⁽¹⁹⁾ Molinia meadows on calcareous, peaty or clayey-silt-laden soils (*Molinia caerulea*) (6410).

⁽²⁰⁾ Habitat types of this group include semi-natural dry grasslands and scrubland facies on calcareous substrates (*Festuco-Brometalia*) (6210) and Nordic alvar and precambrian calcareous flatrocks (6280).

Map 3.6 Spatial distribution of habitat conservation status trends at Member State level in a 10 km × 10 km grid.




Spatial distribution of species conservation status trends at Member State level represented in a 10 x 10 km grid



Notes: Details on the methodology available from Röschel et al., 2020. The map shows the percentage of reports with improving (+), stable (=) and deteriorating (-) trends for each 10 km × 10 km grid cell. Reports with unknown trends are not included in the triangle; thus, grid cells with 100 % unknown trends appear in blue.

Source: Article 17 Member States' reports and assessments.

Figure 3.16 Conservation status trends for habitats not in good status at Member State level



Note: Conservation status trends are based on habitat assessments showing an unfavourable or unknown status. The number of assessments per Member State is indicated in parentheses. The total number of assessments is 2 468.

Source: Article 17 Member States' reports and assessments.

Box 3.6 Closer look: Recovery of the Loggerhead Turtle




Photo: Loggerhead Turtle © Howard Hall, IUCN Red List

The Loggerhead Turtle (*Caretta caretta*) occurs primarily within the Mediterranean Sea as well as in the north-east Atlantic along the coast of Spain, Portugal, the Netherlands, and the United Kingdom. They are threatened by fishing (bycatch), destruction of their nesting grounds and water pollution. To improve the status of the species, the project LIFE+Migrate (LIFE11 NAT/MT/001070) aimed to improve the state of knowledge in terms of the species' population status and to identify important areas for feeding grounds or migratory routes. To achieve this aim, the project designated three protected areas (Sites of Community Importance) in Malta and assisted in closing important knowledge gaps by establishing the conservation status of the turtle populations. The project 'Reduction of mortality of *Caretta caretta* in the Greek seas'

(LIFE02 NAT/GR/008500), which was awarded best LIFE project in 2008, upgraded rescue facilities for wounded turtles, established first aid stations, and used a combination of modelling available oceanographic data and direct boat-based observations to monitor the released turtles.

Box 3.7 Trends in species population sizes and habitat surface area


The population size of species and the surface area of habitats are essential parameters in assessing overall conservation status. The following analysis highlights the key results for the trends reported by Member States for these two frequently used parameters.

Figure 3.17 Population size trends for species at Member State (MS) level

Note: The total number of species assessments is 5 222.

Source: Article 17 Member States' reports and assessments.

Population size trends are predominantly unknown for most of the species groups, averaging around 40 % of assessments (Figure 3.17). More than half of the trends are unknown for invertebrates other than arthropods (66 %) and for mammals (52 %). Information is scarce predominantly for small mammals, such as the Forest Dormouse (*Dryomys nitedula*) and Hazel Dormouse (*Muscardinus avellanarius*), as well as for marine mammals. Although this prevents us from drawing conclusions, the results show a comparatively high proportion of increasing population trends for mammals. Bat populations, the Eurasian Beaver (*Castor fibre*) and large mammals such as the Grey Wolf (*Canis lupus*) are among the main beneficiaries. For amphibians, fish and molluscs, further decreasing population sizes are reported for around 30 % of species groups.

Figure 3.18 Area coverage trends for habitats at Member State (MS) level

Note: The total number of habitat assessments is 2 472.

Source: Article 17 Member States' reports and assessments.

Most of the habitats show stable area trends (average 50 %), while an increase in area was reported for comparatively few (average 7 %) (Figure 3.18). The highest shares of increasing trends in coverage were identified for freshwater habitats, such as for alpine rivers (3220, 3230, 3240), although the differences between most habitat groups are quite small. Overall, decreasing trends in habitat area are reported for an average of 30 % of the habitats. For grasslands and dune habitats, however, the decreasing trend is significantly higher (45 % and 38 %, respectively). Grassland areas such as semi-natural dry grasslands and scrubland facies on calcareous substrates (6210) and lowland hay meadows (6510) are particularly affected across countries and biogeographical regions.

3.3.2 Species trends

At the EU level (see Figure 3.19), 35 % of the 2 049 species assessments with an unfavourable or unknown status indicate a deteriorating trend. Only 6 % show an improving trend in conservation status. The trend is unknown for an additional 31 % of species.


As stated above, a rather low number of species show an improving trend at unfavourable conservation status. However, there are some differences between species groups (see Figure 3.20).

- Except for mammals, fish and non-vascular plants (10 %, 9 % and 6 %), improvements in unfavourable conservation status are below 5 %.
- While fish have higher improving trends than other species groups, they also have — together with amphibians — the highest proportion of deteriorating trends (close to 50 %).

- Deteriorating conservation trends are also reported for species in other groups, e.g. grassland habitat specialists such as the Marsh Fritillary (*Euphydryas aurinia*) and the lower plant *Jurinea cyanooides*.

Table 3.2 presents the conservation status of and the trend in one species characteristic of each biogeographical and marine region. Contrasting the conservation status and trends for single species at the Member State level and at the EU level also highlights the fact that the population size of species can vary greatly among countries. Those with a large population size therefore have a large influence on the status of and trends in species at the EU level and have a special responsibility. France, for example, reported on nearly 90 % of the Atlantic population of the Marsh Fritillary, which led to a decreasing EU trend, despite its stable or improving trend in the remaining countries.


Figure 3.19 Conservation status trends of species not in good status at EU level



Note: Conservation status trends are based on EU species assessments.

Source: Article 17 reports and assessments.
























































Figure 3.20 Conservation status trends of species not in good status at EU level



Note: Conservation status trends are based on EU species assessments. The number of assessments is indicated in parentheses. The total number of assessments is 2 049.

Source: Article 17 reports and assessments.

Table 3.2 Examples of species' conservation status and trends for each biogeographical and marine region

Species	Member State	Conservation status	Conservation status trend	Conservation trend at EU level
Atlantic				
	France		↘	↘
	Germany		↗	
	Ireland		↗	
	Portugal		?	
	Spain		→	
	United Kingdom		→	
Alpine				
	France		↘	↘
	Austria		↘	
	Italy		→	
Boreal				
	Finland		?	?
	Sweden		↘	
Black Sea				
	Romania		↘	↘
Continental				
	Austria		↘	↘
	Germany		↘	
	Romania		↘	
	Slovenia		↘	
Macaronesian				
	Spain		→	→
	Portugal		↘	
Mediterranean				
	Croatia		?	→
	Greece		→	
	Italy		→	
Pannonian				
	Czechia		→	→
	Hungary		→	
	Slovakia		→	
	Romania		→	
Steppic				
	Romania		↘	↘
Marine Atlantic				
	France		?	?
	Ireland		→	
	Portugal		?	
	Spain		?	
	United Kingdom		?	
Marine Baltic				
	Denmark		→	?
	Germany		↗	
	Poland		→	
	Sweden		?	
Marine Black Sea				
	Bulgaria		↘	↘
	Romania		?	
Marine Macaronesia				
	Portugal		→	→
Marine Mediterranean				
	France		?	?
	Greece		?	
	Spain		?	

Conservation status

 Good  Poor  Bad  Unknown

Conservation status trends

↗ Increasing → Stable ↘ Decreasing ? Unknown


Source: Article 17 reports and EU assessments

Looking at the species trends across the EU, some regions show more stable and increasing trends for species than for habitats (see Map 3.7). Based on Member States' reporting, some regions in Germany, Denmark, northern France and central Spain reveal predominantly deteriorating trends for species.

Species conservation status trends vary across Member States (see Figure 3.21 and Map 3.7). Those with the highest proportion of improving trends are Estonia, Luxembourg, Belgium and Denmark (over 20 %). Single species that show particularly positive trends within these countries are bat species, such as the Western

Barbastelle (*Barbastella barbastellus*), or the Eurasian Otter (*Lutra lutra*). Deteriorating species trends do not score as highly as in the habitat assessments: Italy is the only country with more than 50 % of deteriorating species trends. However, over 80 % of Member States report deteriorating trends for more than one fifth of their species assessments. Cyprus is the only Member State that did not report a single deteriorating trend, but unknown assessments exceed 75 %. Several Member States did not indicate any species assessment with improving trends (Bulgaria, Slovenia and Malta). They are also among the countries reporting more than 40 % of their conservation status trends as unknown.

Figure 3.21 Conservation status trends of species on Member State level




Note: Conservation status trends are based on species assessments. The number of assessments per Member State is indicated in parentheses. The total number of assessments is 7 589.


Source: Article 17 Member States' reports and assessments

Key messages


- 81 % of the **habitat** assessments show a poor or bad conservation status, of which only 9 % show improving trends and 36 % continue to deteriorate at the EU scale.
- Grasslands, dunes and bog, mire and fen habitats have the highest proportion of deteriorating trends (each over 50 %).
- Forest habitats exhibit the highest proportion of improving trends among the assessments (13 %).
- Only 6 % of all **species** assessments show an improving conservation status trend, whereas more than one third are still deteriorating. The trend is unknown for an additional 31 % of species.
- While fish have more improving trends than other species groups — together with amphibians — around half of their assessments show further deterioration.

 **DASHBOARD**
with detailed information by Member State:
conservation status and trends

Map 3.7 Spatial distribution of species conservation status trends at Member State level in a 10 km × 10 km grid



Spatial distribution of species conservation status trends at Member State level represented in a 10 x 10 km grid



Notes: The map shows the percentage of reports with improving (+), stable (=) and deteriorating (-) trends for each 10 km × 10 km grid cell. Reports with unknown trends are not included in the triangle; thus, grid cells with 100 % unknown trends appear in blue. In some exceptional cases, such as widely ranging but poorly known cetaceans, Member States submitted maps based on a 50 km × 50 km grid.

Source: Article 17 Member States' reports and assessments.

3.4 Improvement in status and trends

Changes in conservation status reported by Member States can be either 'genuine' or 'non-genuine'. Genuine changes refer to real changes in nature, rather than changes that are due to improved data or knowledge, taxonomic rearrangements or the use of different monitoring methods between subsequent reporting periods (non-genuine changes). The results for habitat and species assessments are very similar, with around 62 % of assessments reporting no change since the 2007-2012 reporting period. Non-genuine changes account for approximately 17 % of all reported changes in both habitat and species groups, due to improved knowledge and the use of different methods (see Figures 3.22 and 3.23).

Genuine changes can include both improvement and deterioration in conservation status. Around 1 % of all **habitat** assessments show improving genuine changes and 3 % show a genuine deterioration. Very few Member States reported genuine improvements in conservation status with the exception of, for example, Austria, France, Germany and the Netherlands.

Around 1 % of all **species** assessments present improving genuine changes and 2 % show a genuine deterioration. In total, 17 Member States reported


genuine improvements and 25 Member States reported deterioration in conservation status. The highest share of genuine changes with improving status were reported by France, Spain and Sweden.

3.4.1 Identifying improvements in status and trends

Beyond the classification of 'genuine' **improvements** in status, other positive developments are evident from Member States' data, namely improving status trends and stabilising status trends following a deteriorating for species and habitats. In order to not miss out on the cases in which improvements are due to conservation efforts, an amended methodology ⁽²¹⁾ was developed, in which improvement is understood as:

- **improved status** category since 2013 (e.g. U2 (bad) to U1 (poor), U1 (poor) to FV (good), U2 (bad) to FV (good)), which are reported by the Member States as 'genuine' changes in conservation status;
- **improving conservation status trends** for habitats/species with unfavourable status (including genuine and other than genuine changes), regardless of what the status was in the previous period; and


Figure 3.22 Type of change in the conservation status of habitats



Note: Non-genuine: method or data/knowledge improvement. No info: no info or N/A. The total number of assessments is 3 246.

Source: Article 17 Member States' reports and assessments.

Figure 3.23 Type of change in species in the conservation status of species



Note: The values include only species that are marked as 'present' or 'extinct'. Non-genuine: method or data/knowledge improvement. The total number of assessments is 7 612.

Source: Article 17 Member States' reports and assessments.

⁽²¹⁾ This methodology corresponds to the approach deployed for the national summaries.

- **stabilised trend** in assessments of unfavourable status in 2018 that had declining trends in 2013 (which are reported by the Member States as genuine changes).

Given the definition of 'favourable conservation status' in the Habitats Directive, changes in the overall conservation status (e.g. from poor or bad to good or from bad to poor) require relatively major changes in the individual conservation status parameters to be noted. The use of trends (improving, deteriorating, stable) in the overall conservation status allows more subtle changes (improvement or deterioration) of the unfavourable categories to be captured (e.g. U1 (poor); U2 (bad)). Improving conservation status trends refer to a situation in which the status is improving during the reporting period, as determined by a balance of the trends in the different parameters.

3.4.2 Improvements in habitats


The methodology described is applied in the following paragraphs and provides the basis for linking improvements to conservation measures (see Section 4.2.3). Figures 3.24 and 3.25 illustrate conservation improvements in habitats and species as reported by Member States. Figure 3.24 shows

that improvements were recorded for a total of 201 habitat assessments, including 20 cases of improving status, 161 cases of improving trends and 20 cases of stabilised trends. In total, 105 habitat types are represented.

On average 6 % of all habitat assessments show improvements. Forests and freshwater habitats each have 10 % of assessments showing improvements, while heath and scrub have 7 % and bogs, mires and fens, as well as dune habitats, have 6 %. In contrast, improvements are rarely reported for rocky habitats (3 % of assessments). Looking at the distribution of these improvements, some patterns can be identified:

- Most of the improvements in diverse forest types were reported from Bulgaria (37 of 73 cases), e.g. for eastern white oak woods (91AA) or beech forests (9110, 9130, 9150). Apart from these, forests in Belgium and Greece also reportedly improved more than average due to an improving trend.
- Freshwater habitats most frequently improved in Germany (8 of 22 cases), largely represented by alpine river habitat types (3220, 3230, 3240).
- Italy reported the highest share of improvements in grasslands (7 out of 26 cases), including for

Figure 3.24 Improvements in habitats reported by Member States



Note: The total number of habitat assessments showing improvement is 201.

Source: Article 17 Member States' reports and assessments.

mountain hay meadows (6520) and lowland hay meadows (6510).

Further illustrative examples of improvements are presented in Table 3.3.

The highest share of improvements with regard to the overall assessments in the group were recorded in the Black Sea (13 %), Atlantic (10 %) and Continental regions (7 %). These findings are promising, as the conservation status of habitats in the two latter regions are particularly critical (see Section 3.2.1). Four out of five marine regions show improvements. No records are included for the Marine Macaronesian region.

There are a variety of reasons underlying the improvement in habitats, such as targeted management

and restoration measures or expanding the habitat area. The conservation status of the *Luzulo-Fagetum* beech forest in Austria, for example, improved in the Alpine and Continental biogeographical regions. The main reasons for these improvements include an increase in its distribution area, improved tree species composition (a decrease in the proportion of non-native tree species, e.g. replacing spruce in the lower montane altitudinal range with native broadleaved trees) and improved management of deadwood and the forest structure.

The Boreal Baltic coastal meadows in Finland (see Box 3.8) illustrate how targeted and strategic nature conservation, restoration and management programmes can successfully contribute to improvements in conservation status.

Table 3.3 Examples of habitats showing improvements in their conservation status and trends

Habitat group	Member State	Habitat group	Habitat code	Habitat name	Improvements in conservation status and trend
Alpine	AT	Forests	9110	Beech forests (<i>Luzulo-Fagetum</i>)	U2 = → U1 +
Atlantic	NL	Dune habitats	2120	Shifting dunes along the shoreline with <i>Ammophila arenaria</i> ('white dunes')	U1 + → FV +
Atlantic	DE	Freshwater habitats	3260	Water courses from plain to montane levels with <i>Ranunculion fluitantis</i> and <i>Callitriche-Batrachion</i> vegetation	U2 = → U2 +
Continental	SI	Freshwater habitats	3180	Turloughs	U1 = → FV =
Mediterranean	FR	Coastal habitats	1510	Mediterranean salt steppes (<i>Limonietales</i>)	U2 = → U1 -
Pannonian	CZ	Heath and scrub	4030	European dry heaths	U1 - → U1 =
Boreal	SE	Forests	9070	Fennoscandian wooded pastures	U2 - → U2 =
Atlantic	UK	Bogs, mires and fens	7140	Transition mires and quaking bogs	U2 - → U2 =

Note Conservation status: FV, good; U1, poor; U2, bad. Trend: '+', improvement, '=', no change; '-'. Incomplete data sets with missing information on 'conservation status' are excluded.

Source: Article 17 Member States' reports and assessments.

Box 3.8 Closer Look: Boreal Baltic coastal meadows in Finland (1630)

Photos: Finnish Boreal Baltic coastal meadows © Mr Tapio Heikkilä, Finnish Ministry of the Environment

Boreal Baltic coastal meadows are low-growing plant communities found around the coastlines of the Baltic Sea on areas subject to land upheaval. Livestock have been grazing in these areas since prehistoric times, preventing the establishment of forests.

Pressures

In Finland, the abandonment of traditional agricultural activities has endangered this habitat. Other pressures include pollution of surface waters and groundwater (eutrophication of the Baltic Sea coastal waters), modification of coastline, estuary and coastal conditions for development and invasive alien species.

Conservation status and trends

Finland reported the overall conservation status of this habitat type as poor but improving (U1). The range of the habitat in Finland has good status, so the main issues to address were increasing the surface area of the habitat type and improving the structure and function of existing areas of habitat. The total surface area increased from 60 to 62 km². The status of the structure and function of the habitat improved from unfavourable-bad improving (U2+) to U1+. Approximately 68 % of the total habitat area is now in good condition and the trend is improving.

Drivers of improvement

The Natura 2000 network plays an essential role in the conservation of coastal meadows. Over 90 % of the total area is located within the network. There are currently 98 designated areas of coastal meadows, and six new designations have been given during the current reporting period. Coastal meadows are very important for breeding birds, given that, of the 98 total sites, 56 are also classified as SPAs under the Birds Directive. National funding will be directed to the management of traditional rural biotopes through a new strategic nature conservation, restoration and management programme (HELMI programme, 2020-2030) and coastal meadows are included within a recently launched LIFE project (CoastNet LIFE, LIFE17 NAT/FI/000544) focusing on the restoration of coastal habitats. However, the key funding element for coastal meadows will continue to be the agri-environment payments through the Finnish rural development programme. The continuation of the positive trend in coastal meadows in Finland is dependent on the next period of the common agricultural policy.

Major achievements

The area of managed coastal meadow in Finland has increased by several hundred hectares since 2007. Coastal meadows are also a key breeding habitat for highly protected bird species (Annex I of the Birds Directive), such as Baltic Dunlin (*Calidris alpina schinzii*), Ruff (*Calidris pugnax*) and Black-tailed Godwit (*Limosa limosa*). Thanks to successful management efforts, especially in the northern parts of Bothnia Bay, the breeding population of these rare species is increasing locally. Environmental education is ongoing, as wetland programmes and weekend tours for families are still taking place 10 years after projects ended (NEEMO LIFE team, 2017). Visitor numbers to the Natura 2000 sites are high; the bird observation towers are very popular and draw thousands of visitors annually.

Source: Keränen et al. (2020).

3.4.3 Species improvements

Improvements were recorded in 419 species assessments, including 108 cases of improving status, 296 cases of improving trends and 15 cases of stabilised trends (as shown in Figure 3.25). In total, 208 individual species are represented.

On average, 6 % of all species assessments show improvements. The species group with the largest number of improvements is mammals, followed by vascular plants, and fish. Approximately 8 % of all recorded mammal assessments show improvements, followed by fish (8 %) and vascular plants (5 %). Amphibians, molluscs, non-vascular plants and reptiles show fewer improvements in their respective groups (between 2 % and 4 %). Nearly 80 % of species showing improvements in status and/or trends are animals and 20 % are plant species.

- For mammals, most of the improvements were reported by France (22 cases), Sweden (20 cases) and Germany (16 cases). Improvements in France include, for example, several bat species (e.g. the Lesser Horseshoe Bat (*Rhinolophus hipposideros*) and the Eurasian Lynx (*Lynx lynx*)). Most of these improvements are due to improving trends.
- Vascular plants improved particularly in Spain (13 of 73 cases), Poland (10 cases) and Austria (9 cases). In Austria, the flowering plant *Dracocephalum austriacum* improved from a bad (U2) to a poor (U1) status in the Continental and Alpine biogeographical regions. In Spain, the endemic plant *Globularia sarcophylla* from Gran

Canaria achieved good conservation status. Poland reported that three plants achieved good conservation status: *Agrimonia pilosa*, *Ligularia sibirica* and the near-threatened *Linaria loeselii*.

- The least improvements were recorded for non-vascular plants, molluscs and reptiles. In detail, 97 out of 208 species (e.g. Grey Wolf (*Canis lupus*) and Eurasian Beaver (*Castor fiber*)) showed improvements in more than one biogeographical region and/or Member State.

In total, 402 of the 419 assessments with improvements refer to terrestrial species. Within the 17 cases of marine species that showed improvements, most are mammals (e.g. Grey Seal (*Halichoerus grypus*) and Harbour Seal (*Phoca vitulina*)) and one reptile, the Green Turtle (*Chelonia mydas*).


Table 3.4 below presents further examples of species that showed an improvement in their conservation status since the last reporting period (2007-2012).

On the biogeographical level, improvements are present in all regions. Most cases are reported for the Atlantic (10 %), Boreal (7 %) and Continental (7 %) regions. Cases from other regions are less frequent. For the marine regions, only three of five marine regions show improvements. No records are included for the Marine Macaronesian or Black Sea regions.

The overall number of improving habitats and species and examples from some selected biogeographical and marine regions are given in Box 3.9.

Figure 3.25 Improvements in species reported by Member States

Number of assessments




Note: The total number of improved cases is 419.


Source: Article 17 Member States' reports and assessments.

Box 3.9 Improvements in habitats and species in biogeographical and marine regions**ALPINE****Wolf (*Canis lupus*), Slovenia**


Improved as a result of an action plan based on scientific knowledge, including activities to improve wolf-human coexistence, e.g. electric fences and shepherd dogs to protect flocks of sheep (SloWolf project).

**ATLANTIC****Rivers with muddy banks with *Chenopodium rubri* and *Bidention vegetation*, Netherlands**






Improved by floodplain restoration and reduction of diffuse pollution to surface waters or groundwaters from agricultural activities.

**BOREAL****Northern Atlantic wet heaths with *Erica tetralix*, Latvia**

Improved by restoration of at least 40 ha of wet heath within the Adazi military training area (mostly covered by Natura 2000 area).

**CONTINENTAL****Agile Frog (*Rana dalmatina*), Sweden**

Improved by creation and restoration of wetlands, creation of hibernation spots, site-specific management plans for future conservation and monitoring, increasing awareness and acceptance of restoration measures (SemiAquaticLife project).

**MACARONESIAN*****Teline rosmarinifolia*, Spain**

Improved as a result of habitat improvement and protection from goats, grazing management, avoidance of trampling due to recreational activities (hunting, fishing), management of harvesting and collection of plants, and close monitoring.

MARINE ATLANTIC**Harbour Seal (*Phoca vitulina*), Netherlands**

Improved as a result of rehabilitation centres with a dispensation to rehabilitate sick, orphaned or injured seals and reintroduce rehabilitated seals and of strict conservation of seal populations and environmental sanitation of neighbouring waters.

PANNONIAN**Forest-steppe Mugwort (*Artemisia pancicii*), Czechia**

Improved by removal of shrubs, mowing, control/eradication of invasive alien species, and reintroduction of sustainable grazing within Natura 2000 sites.

MEDITERRANEAN**Coastal lagoons, France**

Improved by reducing the impact of multipurpose hydrological changes and managing habitats (other than agriculture and forest) to slow, stop or reverse natural processes.

Note:

The box shows the exact number of habitats and species for each biogeographical and marine region, not the number of improved assessments (including improving trends, improved statuses and stabilised trends). Further regions, not shown in the figure include Steppic (habitats 0; species 3), Black Sea (habitats 9; species 1), Marine Black Sea (habitats 0, species 0), Marine Baltic (habitat 1; species 6) and Marine Mediterranean (habitats 0; species: 2).

Table 3.4 Examples of species showing improvements in their conservation status and trend by species group

Taxonomic group	Examples			Improvements in conservation status and trend
	Biogeographical regions	Member State	Species name	
Amphibians (23)	Continental	FR	Moor Frog (<i>Rana arvalis</i>)	U2- → U2=
	Continental	LU	Yellow-bellied Toad (<i>Bombina variegata</i>)	U2= → U2+
Arthropods (55)	Atlantic/Mediterranean	ES	White-clawed Crayfish (<i>Austropotamobius pallipes</i>)	U2+ → U1-
	Alpine	IT	<i>Saga pedo</i>	U1= → U1+
Fish (71)	Continental	CZ	European Bitterling (<i>Rhodeus amarus</i>)	U2= → U1=
	Mediterranean	IT	Mediterranean Trout (<i>Salmo cetti</i>)	U2- → U2+
Mammals (164)	Boreal	SE	Natterer's Bat (<i>Myotis nattereri</i>)	U2- → U1+
	Continental	CZ	Eurasian Lynx (<i>Lynx lynx</i>)	U1= → U1+
Molluscs (9)	Continental	CZ	Geyer's Whorl Snail (<i>Vertigo geyeri</i>)	U2+ → FV=
Non-vascular plants (10)	Atlantic	NL	Slender Green Feather-moss (<i>Hamatocaulis vernicosus</i>)	U2+ → U1+
	Continental	CZ	<i>Dicranum viride</i>	U1= → U1+
Reptiles (12)	Mediterranean	ES	Hermann's Tortoise (<i>Testudo hermanni</i>)	U2- → U1-
	Atlantic	FR	<i>Hierophis viridiflavus</i>	U1= → U1+
Vascular plants (73)	Mediterranean	FR	<i>Arenaria provincialis</i>	U1= → FV+
	Macaronesian	PT	<i>Prunus azorica</i>	U2= → U2+

Note: Conservation status: FV, good; U1, poor; U2, bad. Trend: '+', improvement, '=', no change; '-', deterioration. Incomplete data sets with missing information on 'conservation status' are excluded. Number of assessments in parentheses; 419 species assessments in total, including 'other invertebrates' (2 cases).

Source: Article 17 Member States' reports and assessments.

Key messages

- On average, 6 % of all habitat and 6 % of all species assessments show improvements in conservation status and/or trends, representing 201 habitats and 419 non-bird species assessments. In total, 105 habitat types and 208 non-bird species show improvements.
- Forests and freshwater habitats show the highest number of improvements among habitats.
- The main reasons for improvements in habitats include targeted management and restoration measures or expansions in the habitat's area.
- Nearly 80 % of species showing improvements in status and/or trends are animals (dominated by mammal and fish) and 20 % are plant species.
- For habitats and species, most improvements were recorded in the Continental region, followed by the Atlantic and Alpine regions.



Figure 4.1 Summary of pressures and responses

Pressures and responses

The analysis of pressures and responses looks jointly at the results from both nature directives' reporting. Member States have reported over 200 different pressures categorised into 15 overarching sectors, and over 100 conservation measures are listed in 13 main categories, corresponding to the pressure sectors identified.

Member States reported over **67 000** individual pressures

With **21 %**, **agriculture** is the most frequently reported pressure for habitats and species. Abandonment of grasslands and intensification is particularly impacting pollinator species, farmland birds and semi-natural habitats

Urbanisation and leisure activities account for **13 %** of all reported pressures, representing 48 % of all marine pressures.

Invasive alien species such as the False Indigo-bush, particularly affect dunes and sclerophyllous scrubs as well as species such as breeding seabirds.

The **modification on water regimes**, physical alterations of water bodies and removal of sediments predominantly affect freshwater habitats and fish.

Forestry activities represent **11 %** of all pressures, particularly affecting forest habitats, and woodland species.


18 % of all pressures for **birds** stem from the exploitation of species, mainly relating to **illegal killing and hunting**. In Europe, the annual hunting bag amounts to at least 52 million birds.

Climate change is reported as a rising threat, particularly due to ongoing changes in the temperature and the decrease of precipitation.

Almost **50 %** of all pressures related to **pollution** can be attributed to air, water and soil pollution caused by agriculture.

How are these pressures addressed?

Over **6 000** targeted **conservation measures** aim to **maintain** or to **restore** the current status.




Birds such as vultures particularly benefit from species action plans, agri-environmental measures and large-scale conservation efforts.



Many **habitats** improved through targeted measures: adapted grassland management, the control of invasive species or the reinforcement of sustainable tourism.

Several **non-bird species**, and especially mammals and fish improved through targeted measures such as reducing certain human activities and pollution from different sources.



4 Pressures and responses

As one of the most densely populated regions in the world, Europe's human activities have been driving the decline and deterioration of many of its species and habitats. To understand the most critical pressures and threats underlying this trend, the nature directives require Member States to report on what they consider to be the principal causes of species loss and habitat degradation per single species and per habitat. Pressures are considered to be factors that have affected habitats and species within the current reporting period, while threats are factors that are anticipated to be likely to have an impact during the subsequent two reporting periods.

At the same time, Member States report on measures taken to maintain or to restore the species or habitats to achieve good conservation status. Under the nature directives' reporting, conservation measures are principally understood to be practical actions to mitigate the impact of past and present pressures, and they are not covered by many legal or statutory measures (e.g. strict species protection or designation of protected sites). The Natura 2000 network (see also Chapter 5) can be seen as the overarching measure to be implemented by Member States and at the same time as a legal framework for applying practical conservation actions. The LIFE programme is the EU's major dedicated funding instrument for implementing such target conservation measures (among other funding objectives). Its environment sub-programme funds nature conservation projects particularly in the areas of biodiversity and of protected habitats and species. It provides grants for best practice, pilot and demonstration projects that contribute to the implementation of the EU Birds and Habitats Directives and the development, implementation and management of the Natura 2000 network.

This chapter gives an overview of the pressures and threats that were reported by Member States (Section 4.1) and of the implementing status of targeted conservation measures (Section 4.2). As restoration activities in particular play a central role in conserving European biodiversity, a dedicated analysis of restoration needs for habitats is presented in Section 4.3. The analyses are

based on Member States' data and were conducted for both directives. Therefore, the results are presented jointly, with birds (all seasons) included in the species category. A more detailed description of the methodology used for the assessment of pressures and conservation measures can be found in the methodological report (Röschel et al., 2020).

4.1 Key pressures for species and habitats

Article 12⁽²²⁾ and Article 17 reporting on pressures and threats is structured into two hierarchical levels, with differing degrees of detail. The first (level 1) comprises 15 overarching categories, while the second (level 2) identifies 203 individual pressures/threats (mostly listed as 'activities'). At the same time, pressures and threats are ranked as being of either 'high importance' or 'medium importance' according to their relative impact; Member States are restricted to selecting a maximum of 10 listed pressures/threats and a maximum of five pressures with a 'high importance'. The following analysis focuses on the 'high importance' category of pressures, given that the results for reported pressures and threats are largely consistent among the categories. The complete results of the analysis, including the frequency of pressures and threats of 'medium importance', can be accessed via the dedicated *State of nature in the EU* web page⁽²³⁾.

Member States reported over 67 000 records from one of the above 203 individual pressures for both species (including birds) and habitats. Around one third of these reported pressures are considered to be of high importance, with slightly more pressure records of high importance for non-bird species (35 %) than for habitats and birds (around 32 %).

The most frequently reported pressures for both habitats and species stem from agricultural activities and urbanisation. While the context and dynamics driving habitat degradation and species decline are highly diverse, agricultural activities (or in some cases,

⁽²²⁾ In the case of Article 12, the reporting on pressures was only mandatory for regularly occurring Annex I species and any other migratory species triggering special protection area designations nationally. So, although Member States were encouraged to provide additional information for Annex II and remaining bird species, this potentially leads to an underestimation of pressures, such as those arising from hunting.

⁽²³⁾ <https://www.eea.europa.eu/themes/biodiversity/state-of-nature-in-the-eu/state-of-nature-2020>

the lack thereof) represent the most common group of pressures. As indicated in Figure 4.2, many terrestrial habitats are severely impacted by agriculture, especially grasslands and freshwater habitats, heath and scrub, and bogs, mires and fens. This is also the case for most of the species groups, including reptiles, molluscs, amphibians, arthropods, vascular plants and breeding birds.

Looking at different habitats and species groups, however, a diverging importance of pressure categories can be identified. For example, wintering and passage birds face different key pressures from those experienced by breeding birds. For the former, the exploitation of species (mostly hunting and illegal killing) is reported to be the major impact. Forestry activities are the second largest pressure category for species, affecting in particular arthropods, mammals and non-vascular plants. In contrast, habitats such as dunes or coastal and rocky habitats are primarily

affected by urbanisation. Compared with species, most habitat groups are particularly vulnerable to natural processes (e.g. succession of semi-natural habitats). However, succession — contrary to the guidance provided — was in many cases a response to other human intervention or management changes, such as abandonment of agricultural land or drainage of bogs and mires.

When interpreting the broad pressures for habitats and species, it is important to note that the categories 'Modification of water regimes' and 'Pollution' include only multipurpose activities and pollution from mixed sources; the pressures from specific activities, e.g. relating to agricultural pollution or hydrological changes, are included under the corresponding sectoral headers (in this example agriculture). When looking at the global distribution across the different level 1 categories, the modification of hydrological regimes (including both multipurpose changes under

Figure 4.2 Distribution of level 1 pressure categories among habitats and species



Note: The size of the squares and their shade reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages.

Source: Article 12 and Article 17 Member States' reports and assessments.

the category 'Modification of water regimes' and also changes attributed to other categories) is in fact one of the most important overarching pressures. In the case of pollution, related pressures are in fact mostly covered under agriculture, which accounts for 48 % of all pollution-related pressures.

Level 2 of the pressures reported provides in-depth information on the underlying impacts that are relevant in each level 1 pressure category. The following section focuses on the most important pressure categories and their effects on different habitats and species reported as level 2 pressures.


4.1.1 Agriculture

About 40 % of the total land area of the EU-28 is agricultural land (Eurostat, 2020a). Results show, that current agricultural practices are by far the

most dominant driver affecting habitats and species (see Figure 4.2). However, the richness and abundance of biodiversity associated with agricultural habitats is strongly correlated with the degree of modification (e.g. draining, ploughing) and the intensification of management (e.g. use of fertilisers, irrigation and pesticides). Extensive agricultural management creates and maintains semi-natural habitats with a diverse fauna and flora. Since the 1950s, however, the intensification and specialisation of the agricultural sector has increasingly contributed to ongoing biodiversity loss.

Changes in agricultural management are thus the most frequently reported type of pressure. When looking at the importance of individual reported pressures (Figure 4.3), terrestrial habitats and non-bird species such as grassland habitats, vascular plants and arthropods such as the Marsh Fritillary (*Euphydryas aurinia*) are particularly impacted by the

Figure 4.3 Distribution of the eight most relevant level 2 agricultural pressures for habitats and species, shown as the percentage of pressures within this level 1 group



Notes: The size of the squares and their shade reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages. Total number of reports is given in parentheses.

Source: Article 12 and Article 17 Member States' reports and assessments.

abandonment of grassland management, most importantly the ceasing of traditional or extensive grassland management. Among others, these pressures also badly affect pollination capacities (see Box 6.2). Reptiles and smaller mammals are especially affected by **fragmentation due to the removal of small landscape features**, which reduces landscape connectivity and leads to a loss of habitat area essential for food supply, shelter and breeding sites. Birds, however, are most affected by the **conversion of one type of agricultural land use to another** (e.g. this ranges from conversion from extensive to intensive agricultural land to more subtle changes such as a change in the type of crop grown) and by **drainage**. Drainage activities undertaken in an effort to increase agricultural land area lead to the loss of specialised habitat, and of food supply and breeding sites for species. Wintering birds are particularly affected by ongoing and past drainage activities.

Fertilisers and the **use of plant protection** products are reported to have a considerable impact on many habitats and species. This holds especially true for plant protection chemicals and their effects on amphibians, insects, mammals — mainly bats but also small mammals such as the European Ground Squirrel (*Spermophilus citellus*) or the European Hamster (*Cricetus cricetus*) — and birds. A report on 576 species of butterflies in Europe, for example, found that fertilisers and pesticides negatively affected 80 % of the (often now threatened) species (Sánchez-Bayo and Wyckhuys, 2019). This also indirectly affects insect-eating birds such as the Common Swift (*Apus apus*), a once common and widespread long-distance migrant, which is undergoing major population declines in most Member States. Recent analyses point to the extensive use of pesticides as the primary factor responsible for the decline of birds in farmland ⁽²⁴⁾ (Mineau and Whiteside, 2013). Moreover, **agricultural pollution of surface waters or groundwaters** has significant impacts on standing waters, rivers, ponds and marine habitats as well as on their species.

Despite the introduction of provisions for the more sustainable management of natural resources and climate action ⁽²⁵⁾ in recent reforms of the common agricultural policy (CAP), these have not significantly

reduced the negative effects of agriculture on biodiversity and have not substantially contributed to the conservation and restoration of species and landscapes. This is acknowledged in both the EU Biodiversity Strategy for 2030 and the Farm to Fork strategy (EC, 2020b, 2020c).

4.1.2 Urbanisation

Urbanisation includes development but equally the use of residential, commercial, industrial and recreational areas and dispersed recreational and leisure activities. It is one of the key pressures affecting habitats and species. According to the Member States' reports, major human-induced pressures from this broad group are **sports, tourism and leisure activities** (Figure 4.4). This includes activities such as outdoor sports, leisure aircraft, drones, human trampling and unregulated wildlife watching. Overall, the extension of urban areas and artificial surfaces is the dominant group of pressures for marine and coastal habitats (e.g. dominant for 28 % of the lagoons and for 38 % of sandy coasts). Marine habitats suffer particularly from sports and leisure activities, especially in the Mediterranean and Macaronesian regions. For coastal habitats such as lagoons, estuaries, sandy coasts and rocky shores, **coastline modifications** are a dominant pressure. Tourism-related activities are one of the major underlying causes behind dune habitat degradation and are often related to the broader modification of coastline conditions for commercial and recreational purposes. Urban development — as a consequence of tourism or other drivers — thus considerably decreases the intactness of marine and coastal habitats, critically affecting the species depending on these habitats. Tourism activities, for example, particularly disturb breeding birds, specifically water birds such as ducks, geese, herons and grebes or threatened raptors such as the Egyptian Vulture (*Neophron percnopterus*) or the Bearded Vulture (*Gypaetus barbatus*) when they occur within their nesting areas.

Another important pressure caused by urbanisation is **the conversion of natural and semi-natural land to housing, settlement or recreational areas**. This


⁽²⁴⁾ The current status of and trends in farmland and grassland-dependent habitats and species are explored in more detail in Section 6.2.1

⁽²⁵⁾ Greening measures under pillar 1 (direct payments) aiming to maintain permanent grassland areas, foster crop diversification and promote areas for nature and habitats, including the maintenance of landscape elements (ecological focus areas). In addition, agri-environmental and climate measures aim to encourage trends towards extensification, promote organic farming and encourage the maintenance of low-intensity management on high nature value farmland.

is particularly relevant for grassland habitats and forests as well as for the species they support, such as insects, reptiles and breeding birds. However, the most frequently reported pressure for non-bird species is **construction or modification within already existing urban or recreational areas**. This

could involve, for example, demolishing structures or deliberately closing damaged roofs by repairing them. Such man-made habitats have become important sheltering areas for some species, such as bats ⁽²⁶⁾; closing damaged roofs during breeding or hibernation can therefore harm dependent populations.

Figure 4.4 Distribution of the eight most relevant level 2 urban pressures for habitats and species, shown as the percentage of pressures within this level 1 group



Note: The size of the squares and their shade reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages. Total number of reports is given in parentheses.

Source: Article 12 and Article 17 Member States' reports and assessments.

⁽²⁶⁾ Bats have a special spectrum of pressures. Pressures related to urbanisation are predominant (28 %), followed by pressures from forestry (24 %) and pressures linked to agriculture (19 %). The reason could be the mobility of these species and their larger ecological valence — use of several different habitat types. Bats use agricultural habitats mostly for foraging, but their resting/breeding places are often in urbanised areas or forest (some species use tree holes).


4.1.3 Forestry

Over the last few centuries, forests managed to varying degrees of intensity have replaced almost all of Europe's natural forests. Currently, less than one third of Europe's forests are uneven-aged, 30 % have only one tree species (mainly conifers), 51 % have only two to three tree species, and only 5 % of forests have six or more tree species (Forest Europe, 2015). These general tendencies are not reflected in the reported pressures on forest habitats and species, as in fact many commercial forests do not classify as Annex I habitat types or do not constitute suitable habitats for protected species. However, the increased extraction of forest products and intensified forestry practices have diverse impacts on the various habitats and species protected under the nature directives. The recently increasing use of forests as a source of renewable energy poses one of the major forest-related policy challenges. Current studies suggest that these harvesting activities lead not only to a decline

in forest area but also to potentially a more than 20 % reduction in their capacity for carbon sequestration (Searchinger et al., 2018; Fern, 2020).

Forest-dependent insects, mammals, non-vascular plants and breeding birds such as the Lesser Spotted Woodpecker (*Dryobates minor*) are most heavily affected by an excessive **removal of dead and old trees or the reduction of old-growth forests** (Figure 4.5). The clear-cutting of forested areas is considered to be the most relevant pressure for breeding birds in the context of forestry. Although new trees should be replanted or allowed to regrow after the forest stands have been clear-cut — as required by national forest acts in Europe — deforestation and clear-cutting without regrowth is occurring in Europe (EEA, 2016). Apart from breeding birds that depend on forest habitats, old trees are particularly valuable for some bats and small mammals, such as the Western Barbastelle (*Barbastella barbastellus*), the Caucasian Squirrel (*Sciurus anomalus*) or the Forest

Figure 4.5 Distribution of the eight most relevant level 2 forestry pressures for habitats and species, shown as the percentage of pressures within this level 1 group



Notes: The size of the squares and their shade reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages. Total number of reports is given in parentheses.

Source: Article 12 and Article 17 Member States' reports and assessments.

Dormouse (*Dryomys nitedula*). For species other than birds and for habitats, however, the most frequently reported forestry pressure is the **removal of dead trees**. Many insects, non-vascular plants, amphibians and reptiles depend on these for food, breeding places and shelter. Dead wood specialists such as the near-threatened Hermit Beetle (*Osmoderma eremita*) need dead, decaying wood for their life cycle. These structures thus represent integral features of healthy forest habitats; their removal can lead to changes in forest structure and diversity (Paillet et al., 2010; Vilén et al., 2015).


Not surprisingly, forestry is the dominant group of pressures reported for most of the Annex I forest types: it amounts to 50 % of all pressures for mixed forests, broadleaved deciduous and coniferous forests. For broadleaved evergreen forests, however, forestry accounts for only 20 % of the pressures, which is equal to the degree of impact arising from agricultural activities.

Forest habitats are especially affected by the removal of dead and dying trees as well as by broader land use changes, such as **conversion to monocultures or other forest types**. Other habitat groups, e.g. grassland and heath habitats, are more significantly affected by ongoing **afforestation dynamics** in the EU, which decrease the area of open landscape structures.

4.1.4 Exploitation of species

The exploitation of species is the most frequent pressure group for wintering and passage birds (see Figure 4.2). Impacts on bird species largely relate to **illegal shooting or killing** (27 %) and **hunting** (19 %) (Figure 4.6). Recent research in 26 European countries has estimated an annual hunting bag of at least 52 million birds, excluding a significant number of killings in areas just beyond the study area and in European countries for which no data were available

Figure 4.6 Distribution of the eight most relevant level 2 pressures caused by exploitation of species for habitats and species, shown as the percentage of pressures within this level 1 group



Notes: The size of the squares and their shade reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages. Total number of reports is given in parentheses.

Source: Article 12 and Article 17 Member States' reports and assessments.

(Hirschfeld et al., 2019). Hunting bag information — although mandatory in Article 12 reporting for those Annex II birds that are hunted nationally — still shows significant gaps: in over 78 % of reports no information on hunting bags was provided. The only countries with more than half of their entries filled in were Malta, France and Poland; the former two are also the countries with the highest share of hunting reported. Three countries (Ireland, the Netherlands and the United Kingdom) did not provide any hunting bag data, and, according to the data reported, no bird species is nationally hunted in the Netherlands. However, hunting pressure is related not only to direct increased mortality of the target species but also disturbance and reducing the amount of prey available.

Non-bird species are also affected by exploitation, with an above average impact on fish, mammals and reptiles. Fish are among the most affected by **marine and freshwater harvesting** (Figure 4.6). The impacts on mammals can be divided into two main groups: large terrestrial mammals that are mostly exposed to illegal killing and marine mammals that are mainly affected by **bycatch and marine harvesting activities**. Terrestrial mammals that are particularly affected include the Grey Wolf (*Canis lupus*), Eurasian Lynx (*Lynx lynx*) and Eurasian Otter (*Lutra lutra*), whereas marine mammals include the Short-beaked

Common Dolphin (*Delphinus delphis*) and Harbour Porpoise (*Phocoena phocoena*), among others. Bycatch and marine harvesting is also a significant pressure on island breeding seabirds (see Box 4.1) and sea ducks (Mergini) and has a significant impact on their breeding success.

The exploitation of species can affect the intactness of their habitats. This is especially relevant for coastal and marine habitats because fish and shellfish harvesting (e.g. by bottom trawling) cause physical loss of and disturbance to seafloor habitats and reduce prey populations, which again disturb marine species.

4.1.5 Invasive alien species

Invasive alien species (IASs) are animals and plants that are introduced accidentally or deliberately into a natural environment where they are not normally found, causing serious negative effects in their new environment. IASs represent a major and increasing threat to native European flora and fauna and cause billions of euros worth of damages every year to the European economy. The present IAS pressure category also includes issues arising from interactions with problematic native species, disease or pathogens.

Box 4.1 Closer look: Balearic Shearwater



Photo: Balearic shearwater © Cabrera Natura

The Balearic Shearwater (*Puffinus mauretanicus*) is an Annex I Spanish-breeding endemic seabird whose entire population is located within the Natura 2000 network. While the species is challenging to monitor (information is based on the extrapolation of data from single colonies using indirect methods that could undervalue the existing breeding population), it has been assessed as critically endangered since 2004.

As with many seabirds, this species continues to be significantly threatened by bycatch and incidental killing (due to fishing and hunting activities), as well as invasive alien species (other than species of Union concern). In particular, these birds are threatened by predation at their breeding colonies by introduced mammals (such


as feral cats and genets and, to a lesser extent, rats) as well as at-sea mortality as a result of interactions with commercial and artisanal fisheries. Further threats reported by EU Member States include mixed source marine water pollution, marine fish and shellfish harvesting (causing a reduction in both species and prey populations and general disturbance), and residential or recreational activities and structures generating various kinds of pollution. Finally, the species is also subject to a lesser extent to roads and related infrastructure, pollution from industrial and commercial activities, and modification of hydrological flows. The population is consequently decreasing in the short and long terms.

To improve the status of the Balearic Shearwater, there is an urgent need for habitat restoration, improvement and management. This needs to be accompanied by better monitoring of the impact of the aforementioned pressures as well as reductions in the impact of mixed source pollution and the threats posed by bycatch, unsustainable fishing and invasive alien species.

In 2015, EU Regulation 1143/2014 on invasive alien species (IAS Regulation; EU, 2014) entered into force and identified **invasive alien species of Union concern** (Figure 4.7). These species cause damage such as predation of adults and chicks/eggs by — inter alia — the Coypu (*Myocastor coypus*) or the Raccoon (*Procyon lotor*), browsing of understory habitats by Reeves' Muntjac (*Muntiacus reevesi*), and competition for food from Red Swamp Crayfish (*Procambarus clarkii*). Within the first reporting period of the IAS Regulation, Member States provided specific information on 48 IASs of Union concern ⁽²⁷⁾. According to Member States' reporting, IASs of Union concern represent around 20 % of pressure reported for invasive species, while much greater impact is reported from **IAS other than species of Union concern** ⁽²⁸⁾. These include, among others, the loss of (native) prey species owing to competition from the Pacific Oyster (*Crassostrea gigas*), decline in the quality of native vegetation owing to the spread of invasive plant species, and the overgrowth of alluvial and lowland humid habitats with invasive plants (e.g. Japanese Knotweed (*Fallopia japonica*) or Himalayan Balsam (*Impatiens glandulifera*)). A prominent example of an invasive predator is the American Mink (*Neovison vison*), which in Europe has decimated seabird colonies and reduced some waterfowl populations, such as those of the Coot (*Fulica atra*), Moorhen (*Gallinula chloropus*) and other Rallidae (Ferrerias and Macdonald, 1999; CABI, 2020). This carnivore not only poses a threat to many internationally important populations of ground-nesting birds but also severely affects local populations of amphibians, reptiles and small mammals, such as the European Water Vole (*Arvicola amphibious*) in the United Kingdom (CABI, 2020). IASs are generally reported as highly relevant for habitats, particularly for dune habitats and sclerophyllous scrub ⁽²⁹⁾. A decline in the quality of native vegetation in the sclerophyllous scrub habitat is, among other causes, due to the spread of the False Indigo-bush (*Amorpha fruticosa*). This fast-growing, deciduous shrub forms a dense thicket that outcompetes the native flora and changes successional patterns (CABI, 2020). The pressures caused by IASs are not distributed homogeneously across the EU. Dune habitats, for example, are particularly affected by IASs in the Atlantic and Pannonian regions and sclerophyllous scrub in the Macaronesian region. In some parts of Europe, floodplains and riparian areas in particular are subject to invasion by IASs.

In these areas, invasive plant species (e.g. Giant Goldenrod (*Solidago gigantea*), Japanese Knotweed and Himalayan Balsam) are spreading along water courses. While forests are generally not as badly affected by IASs as other habitat types, broadleaved evergreen forests are the exception. For them, IASs other than of Union concern is the single most

Figure 4.7 Distribution of level 2 pressures caused by invasive alien species for habitats and species, shown as the percentage of pressures within this level 1 group



Notes: The size of the squares and their shade reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages. Total number of reports is given in parentheses.

Source: Article 12 and Article 17 Member States' reports and assessments.

⁽²⁷⁾ In August 2019, an extension brought the list to 66 species (36 plants, 30 animals).

⁽²⁸⁾ This group includes species introduced in the modern period that are established in the wild outside their natural range and excluded from the IAS Regulation.

⁽²⁹⁾ The reporting of invasive alien species is still very uneven across Member States and the nature directives; some Member States reported invasive alien species as a problem in many instances and some did not report any problems at all.

reported pressure, amounting to 17 % of all reported pressures. IASs of Union concern are most often reported as pressures for coastal habitats, followed by forest and freshwater habitats.


Amphibians, fish, vascular plants and birds — especially breeding seabirds such as shearwaters or storm-petrels — are the species that are most affected by IASs (see Box 4.1). Amphibians are also affected by animal diseases to an exceptionally high extent. As an example, a recent study on the amphibian fungus *Batrachochytrium salamandrivorans* concluded that the disease is more widely distributed than previously thought and can cause localised extinction of amphibian species (Spitzen-van der Sluijs et al., 2016).

Compared with the last reporting period (2007-2012), the impact of IASs increased in importance for both habitats and species. IASs are also considered to be a growing threat in the future.

4.1.6 Natural processes

Natural processes refer to, for example, vegetation succession and biocenotic evolution, abiotic natural processes and interspecific faunal relations. These processes are ongoing and are critical to sustain natural life. Habitats and species are subject to such natural processes over long time-frames as they evolve together, with strong impacts on species diversity and habitat composition. However, only when an ecosystem's natural balance is distorted as a result of accelerating climate change or direct human intervention (e.g. by eliminating disturbance regimes like natural floods, the presence of wild herbivores or large carnivores, or by confining dynamic early succession habitats to static fragments in an otherwise unavailable agricultural and forest landscape) can they become considerable pressures. The most relevant natural process-related pressure is **natural succession** (Figure 4.8), resulting in changes in species

Figure 4.8 Distribution of level 2 pressures caused by natural processes for habitats and species, shown as the percentage of pressures within this level 1 group



Notes: The size of the squares and their shade reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages. Total number of reports is given in parentheses.

Source: Article 12 and Article 17 Member States' reports and assessments.

composition. However, this pressure has often been mistakenly reported by Member States as indicating a lack of appropriate management of certain habitats (e.g. abandonment of grazing, lack of hay cutting).

Bogs, mires and fens are among the habitats most vulnerable to natural succession, usually indirectly caused by human intervention (e.g. lowering water tables by drainage). Other habitats affected by natural processes are dunes and grasslands. Birds, however, according to Member States' reports, are mostly exposed to **interspecific relations** largely associated with competition, predation and parasitism. Certain groups of birds, such as waders, gulls, shearwaters and storm-petrels, are particularly affected by interspecific relations. These include, among others, predation by other species and competition for nesting sites or food (see Box 2.2).


Of the Article 17 species, vascular plants most frequently face pressures from natural processes. The Lady's Slipper Orchid (*Cypripedium calceolus*), Fen Orchid (*Liparis loeselii*) or the Water Shamrock (*Marsilea quadrifolia*), for example, experience interspecific relations as their predominant overall pressure. In the case of the Fen Orchid, the plant depends on the natural processes of land upheaval around the Baltic Sea: old sites become overgrown when they rise too high above sea level.

4.1.7 Modification of water regimes

Pressures concerning modifications to the water regime are highly fragmented among the different pressure groups, overall accounting for 11 % of all reported pressures. This specific pressure group only accounts for pressures that are not directly related to any other pressure group. For example, drainage activities that are part of agricultural activities are included in the agricultural pressure group, and hydropower installations are included in the energy pressure groups, making up 14 % and 13 % of water-related pressures. Following this logic, the present pressure group covers only 58 % of all reported pressures related to human-induced changes in water regimes, i.e. only those that are not covered by others.

Unsurprisingly, pressures related to this group are particularly relevant for freshwater habitats and fish. **Modifications of the hydrological flow** (Figure 4.9) are reported to be the pressure with the most significant impacts on European freshwaters. This hydrological pressure is mostly related to running water and includes, for example, modification of flooding regimes or cutting of aquatic and bank

Figure 4.9 Distribution of level 2 pressures caused by modification of water regimes for habitats and species, shown as the percentage of pressures within this level 1 group




Notes: The size of the squares and their shade reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages. Total number of reports is given in parentheses.

Source: Article 12 and Article 17 Member States' reports and assessments

Box 4.2 Closer look: Pressures on wetland habitats

For this analysis, Annex I habitats were classified into six groups: bogs and mires, calcareous fens, humid meadows, inland salt marshes, wet heaths and shrubs, and wet forests. From Figure 4.10 it can be seen that pressures related to agriculture are the most frequently reported group of pressures in all habitat types except wet forests. For humid meadows they represent 67 % of all pressures reported. Pressures related to forestry are the main pressure group for wet forests. Modification of water regimes (for purposes other than agriculture or forestry) is important especially for bogs and mires and calcareous fens and less important for inland marshes and wet forests. Urbanisation is the second most important pressure category for inland marshes. Invasive alien species were reported as a pressure especially for wet forests and wet heaths and shrubs, less so for inland marshes and humid meadows and rarely for calcareous fens and bogs and mires. Pollution is not a dominant pressure: it represents **up to 5 % of pressures reported for bogs and mires and wet heaths and shrubs**.

Figure 4.10 Main pressures for wetland habitats

Source: Article 12 and Article 17 Member States' reports and assessments.

At the level of individual pressures, the following pressures are most important for wetland habitats:

- **Abandonment of grassland management** (but not significant for wet forests and wet heaths and shrubs), **natural succession** resulting in species composition change (not important for inland salt marshes and wet forests), and **intensive grazing or overgrazing by livestock** (not important for wet forests) are overall the most important pressures.
- **Drainage** belongs to the group of top pressures for calcareous fens and for bogs and mires.
- The most important pressure on wet forests is **other invasive alien species** followed by **conversion to other types of forests** including monocultures and **modification of hydrological flow**.
- The top pressure for inland salt marshes is the **conversion from one type of agricultural land use to another**.

vegetation to improve water flow. Next, **physical alterations of water bodies** is one of the main pressures on freshwater fish. Physical alterations encompass, inter alia, the removal of sediments, building of dams and weirs, canalisation and water deviation.


Aside from freshwater habitats, severe pressures related to human-induced changes in water regimes also impact bogs, mires and fens and the species associated with them (e.g. Aquatic Warbler). **Drainage** activities reported under this category that are not attributed to a specific sectoral driver (e.g. agriculture) account for 10 % of the overall pressures reported for bog habitats. Consequently, these activities also affect rare and specialised species that are a part of bogs, such as the Moor Frog (*Rana arvalis*) or Peat Moss (*Sphagnum spp.*). Hydrological changes caused by drainage also affect birds such as herons and storks

that often prey on species dependent on such habitats. Other birds such as waders use wet grasslands for nesting (see Box 2.3). As a result, any changes in water levels may pose a significant threat to their breeding success.

4.1.8 Energy production

In 2018, the EU produced around 42 % of its own energy. Renewable energy (34 % of total EU energy production) was the largest source contributing to energy production in the EU at that time, followed by nuclear energy (31 %) and solid fuels (22 %) (Eurostat, 2020b). Renewable energy sources include solar (thermal and photovoltaic) energy, hydropower (including tide, wave and ocean energy), wind, geothermal and all forms of biomass energy. Although these renewable energy sources are highly important

Figure 4.11 Distribution of level 2 pressures caused by energy production for habitats and species, shown as the percentage of pressures within this level 1 group



Notes: The size of the squares and their shade reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages. Total number of reports is given in parentheses.

Source: Article 12 and Article 17 Member States' reports and assessments.

to mitigate climate change and thereby reduce negative impacts on biodiversity, their construction and operation nevertheless have an impact on habitats and species. In contrast to the pressures caused by renewable energy sources, the impacts of fossil fuel energy (oil, coal and gas) production are more dispersed: water-related impacts (e.g. lowering the water table) or the impacts of extraction activities and the operation of power plants are rather reported under level 1 pressure groups such as 'Extraction of resources' or 'Mixed source pollution'.

As noted in the previous section, the pressures related to human-induced hydrological impacts are dispersed among the different level 1 pressure groups. Therefore, this group reflects only 45 % of all energy-related pressures; most of the other pressures are classified as human-induced changes in the water regime (largely dams). Of all the energy-related pressures, **hydropower installations** represent the single most important impact for (migratory and freshwater) fish (Figure 4.11). This includes not only dams or weirs as physical structures but also the changes they cause in the hydrological functioning of rivers, river morphology (large stretches canalised) and the chemical and thermal properties of water. Aside from freshwater habitats that are exposed to hydropower installations (such as Alpine rivers and their ligneous vegetation with *Salix elaeagnos*, 3240), other habitats seem much less affected by energy production.

Pressures from **wind, wave and tidal power** are the dominant energy-related pressures for species. Wind power may have a significant impact, as was reported for bats such as the Lesser Noctule (*Nyctalus leisleri*), Nathusius's Pipistrelle (*Pipistrellus nathusii*), and the Paricoloured Bat (*Vespertilio murinus*) as well as birds such as the Razorbill (*Alca torda*), the Greater White-fronted Goose (*Anser albifrons albifrons*) or the Golden Eagle (*Aquila chrysaetos*). This is mainly due to direct collision and fatal barotraumas from the rotating blades of turbines. However, species can also be affected by displacement through habitat reduction and substantial habitat loss or damage, barrier effects that alter migration flyways or local flight paths, and other indirect effects, such as reductions in prey species (Arnett et al., 2008; Rollins et al., 2012; Gove et al., 2013; Garthe et al., 2017). The impacts of wind farms on birds are highly dependent on, for instance, species ecology and flight height. As an example, little direct impact has been identified for the gannet population, but large raptors and other large soaring species have a higher risk of collision (Furness et al., 2013; Gove et al., 2013; Warwick-Evans et al., 2017). Oceanic wave and tidal resources, while still few in operation, may result in the displacement

of marine birds from important feeding grounds either through direct disturbance of the birds themselves or through disruption to their prey as developments expand in the coming decades (BirdLife International, 2012). In addition, **offshore wind farms** and other types of energy production severely affect marine mammals because of the noise pollution caused during the construction process; affected species include the Harbour Porpoise (*Phocoena phocoena*) and the Grey Seal (*Halichoerus grypus*) (BfN, 2020).

EU energy production from renewable energy sources is expected to increase significantly over the coming decades, as it is a major pillar of fully decarbonising the EU economy by 2050. For 2030, the EU currently aims to increase the share of renewable energy in total final energy consumption to 32 % by 2030, an increase of 60 % compared with 2015 (EC, 2018). In its proposal for a new European Green Deal, the European Commission has proposed to increase the EU greenhouse gas reduction target for 2030 from currently -40 % compared with 1990 to a target in the range between -50 % and -55 % (EC, 2019a). If such an increase is adopted by the EU, renewable energy targets for 2030 would also have to increase. Therefore, it is fundamental that climate mitigation and biodiversity policies and measures are developed and implemented in a coherent and coordinated manner to avoid, or minimise, further impacts.

4.1.9 Climate change

Climate change is already happening, with noticeable impacts such as rising temperatures (in air, sea and freshwater), more frequent periods of drought and wildfires, shifting rainfall patterns, melting glaciers, less snow and a rising global mean sea level. These effects have both direct and indirect impacts on species and habitats. Direct impacts include changes in phenology, species abundance and distribution, community composition, habitat structure and ecosystem processes, and the desynchronisation of ecological relationships (EEA, 2017; IPBES, 2019). Other impacts include northwards and uphill range shifts, as well as local and regional extinctions of species (Keller et al., 2020).

In the current reporting period, the most relevant pressure related to climate change was **droughts and decreases in precipitation** (see Figure 4.12). This pressure accounts for 5 % of all reported pressures affecting amphibians (e.g. the Yellow-bellied Toad (*Bombina variegata*)). Amphibians are particularly sensitive to **temperature** and **changes in precipitation** because of their central

position in food webs, their strong dependence on both aquatic and terrestrial systems, and their moist permeable skin acting as a sensitive respiratory organ (Olson and Saenz, 2013). Molluscs and some mammals, e.g. bats such as Botta's Serotine (*Eptesicus anatolicus*), are affected by rising temperatures, as this hampers their reproduction and foraging. Drought and decreased precipitation make up 2 % of the reported pressures for birds associated with reedbeds and reedy ponds, such as the Common Little Bittern (*Ixobrychus minutus*).

Several habitats also face severe pressures from decreases in precipitation, such as bogs, mires and fens. Coastal habitats, such as those in the Atlantic and Boreal regions, mainly face the pressure of **changes in sea level and wave exposure**. Although other climate-related pressures may not have been

reported as high-level pressures in many instances, the **decline or extinction of related species** (e.g. prey species) or the **change in habitat locations** (e.g. for coastal and pelagic seabirds) already indicate a significant impact on the reproduction and survival rates of species and bird populations.

Although climate change is not reported as a particularly relevant pressure for the period from 2013 to 2018, research on future scenarios predicts that climate change will have a dramatic effect on European plants and animals in the years to come and lead to an acceleration in biodiversity loss in many areas (EC, 2020f). Climate change is also likely to exacerbate the problem of invasive species in Europe (especially in the northern parts), and some locations will become more favourable for previously harmless alien species.

Figure 4.12 Distribution of level 2 pressures caused by climate change for habitats and species, shown as the percentage of pressures within this level 1 group





Figure 4.13 Distribution of level 2 pressures caused by pollution for habitats and species, shown as the percentage of pressures within this level 1 group



Notes: The size of the squares and their colour reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages. Total number of reports given in parenthesis.

Source: Article 12 and Article 17 Member States' reports and assessments.

Reflecting these predicted changes, climate change is seen as an emerging threat. This is also mirrored in Member States' reports under the nature directives. In this context, threats are defined as projected pressures within 12 years of the end of the current reporting period. For climate change, the reporting suggests a steep rise (80 %) in related pressures, further amplifying the present impacts of droughts and decreases in precipitation as well as changes in temperature. In addition, changes in habitat location, size and/or quality is a rising threat with particular relevance for species, most notably in the Continental and Alpine regions. According to Member States' reporting, water-dependent species and habitats will be specifically affected, such as waterbirds (e.g. geese, diving ducks (*Aythya spp.*), herons), freshwater species such as fish (e.g. Grayling (*Thymallus thymallus*) and Atlantic Salmon (*Salmo salar*)), amphibians, butterflies, crayfish and the Pearl Mussel (*Margaritifera margaritifera*) and diverse habitats such as forests (e.g. several temperate beech and oak forests and riparian mixed forests), coastal and marine habitats (e.g. coastal lagoons, mudflats, reefs), glaciers, and alkaline fens, petrifying springs and quaking bogs.

4.1.10 Pollution

The number of pollution records from the level 1 group 'Pollution' was relatively few (see Figure 4.2). However, this group includes only pollution from mixed sources that could not be attributed to a specific origin or activity (see Figure 4.13).

Considering the pollution entries reported across different level 1 groups reveals the true importance of this pressure. These pressures account for 7 % of the over 31 000 individual pressure records. The top three level 1 groups under which pollution pressures were reported are 'Agriculture' with almost half of the records (48 %), 'Mixed source pollution' (28 %) and 'Urbanisation' (21 %). This clearly shows the importance of agricultural activities as a key source of pollution (air, water and soil) negatively affecting the status of and trends in many habitats and species. Because of the taxonomy of pressures used in the reporting, atmospheric emissions and air pollution did not stand out and were underestimated; however, they have a significant impact in terrestrial habitats especially reactive to nitrogen deposition. Across Europe it is predicted that, with current policy in 2020, over 70 % of the area of EU ecosystems will receive

more atmospheric nitrogen than they can take and continue to function naturally (Hettelingh et al., 2017).

As a policy response to the critical issue of pollution, both the EU Biodiversity Strategy for 2030 and the

Farm to Fork strategy aim to significantly reduce the input of chemical pesticides (50 % reduction) and to promote less intensive farming practices, including a significant reduction in fertiliser use (at least 20 %).

Key messages

- Agricultural activities represent the most common pressure group across habitats and species: abandonment of extensive management and intensification are the most frequent pressures.
- Agriculture is also the main sector contributing to air, water and soil pollution (almost half of all pollution reports), with significant impacts on standing waters, rivers and marine habitats and their species.
- Marine and coastal habitats are particularly affected by pressures related to sports, tourism and leisure activities.
- Species exploitation is the largest overall pressure for wintering and passage birds, relating to illegal shooting or killing and hunting, as well as incidental killing; non-bird species are also affected, with fish, mammals and reptiles being affected more than average.
- Invasive alien species affect habitats more than species, but they do affect amphibians, fish, vascular plants and breeding seabirds.
- Modifications of the hydrological flow are the most significant water regime modification pressure for European freshwaters and bogs, mires and fens, while physical alterations of water bodies are the dominant pressure for freshwater fish.
- Of energy-related pressures, hydropower installations are the most damaging for (freshwater) fish; most birds are especially vulnerable to electricity and communication transmission infrastructure.
- The most relevant pressure related to climate change is droughts and decreases in precipitation, accounting for 5 % of all reported pressures affecting amphibians; climate change is growing in importance and Member States reported significant increases in threats.



DASHBOARD
with detailed information by Member State:
Pressures Article 12



DASHBOARD
with detailed information by Member State:
Pressures Article 17


4.2 Conservation measures

4.2.1 Application of measures

The nature directives emphasise the need for conservation measures to maintain or restore the natural habitats and populations of wild fauna and flora. Establishing conservation measures in Natura 2000 sites is a mandatory responsibility of the competent authorities in each Member State⁽³⁰⁾ and is the focus of this chapter. If it is necessary to set up a formal framework for implementing practical conservation actions (conservation measures) — which represent the core of this section — Member States are asked to (1) develop appropriate management plans specifically designed for their sites or integrate them into other development plans, or (2) take appropriate statutory, administrative or contractual measures corresponding to the ecological requirements of the Annex I habitats and Annex II species present on the sites. For Annex I birds and other migratory birds for which the Natura 2000 sites are classified, Member States have to avoid pollution and deterioration of habitats or any disturbances affecting them.

Although the directives require conservation actions to be established within the Natura 2000 sites, conservation measures are also applied by Member States outside the network to contribute to achieving good conservation status of habitats and species in need of management. However, as Figure 4.14 shows, only a small number of measures are only taken outside Natura 2000 sites. Instead, most conservation measures are applied both inside and outside the Natura 2000 network.

Figure 4.14 Main location of measures for habitats and species




Source: Article 12 and Article 17 Member States' reports and assessments.

Within the nature directives' 2013-2018 reporting period, Member States provided an overview of conservation measures, indicating whether or not the majority of the measures needed were taken. For each habitat and species, this consisted of an overarching evaluation of the status of the main conservation issues targeted by the measures in place and of the expected overall response time of all the measures taken. This broad evaluation did not look specifically at the status, purpose or response time of each individual measure in place but was a global assessment of a habitat or species. This enables the link to be established between the status of measures and the reported conservation status or trends. In addition, Member States provided a detailed list of measures taken, which serves to highlight what has been done to counteract pressures and threats and to conduct an overall evaluation of the state of conservation measures.

Habitats

- Across terrestrial and marine habitats, the majority of conservation measures are already taken for around 60 % of Member States' habitats (Figure 4.15). Another 10 % of habitats were reported as not requiring any conservation measures.
- Forests, followed by grasslands, freshwater and coastal habitats more frequently have all measures needed in place in comparison to other habitat groups. Eight Member States (Bulgaria, Denmark, Estonia, Italy, Luxembourg, Latvia, Slovakia and the United Kingdom) report that, for more than 90 % of their habitats, conservation measures have been identified and taken.

Figure 4.15 Implementation status of habitat conservation measures



Source: Article 12 and Article 17 Member States' reports and assessments.


⁽³⁰⁾ Reporting on conservation measures does not cover all species listed in the nature directives; this information is available only for Annex II species from the Habitats Directive, Annex I birds and birds for which Natura 2000 sites are classified under the Birds Directive.

- Habitats for which the necessary measures have been identified but most of them not yet taken comprise an additional approximately 22 % of reports. These largely stem from coastal, freshwater and dune habitats and the largest share is in the Marine Mediterranean region, followed by the Mediterranean and Boreal regions.
- For those habitats where particular conservation actions were reportedly not needed, the highest proportion is in Cyprus followed by Greece and Romania.

Member States have different underlying reasons for applying conservation measures, including **restoring** the current status — increasing the habitat area, restoring its structures and functions — or just **maintaining** the current status. For the large majority of habitat reports, 'measures taken' aim to maintain the current status, followed by restoring structure and functions (72 % and 23 % respectively); only around 4 % of the reports have measures aiming to increase the habitat area.

In relation to measures taken and their main purpose, the habitats reports are also classified according to the time-frame of the expected response of a habitat to measures applied. The majority of habitats will respond to the measures in the mid-term (68 %) and the rest either in the short term (14 %) or the long term (18 %). These categories were all present in each of the habitat groups but varied in their frequencies. Short-term responses to measures are expected in particular for dune habitats and heath and scrub, mid-term responses are generally high across all habitat groups, with freshwater habitats having the highest

Figure 4.16 Implementation status of species conservation measures



Source: Article 12 and Article 17 Member States' reports and assessments.

share. Long-term responses are expected for forests in particular.

Species

For approximately half of the birds and non-bird species, measures were 'identified and taken' and one third have 'identified but not yet taken' the necessary measures (see Figure 4.16). The share of these categories is similar across marine and terrestrial species.

Conservation measures relating to **non-bird species** revealed several noteworthy findings across the different categories:

- Measures 'identified and taken' have the highest reported shares in Belgium, Bulgaria, Luxembourg, Slovakia and the United Kingdom at around 90 %; the majority of necessary measures were taken for vascular plants, arthropods, mammals and fish, with around 80 % aiming to maintain the current status.
- Species with measures 'identified but not yet taken' make up the largest share of the Macaronesian region species; the highest total counts were reported by Croatia, Spain and Portugal.
- 'Measures not needed' comprise around 70 % of the Marine Black Sea reported measures for species; the largest share of cases was reported by Cyprus.

Almost 80 % of the measures taken for non-bird species aim to maintain their current status. Restoration measures, on the other hand, play a less prominent role. The majority of measures are estimated to have short-term effects (60 %). The highest share of those with long-term responses are reported for fish, mammals, molluscs and non-vascular plants.

Regarding **birds**, 'measures not needed' account for around 30 % of all cases. The remaining species are largely reported as having the measures 'identified and taken' (40 %) or 'identified but not yet taken' (30 %):

- Cases with measures 'identified and taken measures' were largely reported from Bulgaria and Poland (more than 160 measures each) as well as Spain, Germany, Hungary and the United Kingdom (more than 100 measures each) and comprise very high shares of the measures reported from Hungary, Malta and the United Kingdom (approximately 90 %).

- Cases with measures 'identified but not yet taken' were largely reported from Croatia.
- Only a few bird species have measures that are 'needed but cannot be identified', comprising a large share of Greece's reported measures. The share of response times of these measures is around 60 % mid-term, 25 % short-term and 15 % long-term.

Among breeding, wintering and passage birds, the highest total number of measures was reported for breeding birds. The distribution across categories is similar for all three groups and is, in decreasing order: maintain current status, increase area, restore structure, expand range. Malta reported almost 100 % of the measures to increase area as taken, while Estonia reported more than 70 % of the measures to enhance the current status by restoring structure and functions as taken.

Key messages

- For most of the habitats and species, measures have been applied both inside and outside the Natura 2000 network, with a smaller share taken only within the network. Measures that have been taken only outside the Natura 2000 sites are limited.
- For 60 % of habitats, measures were 'identified and taken'; these mostly aim to maintain the habitat's status (e.g. for forests, grasslands, freshwater and coastal habitats).
- For non-bird species, around 50 % of the reports indicate that measures have been taken; these largely target vascular plants, arthropods, mammals and fish, with the majority of measures aiming to maintain their status.
- For birds, approximately 40 % of the reports indicate that measures have been taken, with the highest total share reported for breeding birds.
- Overall, the majority of habitats and bird or non- bird species have measures that have been taken to maintain the current status.
- Reports indicate that, for the majority of habitats and species, measures are needed and these had been taken; however, conservation status and trends have not significantly improved and deterioration is ongoing. This may indicate that measures not yet taken are key to delivering conservation objectives and also that measures not fully implemented either need to be scaled up or were not effective.



DASHBOARD
with detailed information by Member State:
Measures Article 12



DASHBOARD
with detailed information by Member State:
Measures Article 17

4.2.2. Measures and pressures


The following section looks at the measures reported that aim to reduce the pressures reported in Section 4.1 to minimise potential impacts from future threats. Corresponding to the pressures, measures are reported for different level 1 groups that (mainly) mirror the level 1 pressure groups.

As shown in Figure 4.17, conservation measures targeting agricultural practices are by far the most

frequent among the measures implemented, corresponding to the negative impacts of agricultural activities, or the decline of extensive management of grasslands and some heathlands. However, measures are more evenly distributed across the categories than pressures targeting a wide range of different sectors.

As for the reporting of pressures, practical conservation measures are further detailed in a second level. The following section therefore reflects measures that are applied to target the pressures listed in Section 4.1.

Figure 4.17 Distribution of level 1 measure categories among habitats and species



Notes: The size of the squares and their shade reflect the percentage of pressures for each group: bigger darker squares indicate higher percentages.

Source: Article 12 and Article 17 Member States' reports and assessments.

Agriculture

With more than 17 % of all measures, agriculture-related activities were the most frequently reported group. Most of the conservation measures applied for habitats concern **adapting mowing and grazing activities, maintaining existing extensive agricultural practices** and **reinstating appropriate practices to address abandonment**. Semi-natural grasslands and especially hay meadows, such as *Molinia* meadows (6410) or lowland hay meadows (6510), are the habitats that are most frequently addressed by such measures.

For freshwater habitats, the majority of measures aim to **reduce diffuse pollution to surface waters and groundwaters**, e.g. through decreased discharge, connection to sewers and improvements in waste water treatment. Bogs, mires and fens, however, are mainly addressed by **management of drainage and irrigation operations** to improve habitat condition, together with **reinstating or adapting extensive management practices**. In addition, airborne pollution caused by agriculture was targeted by a number of measures, e.g. by integrated nitrogen management approaches, low-emission application techniques or replacing chemical fertilisers with organic equivalents.

For most species' groups, **reinstating, adapting and maintaining extensive management activities** were also the conservation measures most frequently reported, especially for insects, vascular plants and birds. However, with 6 % of all reported measures taken for mammals, **managing the use of natural fertilisers and chemicals** is the predominant measure to address the related pressures reported mostly for bats and small mammals.

Among a wide variety of measures applied, Member States should also provide specific information on measures aiming to **restore or re-create the landscape features of agricultural landscapes** or habitats. This type of measure is implemented relatively rarely for most of the habitat or species groups, but these proactive conservation actions are rather frequent for agricultural and dune habitats, reptiles, amphibians and mammals.

Urbanisation

Overall, **reducing the impact from outdoor sports, leisure and recreational activities** is the most frequent measure related to this category targeting habitats and species. Such reductions particularly target marine and coastal habitats, dune habitats and rocky habitats

Box 4.3 Closer look: Pastures as ecological corridors for mammals



Photo: Corsican Red deer © NEEMO

The Corsican Red Deer (*Cervus elaphus corsicanus* Erxeleben) is a subspecies of the European red deer and is endemic to Sardinia and Corsica. The species is mainly threatened by extensive logging, hunting, forest fires, the diffusion of farming and the spread of livestock. These factors have led to habitat fragmentation and severe population decline across the deer's territory. The LIFE project One Deer Two Islands (LIFE11 NAT/IT/000210) aimed to improve the conservation status of the Corsican red deer in Sardinia and Corsica by reducing conflict with human activities and promoting awareness of the species' ecological and economic value. Eighty deer captured in Sardinia in the source area of the Costa Verde Oasis (Arbus) at the Monte Arcuentu and Rio Piscinas Site of Community Interest were reintroduced into the former

Sardinian province of Ogliastra to repopulate a territory that in the past housed a large number of red deer. Ecological corridors have been set up by creating 80 ha of pastures and herbaceous meadows to feed the deer in south-east Sardinia. The pasture and meadows keep deer away from crops that are intended for domestic livestock and encourage them to migrate to less populated areas. As the project only ended in 2019, the impact cannot yet be measured. Considering the endemism characterising the species, it is highly probable that the project has contributed to the recent positive trend and a good conservation status.

(predominantly in the Alpine region) and species such as mammals (e.g. Greater Mouse-eared Bat (*Myotis myotis*)), birds (e.g. Golden Eagle (*Aquila chrysaetos*)) and vascular plants. One of the main focuses of these measures is to develop sustainable tourism.

Measures related to **managing the conversion of land for construction and development of infrastructure** are frequently applied to grasslands and related species such as insects, birds and reptiles.

Proactive measures such as **restoring the areas impacted by urbanisation or leisure use** are predominantly implemented for marine and coastal habitats. For dunes, these measures may include controlling visitor numbers and raising awareness of their value.

Forestry

Overall, the most common forestry measures relate to **adapting and changing forest management and exploitation practices** (25 % of all forestry measures taken). Not surprisingly, this type of measure particularly targets forest habitats. In practice, these measures can involve, for example, adapting and changing management practices to secure or develop

old stocks of trees, retaining dead and dying trees and stumps, preserving habitat continuity or preventing forest wildfires. One of the prominent approaches to increasing the multifunctionality of forests is the 'close-to-nature-management' that promotes alternatives to even-aged monocultures.

Preventing the conversion of (semi-) natural habitats into forests and (semi-) natural forests into intensive forest plantation is of particular relevance for many habitat groups and some specific non-bird species, each of which have close to 20 % of all forestry-related pressures falling within this category. Examples include grasslands, heath and scrub, dune habitats and bogs, mires and fens, as well as species such as insects and vascular plants.

The species targeted by forestry measures are largely mammals, insects and vascular plants as well as forest-dependent birds such as the Hazel Grouse (*Bonasa bonasia*) or the Black Stork (*Ciconia nigra*).

Exploitation of species

Measures in this group are most frequently implemented for fish and birds as it includes, 16 % and 14 %, respectively, of all measures taken for fish

Box 4.4 Closer look: Tackling invasive alien species in England



Photo: Japanese Knotweed © MdE

(Himalayan Balsam and Japanese Knotweed) and the use of male sterilisation to control signal crayfish. Regarding the actual management and control of IAS dispersal, RAPID LIFE implemented demonstration projects at catchment and local scales. At catchment scale 12 000 m² and 8 000 m² of Himalayan Balsam was removed manually by volunteers, while, at the small scale, biocontrol methods were used to clear 100 m² of Himalayan Balsam and 300 m² of Japanese Knotweed. Biocontrol methods were used to combat Himalayan Balsam at nine sites and Japanese Knotweed at 13 sites. The project also successfully combatted Signal Crayfish at three sites and released White-clawed Crayfish reared in captivity to restore the population balance.

The LIFE project Rapid Life (LIFE16 NAT/UK/000582) aims to deliver a package of measures to reduce the impact and spread of invasive alien species (IASs) in freshwater aquatic, riparian and coastal environments across England. It addresses different pillars of the EU IAS Regulation, such as prevention, early detection and response to and management of widely spread IAS species at the regional level. The project produced IAS management toolkits and protocols for the prevention, detection and control of IASs. Moreover, the project successfully implemented and demonstrated novel methodologies for the detection and monitoring of IASs, such as the use of e-DNA to detect signal crayfish, and innovative technologies for the removal and/or eradication of IASs, such as the use of novel biocontrol methods to eradicate invasive weeds

and birds. For birds, 40 % of these measures include the **management of hunting, recreational fishing and recreational or commercial harvesting or collection of plants**, which is also the most frequent measure taken for fish. Other species that are targeted by measures relating to their exploitation are mammals and reptiles, especially through the **control/eradication of illegal killing, fishing and harvesting** — mostly achieved by increased enforcement. This measure is also frequently applied in the conservation of fish and birds.

Invasive alien species

Around 9 % of all measures reported were implemented to **reduce the impacts of invasive alien or problematic native species**. Habitats, especially coastal habitats, dunes, forests, grasslands and freshwater habitats, and equally non-bird species, are mainly targeted by measures related to controlling and eradicating invasive alien species, whereas measures for birds mostly involve managing problematic native species. Practical implementations include the elimination of key invasive species such as Himalayan Balsam (*Impatiens glandulifera*) or Japanese Knotweed (*Fallopia japonica*) or the control and containment of invasive animal species.

Natural processes

Natural process-related measures largely target habitats through their **management to slow, stop or reverse natural processes**. Such measures are typically applied to bogs, mires and fens as well as to forests and grasslands.

Mixed source pollution and the modification of water regimes

This group of measures, targeting in particular the aquatic environment and wetlands, contains the measures overall most frequently applied for the conservation of fish populations. Here, the main focus lies on **reducing the impact of multipurpose hydrological changes**, which is the single most often applied measure, of all measures, dedicated

to conserving fish. Direct measures may include the removal of sediments, canalisation, water deviation or modification of the flooding regime. These measures are also frequently applied for the conservation of birds, for example wading birds such as the Eurasian Bittern (*Botaurus stellaris*).

Moreover, measures from this group are equally important for freshwater habitats and bogs, mires and fens that are extensively targeted by **reducing the impact of multipurpose hydrological changes** and **mixed source pollution** as well as by **restoration measures**. This includes renaturalising rivers from plain to montane levels with *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation (3260) (see Box 4.5) or oligotrophic to mesotrophic standing waters with *Littorelletea uniflorae* and/or *Isoëto-Nanojuncetea* vegetation (3130). Measures related to mixed source pollution address emissions to air of ammonia as well as greenhouse gas and water-borne pollution by local action and regional policy implementation.

Energy production and resource extraction

For species, the most frequent measures addressing energy- and extraction-related pressures are **reducing the impact from hydropower operation and infrastructure**. This includes, inter alia, building and managing fish passages or regulating water flows, which is particularly relevant for fish such as the Asp (*Aspius aspius*). On the other hand, **adapting and managing renewable energy installation, facilities and operation** is most relevant for birds and aquatic mammals such as the Harbour Porpoise (*Phocoena phocoena*) or the European Otter (*Lutra lutra*) as well as bat species. Birds such as the White Stork (*Ciconia ciconia*) are further targeted by measures aiming to **reduce the impact of service corridors and networks**, such as by modifying power lines to prevent birds being electrocuted or developing green or blue infrastructure to improve connectivity.

Measures that focus on **adapting and managing the extraction of non-energy resources** are especially applied to coastal habitats. These measures target,

Box 4.5 Closer look: Increasing river connectivity in Estonia

Since 2012, Estonia has reported genuine stability in the unfavourable-inadequate conservation status of water courses from plain to montane levels with *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation (3260). This is in part thanks to the contribution of two consecutive LIFE projects HAPPYFISH (LIFE07 NAT/EE/000120) and LIFE HAPPYRIVER (LIFE12 NAT/EE/000871), which both restored habitats affected by hydrological changes by physically removing sediments in the water and managing riparian vegetation. HAPPYFISH was awarded Best of Nature in 2013.

for example, the extraction of minerals such as sand, gravel, loam, clay, peat and salt.

Climate change

Conservation measures targeting the effects of climate change involve either mitigation or adaptation activities. Of the reported measures that have been taken in this group, the majority addressing mitigation are reported for birds. However, measures focusing on adaptation are largely reported for habitats, including forests, freshwater and coastal habitats as well as bogs, mires and fens, e.g. by adapting

fire management or by adaptive management and restoration to strengthen their resilience and enhancing their heterogeneity and protection against natural hazards (through genetic diversity or adapted tree composition). In the Netherlands, for instance, the Greater White-fronted Goose (*Anser albifrons*), the Great Bittern (*Botaurus stellaris*) and the near-threatened Corncrake (*Crex crex*) are targeted with so-called 'climate buffers' as part of a landscape management approach, including restoring overflow areas and reed marshes in the alluvial plain and broadening tributary river mouths to slow outflow into the river (BirdLife International, 2009).

Key messages

- 17 % of all measures target agricultural activities, most prominently by reinstating, adapting and maintaining extensive management activities such as mowing and grazing.
- Measures for marine and coastal habitats, dune habitats and rocky habitats most frequently relate to reducing the impact from outdoor sports, leisure and recreational activities.
- Most common forestry measures relate to adapting and changing forest management and exploitation practices, which also relates to climate change adaptation.
- The management of hunting and recreational fishing and the control/eradication of illegal killing and fishing mainly target birds and fish. Overall, the most frequent measure targeting the conservation of fish, however, is reducing the impact of multipurpose hydrological changes.

4.2.3 Effects of measures

Linking status and trends to measures


A link between the measures taken and good conservation status is apparent for nearly all **habitat groups**, with a statistically significant positive correlation for rocky habitats and sclerophyllous scrub (see Figure 4.18). These two groups include many natural habitats in which the main conservation challenge is to regulate or avoid human activities and also habitats requiring agricultural use. Conservation measures applied to these habitats include regulating and maintaining extensive agricultural use, regulating leisure activities, regulating extraction activities, reducing the impact of the transport infrastructure, and habitat restoration. Raised bogs, mires and fens, however, show the opposite tendency. Only about 5 % of bogs in which measures have been taken are reported to have a good conservation status; this is significantly lower than for bogs where no measures

have been taken. In general, as seen in Section 4.1, the majority of conservation measures taken during the reporting period will only achieve the expected result in the near future. This may be one of the reasons for the missing correlation between measures and status for these habitat groups, but it is also an indication that further restoration effort is needed (see Section 4.3). Similarly, the majority of habitats show improving and stable trends as a result of the measures taken.

Transboundary pressures, such as air pollution, affect habitats particularly sensitive to nitrogen deposition; however, this kind of pressure cannot be addressed at local level and may explain the poor response of those habitats to restoration measures.


(Non-bird) species also reflect this correlation, having a higher likelihood of having a good conservation status when conservation measures are implemented, often just by maintaining their good status from the

Figure 4.18 Percentage of good conservation status if measures are taken or not taken for Annex I habitats



Source: Article 17 Member States' reports and assessments.

Figure 4.19 Percentage of good conservation status if measures are taken or not taken for Annex II species



past. Approximately 30 % of species with a good status have benefited from such measures, while only 20 % of species for which no measures were implemented (although they are needed) have a good status (see Figure 4.19). Similarly, over one third of species that did not have conservation measures in place have a bad conservation status, compared with only one quarter of those species for which measures were applied. The strongest correlation between implementing conservation measures and status is observed for mammals, arthropods, reptiles and non-vascular plants as well as, to a lesser degree, fish. The proportion of deteriorating trends in conservation status is significantly higher for almost all species groups if measures are not taken, with the exception of amphibians and molluscs.

For **birds**, Annex I taxa and special protection area trigger species are more likely to have increasing population trends when conservation measures are implemented; this is true for all bird groups, including breeding, wintering and passage birds (see Figure 4.20).

Linking improvements to measures


Looking only at the link between measures and the status of Habitats Directive species and habitats does not show the full picture, especially as the changes (represented as a change between either conservation status categories or conservation

status trends) that may result from implementing conservation measures are not taken into account. In addition, many habitats and species populations are targeted by measures as a result of their bad conservation status. Finally, measures sometimes need longer time-frames to take effect. This is the case with, for example, degraded bogs, mires and fens where — depending on the hydro-ecological situation before restoration and the methods applied — it may take several decades before the desired peat-forming vegetation is sustainably restored. The following section therefore looks at these changes, building on the approaches taken to identify improvements for habitats and non-bird species (Section 3.4) and for birds (Section 2.4).

Applying conservation measures increases the likelihood of having a positive trend in conservation status (improved status as well as improving trend; both are grouped under the 'improvement' category in this analysis). However, improving cases for habitats are rather rare, as shown in Section 3.4.2. In cases where measures are needed but not taken, the conservation status and/or trend is more likely to be bad or further deteriorating (Figure 4.21).

For all improving habitat assessments, more than 80 % of the improvements can be linked to the measures taken. However, habitats showing a deteriorating status despite the application of measures may indicate that

Figure 4.20 Percentage of increasing population trends if measures are taken or not taken for Annex I and special protection area trigger bird species



Source: Article 12 Member States' reports and assessments.


the measures taken have not been effective or require more time for their impact to be seen.

It can be expected that, for at least some habitat types that were previously degraded or depleted, the positive trends will be more frequently associated with restoration measures (if the response time is long enough) than with measures aiming to maintain the current state.

This correlation between restoration measures and positive trends is evident in the data reported for certain habitat groups and locally, for example:

- bogs, mires and fens (mainly in Belgium and the United Kingdom), e.g. through buying out an industrial peat extraction site for restoration in the case of the United Kingdom;
- coastal habitats, e.g. sandbank (1110) restoration in Denmark and the Netherlands;
- freshwater habitats by restoring water courses from plain to montane levels with *Ranunculon fluitantis* and *Callitricho-Batrachion* vegetation (3260) in Estonia or Germany (see also Box 4.6);
- heaths, e.g. through restoration activities in Latvia (4010);

Figure 4.21 Measures reported and changes in conservation status of habitats



Source: Article 17 Member States' reports and assessments.

- grasslands, e.g. through proactive management in Belgium and the Netherlands;
- forests, mainly various beech forests in Austria, Belgium and the United Kingdom.

There are limited cases of **Annex II** species having had improvements to their conservation status reported (see Section 3.4.3). Similar to the findings for habitats, the data show that applying measures correlates with a higher likelihood of having good or improving conservation status. Around 80 % of the improvements identified can be linked to the measures taken (see Figure 4.22). In cases where measures are 'needed but not taken', conservation statuses are more likely to be poor, bad or further deteriorating.

Of the various species groups, mammals and fish have shown the most frequent improvements in conservation status as a result of measures taken. A link between restoration measures (mainly promoting the population growth or restoring the species' habitats) and positive trends in global conservation status assessments has been recorded for some species or species groups, for example:

- large carnivores such as the Brown Bear (*Ursus arctos*), Grey Wolf (*Canis lupus*), Iberian and Eurasian Lynx (*Iberian lynx* and *Lynx lynx*) in several Member States with depleted populations;

Box 4.6 Management measures to improve the hydrological regime in restored freshwater habitats in Germany (3260)




Photo: *Ranunculus fluitans* © Rutger Barendse

Various LIFE projects contributed to this success, e.g. Bachtäler Arnsberger Wald/Rehabilitation of streams in the Arnsberger Wald (LIFE07 NAT/D/000214) and Rur und Kall/Fluvial habitats (LIFE10 NAT/DE/000008).

Source: Tucker et al. (2019).

The watercourse habitat from plain to montane levels with *Ranunculus fluitans* and *Callitriche-Batrachion* vegetation in Germany was widely assessed as being in bad condition. Measures to improve the status of the watercourses and their species included improving and restoring the water quality and the hydrological regime and establishing protected areas. The measures taken to improve the status of the habitat had wider positive effects and enabled the populations of the associated species to increase across Germany. In addition to reducing habitat-related pressures on the species, measures were taken to regulate fishing practices. The management recommendations were drawn up by the Federal Agency for Nature Conservation for specific endangered species.

Figure 4.22 Measures reported and changes in conservation status of Annex II species



Source: Article 17 Member States' reports and assessments.

- several bat species, mainly in Belgium and France;
- among plants, such as the Fen Orchid (*Liparis loeselii*) in Austria, Belgium and the United Kingdom;
- insects, e.g. saproxylic beetles in Italy;
- migratory fish, including anadromous fish and lampreys such as the Atlantic Salmon (*Salmo salar*).

Fish and mammals also respond to both restoration and maintaining measures, whereas the improvements among arthropods, molluscs and vascular plant groups are more frequently associated with measures aiming to maintain the current state. For amphibians, non-vascular plants and reptiles, the opposite is true, and improvements are more frequently connected to restoration measures.

The share of targeted measures applied for **birds** is around 50 % (20 % to restore and 31 % to maintain), which is significantly lower than that for habitats and non-bird species. Overall, the link between the trend category and the implementation of measures is less evident. However, as is the case for Annex II species, most of the measures are expected to have an impact in the mid-term (i.e. by 2030) rather than in the short term.


Box 4.7 Closer look: Recovery of the Aquatic Warbler

Photo: Aquatic warbler © Dušan Boucný, IUCN

The Aquatic Warbler (*Acrocephalus paludicola*) is an Annex I species for which an improved conservation status was reported, moving from vulnerable in 2015 to near threatened in the most recent reports. Once widespread, the aquatic warbler almost entirely disappeared from western Europe as fen, mire and wet meadow habitats were lost to agriculture. Since 2011, however, the population has started to stabilise. These status improvements can be traced back to targeted conservation actions, including LIFE projects (e.g. the Aquatic Warbler project — Conserving *Acrocephalus paludicola* in Poland and Germany, LIFE05 NAT/PL/0001016, and LIFE Aquatic warbler and biomass, LIFE09 NAT/PL/000260), an international species action plan (2008), the international memorandum of understanding on

the species under the Bonn Convention for Migratory Species in 2003, and protection under the Natura 2000 network. While most countries with low Aquatic Warbler numbers reported decreasing trends, Poland reported an increase in their population following intensive conservation actions. In the case of the Polish LIFE project, the area of habitat suitable for the aquatic warbler in eastern Poland was increased and ongoing management of major parts of the project sites was secured through income generated from biomass. This suggests that the species is highly dependent on conservation management.


Source: Polish Society for the Protection of Birds (2020).

Although the link between increased trends and measures is not always clear, decreasing trends are more likely when measures have not been taken (see Figure 4.23). In terms of increasing population trends, there is a link at Member State level for some species groups:

- For raptors and owls, for example, more than half of the taxa with an increasing population trend at Member State level (winter and breeding) have had measures taken.
- for seabirds from the petrel, storm-petrel and shearwater families (Procellariiformes), a large part of the increasing trend is associated with the measures taken.

Further differences are evident based on the type of measure taken, i.e. restoration or **maintenance** measures. Some groups benefited noticeably more from a given type more than others. **Maintenance** measures benefited, for example, ducks, geese and swans (Anseriformes), stork-like birds (Ciconiiformes), loons (Gaviiformes), passerines (Passeriformes) and grebes (Podicipediformes) more than restoration measures. Hoopoes and hornbills (Bucerotiformes), bustards (Otidiformes) and the colourful Coraciiformes families, on the other hand, benefited more from restoration measures.

Figure 4.23 Measures reported and changes in bird population trends (Annex I and special protection area trigger species)



Source: Article 12 Member States' reports and assessments.

Key messages

- For most habitats and species, there is a positive link between the measures taken and good conservation status (except for raised bogs, mires and fens, vascular plants, amphibians, molluscs, and passage birds).
- Applying proactive restoration measures (e.g. to restore structure and functions) and measures to maintain the current status drives improvements in status.
- Restoration measures strongly benefited bogs, mires and fens, heath and scrub, freshwater habitats, grasslands and forests (mostly locally) as well as amphibians, reptiles and birds of the hoopoe and bustard groups.
- Fish and mammals benefited equally from both restoration and maintenance measures.
- Maintenance measures are key for arthropods, molluscs and vascular plants and strongly benefited ducks, geese and swans, storks and flamingos, loons and passerines.

4.3 Restoration needs of habitats

The conservation status of a habitat depends on its natural range, area, structure and functions, and future prospects (see more detail on parameters in Box 3.1). While restoration efforts need to address all of these parameters, this section focuses only on Annex I habitat types and the restoration efforts that are needed to re-establish their natural range, increase their surface area and improve their structure and functions (including within the Natura 2000 network). In the context of this report, restoration has a dual meaning and refers to:


- **improving** the condition (used to assess structure and functions) of existing habitats through targeted management measures (e.g. for overgrazed grassland with surplus nitrogen in the soil and reduced floristic composition);
- ensuring sufficient availability of habitat area through habitat **(re-)creation** (i.e. creating additional areas of a given habitat, e.g. wetland habitat from a drained agricultural field).

This section estimates the minimum area that needs to be restored to achieve favourable conservation status for the habitats listed in Annex I of the Habitats

Directive. It should be kept in mind that restoration alone is, however, not sufficient and that future prospects (i.e. pressures and measures) also need to be favourable to achieve the desired effects.

4.3.1 Overall need for restoring EU habitats

The analysis shows that at least 11 000 km² of Annex I habitats **need to be (re-)created** and added to the habitat area that already exists to ensure the long-term functioning of each habitat. This figure was estimated based on the minimum possible area required to achieve the status of a favourable reference area reported by each Member State (for more details, see Röschel et al., 2020), and it is likely that more will be needed to achieve good status for all habitats. It should be noted that the favourable reference area (and the condition of habitats used for the estimation of improvement needed below) is largely missing from several Member States — particularly for marine habitats — which might lead to a considerable **underestimation** of the actual area needing to be restored (see Figure 4.24). The habitat groups identified as having the largest areas to be (re-)created are forests (4 600 km²), grasslands (1 900 km²), bogs, mires and fens (1 700 km²), and coastal habitats (1 400 km²). EU-wide, this corresponds to 1-1.5 % of the total existing area of these habitat groups.

Figure 4.24 Percentage of Annex I habitat areas to be improved per habitat group

Note: Proportion of the total area reported by Member States that needs to be improved and proportion of unknown or not reported condition for each habitat group. The United Kingdom and habitat 8310 were excluded from the calculations.

Source: Article 17 Member States' reports and assessments.

The current area of habitats that **need to be improved** (i.e. areas reported as being in 'not good' condition) is estimated to be between 167 000 km² and 263 000 km², or 4-6 % of the EU territory. It should be noted that the quality of available data ⁽³¹⁾ on habitat condition on which this analysis is based is a limiting factor. Largely because of insufficient monitoring data, the condition of many habitats used in this calculation could only be determined for part of the habitat area (e.g. the condition of 63 % of coastal habitats and 50 % of dune habitats remains unknown). This means that the actual habitat areas in need of restoration are **likely to be much bigger** than those that have been estimated.

The average areas to be improved for each habitat type are as follows:

- Forests have the largest area in need of restoration targeted at improving their condition (approximately 100 000 km²); this equates to near 20 % of the total area of Habitats Directive forests needing to be improved.
- Coastal habitats need 16 % of their total area to be restored (approximately 46 000 km²).
- Grasslands need around 13.5 % of their total area to be restored (approximately 33 000 km²).

- Freshwater habitats need approximately 10.5 % of their total area to be restored (approximately 13 500 km²).
- Bogs, mires and fens need around 9 % of their total area to be restored (approximately 10 900 km²), including bog woodland, natural eutrophic and dystrophic lakes, oligotrophic waters, transition mires and quaking bogs.

The areas in need of restoration (improvement of their condition) listed above are shown in Figure 4.24 as a proportion of the total habitat area reported by Member States. In addition to maintaining areas that are already in good condition, restoration efforts also require increased investment from existing funding streams accompanied by appropriate policies to facilitate and prioritise the required restoration actions. This is confirmed by the EU Biodiversity Strategy for 2030 — Bringing nature back to our lives (EC, 2020d) — adopted by the European Commission, which puts restoring nature at its core.

As mentioned above, the incompleteness of the data reported confirms the need to further support, develop and implement appropriate monitoring schemes in Member States.

⁽³¹⁾ The calculations exclude the United Kingdom, given the relevance of these estimations for the post-2020 biodiversity strategy. Habitat 8310 (Caves not open to the public) was also excluded, taking into account its linear and underground characteristics.

Box 4.8 Closer look: Active restoration of habitats**Boost for alvar grasslands in Estonia**

Nordic alvar and Precambrian calcareous flatrocks (priority habitat type 6280*) are found around the Baltic Sea. The habitat is mainly under pressure because traditional low-intensity grazing has been abandoned. In Estonia, the project LIFE to Alvars (LIFE13 NAT/EE/000082) and State Forest Management Centre land management agreements have achieved substantial improvements in the country's alvar grasslands by implementing restoration measures. Key success factors were the fast, efficient large-scale mechanical restoration technique, improved communication between the local people and the state organisation (which facilitated restoration and grazing arrangements), availability of targeted agri-environment support, and the project team's efforts to enable local livestock owners to sign restoration agreements and agri-environment contracts. The habitat improvement was enabled by developing integrated coastal zone management and also by the local population's enhanced awareness of sustainable development and the benefits of nature conservation in this biosphere reserve.



Photo: Fixed dunes in Sweden © NEEMO

the surface area of fixed dunes Here, the project SandLife (LIFE11 NAT/SE/000849) addressed the problems of historical over-stabilisation of coastal and inland dunes and was instrumental in stopping the further decline of the fixed dune habitat. The project coordinated actions across 23 Natura 2000 sites in the south of the country and cleared 550 ha of scrub and trees, opened up 200 ha of dunes by disturbing the soil with tractor-mounted harrows and ploughs, and dug up 40 ha of the invasive Japanese Rose (*Rosa rugosa*).



Photo: Peatland restoration in Belgium © NEEMO

approximately to the total area of degraded raised bogs still capable of natural regeneration (7120) in continental Belgium. In regenerating bog habitats it can take up to several decades before the desired peat-forming vegetation is sustainably restored. However, it can be assumed that the large scope of the series of LIFE projects in the Ardennes plateau and the advanced stage of development of pioneer mire vegetation will soon lead to an improvement in the conservation status of the targeted mire habitats on a national scale.

Restoring fixed dunes in Sweden

Fixed — or grey — dunes (priority habitat type 2130*) comprise a secondary succession stage of dune formation. They are characterised by a perennial open vegetation of grasses, herbs, mosses and lichens, attracting specialised fauna, mostly invertebrates. Although they are present along most of Europe's coasts, three quarters are found in the Atlantic region. In all regions fixed dunes have a bad conservation status with mostly negative trends. The habitat is mainly threatened by encroachment of tall herbs and grasses, shrubs and trees, mostly as a consequence of planting trees for coastal defence or land abandonment. In the Continental region of Sweden there is a positive trend in

Peatland restoration in Belgium


The majority of the total habitat area of bogs and mires and grasslands has been lost over the last century as a result of human interventions, such as agricultural intensification, draining peatlands and land take/urbanisation. To overcome these pressures and restore peatlands, a series of six LIFE Nature mire restoration projects were successfully implemented in the Belgian Ardennes between 2003 and 2019, and more than 80 % of peatlands in Wallonia and about 40 % of all peatlands nationally were mapped. As a result, a total area of over 2 500 ha of peatlands show improved peat soil hydrology. The area on which restoration measures (mainly deforestation) have been completed corresponds

4.3.2 Restoration estimations per biogeographical and marine region

The biogeographical regions with the largest areas of habitats needing to be actively (re-)created are the Boreal, Mediterranean, Continental and Alpine (see Figure 4.25). The relatively small area needing to be (re-)created in the Atlantic region may indicate the physical and ecological impossibility of (re-)creating additional habitat areas because of current conditions, such as the high degree of urbanisation and intensive agricultural land use. However, the figures need to be read in relation to the total areas of Annex I habitat in the regions, where the total area in the Atlantic region (without the United Kingdom) is between 50 km² and 71 000 km², while for the Continental region it is between 250 km² and 280 000 km².

In terms of improving the condition of existing habitat areas, the biogeographical regions with the largest estimated needs are the Continental, Mediterranean, Atlantic, Marine Atlantic and Boreal regions. In the Continental region, the vast majority of habitat areas in 'not good' condition are beech and oak forests, lowland and hay meadow grasslands and freshwater habitats, in particular natural eutrophic lakes and lowland rivers. The main areas in the Mediterranean and Boreal regions to be improved are degraded forests, grasslands, scrub and heath (Mediterranean) and bogs and mires and freshwater habitats (Boreal). Degraded sandbanks, large shallow inlets and bays, and reefs require significant restoration in the Marine Atlantic region. More than half of the area to be restored in the Atlantic region is forests (mostly beech), as well as temperate heath, bogs and mires.

Figure 4.25 Estimation of Annex I habitat areas to be improved and area to be created per region



Note: 'Low' and 'high' range refer to the minimum and maximum areas in 'not good' condition as reported by Member States; 'additional' means the minimum areas to be re-created. The United Kingdom and habitat 8310 (caves) were excluded from the calculations.

Source: Article 17 Member States' reports and assessments.

Table 4.1 Estimation of carbon-rich Annex I habitat areas to be improved and area to be created per habitat group

Ecosystem	Low range (km ²)	High range (km ²)	Additional (km ²)
Coastal habitats	4 695	6 670	400
Heath and scrub	4 055	9 770	313
Grasslands	15 112	50 581	1 852
Bogs, mires and fens	9 331	11 567	1 690
Forests	84 455	109 978	4 648

Note: 'Low' and 'high' range refer to the minimum and maximum areas in 'not good' condition as reported by Member States; 'additional' means the minimum areas to be re-created. The United Kingdom and habitat 8310 (caves) were excluded from the calculations.

Source: Article 17 Member States' reports and assessments.

4.3.3 Restoration and climate change mitigation

Natural solutions are essential to confront the climate change and biodiversity crises. Here, many Annex I habitats that are particularly carbon rich offer the potential to store and sequester carbon and thereby contribute to mitigating climate change in addition to their biodiversity value. However, this capacity depends on their physical scale of coverage, condition and conservation status. Historically, land use decisions have been the primary driver of carbon stock changes across ecosystems, serving to decrease carbon stocks through land conversion (such as converting forest to cropland) or increase capacity through restorative measures (Goldstein et al., 2020).

Habitats considered 'carbon rich' were selected from the Habitats Directive to identify their contribution towards climate change mitigation targets. These include all Annex I forests, bogs, mires and fens, grasslands and temperate heath and scrub, as well as several coastal and halophytic habitats (mostly coastal marshes and mudflats ⁽³²⁾) and *Posidonia* beds (see Table 4.1). These habitats represent 148 out of 233 habitats in Annex I, covering an area of approximately 950 000 km² or around two thirds of the total habitat area reported across Member States. Here, carbon is stored in the above and below ground

biomass and (mainly) in the soil, with variations across sites and habitat types.

By maintaining and restoring such carbon-rich habitats or promoting active accretion of sediments in intertidal systems, climate change can be mitigated by directly reducing greenhouse gas emissions, safeguarding carbon stores and — in some cases — re-starting sequestration functions (Alonso et al., 2012). Policies and conservation measures must therefore prioritise improving the degraded ecological conditions of carbon-rich nature areas across Europe. Taking this into account, it is estimated that between 13 % and 19 % of the carbon-rich surface areas need to be improved (118 000-189 000 km²). In addition, it is estimated that a minimum of 9 000 km² of these habitats would need to be (re-)created to achieve good status in terms of distribution (range) and surface area.

In some cases, anticipated restoration measures may create trade-offs between increasing carbon sequestration and protecting biodiversity (e.g. the removal of trees to restore lowland heathland) (Alonso et al., 2012). All available information is thus necessary to underpin decision-making, including a comprehensive mapping of carbon- and nature-rich areas, the effects of management, habitat condition and factors such as soil and sediment characteristics (Alonso et al., 2012).

Key messages

- At least 11 000 km² of Annex I habitats need to be (re-)created, in addition to the area that currently exists, to ensure the long-term viability of each habitat type; this figure, however, is greatly underestimated because of the lack of reported data.
- The current area of habitats needing to be improved is estimated to be approximately 215 000 km² (or 5 %) of the EU-27 (EU-28 minus UK) territory; this is also underestimated given the incompleteness of the reported data.
- The biogeographical regions with the largest areas of habitats estimated to be actively needing to be (re-)created are the Boreal, Mediterranean, Continental and Alpine regions.
- Approximately 16 % of the carbon-rich surface areas of Annex I habitats need to be improved (154 000 km²). Restoring and maintaining carbon-rich habitats can make a significant contribution to climate change mitigation.



DASHBOARD
with detailed information by Member State:
condition of habitats

⁽³²⁾ Habitat codes 1120, 1130, 1140, 1310, 1320, 1330, 1410, 1420, 1430, 1510 and 1630.




Figure 5.1 Natura 2000 and conservation status

Natura 2000

Member States provide information* on the total number and area of sites classified under the Birds Directive (Special Protection Areas, SPAs) and proposed and designated under the Habitats Directive (Sites of Community Importance, SCIs, and Special Areas of Conservation, SACs) for both land and sea.


*Official statistics on Natura 2000 are provided on the dedicated EEA Natura 2000 Barometer.

What are the main facts on the EU Natura 2000 network?

The Natura 2000 network includes **27 852** sites
with an area of **1 358 125 km²**

Largest terrestrial area is Vindelfjällen in Sweden with **5 547 km²**; the **smallest** are the bat quarters in Walpersberg (Germany) with only **100 m²**

Largest marine area is the SPA in the French Mers Celtiques with **17 861 km²**; the **smallest** is Leixão da Gaivota (Portugal) with **1 600 m²**



What are the main results of the nature reporting?

Natura 2000 coverage

With a mean of 70 %, non-bird species have the highest **coverage**, especially **vascular plants and reptiles**.

Fish species are the least well covered (40 %).

For terrestrial habitats, **sclerophyllous scrubs and heaths** are best covered by Natura 2000 sites; **forests** on the other hand have the least coverage with 56 %.

Passage birds have the highest coverage among birds.

Effectiveness of Natura 2000 network

Measuring the effectiveness of Natura 2000 and its measures **remains limited** because of the lack of appropriate monitoring.

Habitats and species with a high coverage frequently show a significantly higher proportion of improvement and less deterioration than habitats with less coverage.

For **birds**, the correlation is less evident.

Land cover changes

Land cover changes are less within Natura 2000 than outside, but habitats are still being lost.

Dominant land cover changes within the Natura 2000 network occurred for **grasslands**.

Within the network, **arable land and permanent crops** have increased.

Full potential of Natura 2000 not yet achieved

Further action is needed to increase its effectiveness such as fostering coherence within and across Member States, improving the policy planning and implementation process and strengthening management and monitoring on the individual sites.



5 Status of the Natura 2000 network

Natura 2000 represents the largest coordinated network of nature conservation areas in the world, covering almost one fifth of the EU's terrestrial land area and approximately 10 % of Europe's seas. This network forms the backbone of EU nature conservation, contributing to maintaining or improving the conservation status of targeted habitats and species. While the terrestrial component of Natura 2000 is considered close to being complete, further marine sites are necessary.

The Natura 2000 network currently comprises almost 28 000 sites, classified either as 'special protection areas' under the Birds Directive or designated as 'Special Areas of Conservation' (and 'Sites of Community Importance') under the Habitats Directive. Special Protection Areas (SPAs) ensure that the 'most suitable territories' for bird species are designated. For sites to be designated under the Habitats Directive, Member States propose Sites of Community Importance (SCIs). The responsible Member States then have 6 years to designate the site as a special area of conservation (SAC), thereby formally enacting the requirement to apply necessary conservation measures.

Many sites are both (fully or partially) SPAs and SACs and are often also protected by other national or international designations (e.g. national parks, World Heritage or Ramsar sites). Sites are, however, not necessarily strict nature reserves and actually often include other uses, such as farmland or exploited forests. While much of the land included in the network is privately owned and managed, the use and management of all SACs and SPAs must comply with the ecological requirements of the species and habitats in question.

This chapter provides more detailed information on the current status⁽³³⁾ of the Natura 2000 network and gives an outlook on the actions needed to further strengthen the network. The current coverage of ecosystems, species and habitats and their trends and conservation status are outlined. Finally, we provide insights on the umbrella effect of the network and its effects on non-target species. The chapter is illustrated with case studies on the ecological coherence and drivers of the successful implementation and management of Natura 2000 sites.

The data for the present analysis includes Member States' reports as well as the Natura 2000 database and Corine Land Cover data.



Photo: © Nuno Alves, Environment & Me /EEA

⁽³³⁾ Using in particular data from the Natura 2000 Barometer (end of 2019).

5.1 Facts and figures

Member States provide information on the total number and area of sites designated under the Birds Directive (SPAs) and proposed and designated under the Habitats Directive (SCIs and SACs) as well as on the terrestrial area of sites and the proportion of the marine area of each site. The official data from the Natura 2000 network, however, are published by the EEA on the Natura 2000 Barometer ⁽³⁴⁾. For the general introductory figures on the current status of the Natura 2000 network, current Barometer data (end 2019) are presented.

As at end 2019, the Natura 2000 network includes 27 852 sites with an area of 1 358 125 km². The terrestrial component of the network comprises

nearly 784 994 km², representing 17.9 % of the EU's land territory. With this extent, the EU has achieved Aichi biodiversity target 11 ⁽³⁵⁾ through its Natura 2000 network alone since 2012. The remaining 573 131 km² include the marine area covered by Natura 2000 sites, amounting to 9.7 % of EU waters (Figure 5.2).

Maps 5.1 and 5.2 show the current number and area of terrestrial Natura 2000 sites per Member State in 2019. These presentations indicate the total area distribution and discloses country-specific geography by the over- or under-representation of their actual extents.

The highest terrestrial overall coverage of Natura 2000 sites is reported by Slovenia (38 %), Croatia (37 %), Bulgaria (36 %) and Slovakia (30 %). The lowest terrestrial coverage is reported by Denmark (8 %), the United



Figure 5.2 Natura 2000 area coverage





Note: Status end 2019.

Source: Natura 2000 database (2019).

Map 5.1 Number of Natura 2000 sites



Map 5.2 Total area of Natura 2000 sites



Note: These maps are cartograms that distort the geometry of regions to convey specific information by resizing them. The bottom right boxes show the Macaronesian islands (Azores, Madeira and Canary Islands). They include only terrestrial Natura 2000 sites for the EU-28 (SPAs, SACs, SCIs and proposed SCIs).

Source: Natura 2000 database (2019).

⁽³⁴⁾ <https://www.eea.europa.eu/data-and-maps/dashboards/natura-2000-barometer>

⁽³⁵⁾ Aichi biodiversity target 11 of the Convention on Biological Diversity's strategic plan 2011-2020 requires the conservation of at least 17 % of terrestrial and inland water and 10 % of coastal and marine areas.


Kingdom (9 %) and Latvia (12 %). Germany has the largest number of Natura 2000 sites (5 200) while Spain has the largest area (222 515 km²), followed by France (203 564 km²). With regard to the marine distribution of sites, France has the biggest Natura 2000 area with 132 689 km² (Map 5.3).

The Natura 2000 network has grown rapidly since the early 1990s, rising from 50 000 km² in 1993 (equivalent to the combined area of Luxembourg and Slovakia)

to 950 000 km² in 2010; the network reached over 1.35 million km² in 2019, which is equivalent to the combined area of France, Sweden and Romania (see Figure 5.3). The recent growth can mainly be attributed to the extension of the marine Natura 2000 network, which has doubled in area since the previous reporting period (2007-2012).

Member States need to ensure that sufficient protection and appropriate measures are implemented


Map 5.3 Natura 2000 network (terrestrial and marine areas)



Reference data: ©ESRI

Note: The map shows SPAs, SACs, SCIs and proposed SCIs (pSCIs).

Source: EEA (2020c).

Figure 5.3 Cumulative surface area of the Natura 2000 network from 1993 to 2019

Note: Many Natura 2000 sites are, partially or totally, both SPAs and SCIs; in calculating the area, the overlaps have been eliminated. Because of limitations in geographical information systems technology in the past, the area taking into account overlaps has only been available since 2011.

Source: Natura 2000 databases. Data are for the EU-28 and include SPAs, SACs, SCIs and proposed SCIs.

in Natura 2000 sites for habitats and species of Community interest and that they form a functional network. Member States adopt different strategies to achieve this goal: some select large areas (mostly in more natural and extensive landscapes, and in this case a Natura 2000 site also includes many non-targeted habitats and even human settlements) and some select small areas targeting only one habitat or the habitat of a species covered by the nature directives.

The range in size of Natura 2000 network sites is quite remarkable: some sites are less than 1 ha in size, e.g. the *Eiskeller in Klötze* (Germany) with just 100 m², while others cover vast areas spanning several thousands of square kilometres (see the lists of the smallest and largest sites). Some SPAs, for example, are classified to protect selected sea cliff habitats, which may be used by many thousands of breeding seabirds. Such sites are nevertheless likely to have a relatively low spatial area due to their vertical nature.

Within 6 years of their designation as SCIs, Member States need to designate these sites as SACs. In the reporting under the Habitats Directive, Member States give updates on the current state of SAC designations. According to this reporting, the number of SAC sites has more than doubled, with 7 262 new designations and around 303 800 km² added since the

end of the last period (2012). The designation status between Member States varies greatly. Whereas some countries report that their SAC designation process is complete, e.g. Austria, Belgium, Denmark, Estonia, Luxembourg and Slovenia, or nearly complete, other countries have only designated less than 20 % of their SAC areas, e.g. Bulgaria, Malta, Latvia, Poland, Portugal and Romania. Looking at the individual progress reported by Member States, some countries increased their SAC area significantly. Spain, for example, reported a more than 10-fold increase and France an increase close to 200 %, while the increase in designation in Portugal was only marginal within this reporting period (3.8 %).

With the designation of SACs, Member States also need to adopt conservation measures that involve, if need be, appropriate management plans and/or other measures that correspond to the ecological requirements of the natural habitat types and the species of Community interest. An analysis of Article 17 reporting shows a cumulative SAC area of 358 000 km² in the current reporting period⁽³⁶⁾ (131 500 km² reported in 2012), which equals over 15 500 Natura 2000 sites with management plans. Because of a change in methodology, a direct comparison between the two reporting periods is of limited value.

⁽³⁶⁾ Data limitations arise from missing Article 17 data from France and from the fact that the management plans are not mandatory measures and Member States are free to use other measures under Article 6.1 that are not included in this analysis.

Box 5.1 Europe's marine realm under protection**Photo:** North Sea © Congerdesign, Pixabay

The marine Natura 2000 network is making steady advances towards reaching the global biodiversity conservation goal of protecting 10 % of the world's oceans by 2020. Over 3 150 sites have been designated across the 23 coastal EU Member States, with exceptional progress being made in recent years, as the marine network area doubled since the last reporting period. The network encompasses over 570 000 km² of coastal and marine areas, which represents around 9.7 % of the total marine area of the EU, and protects 66 marine bird species, seven marine mammal species, two sea turtle species, 19 anadromous fish species ^(*) and nine marine habitat types (EC, 2019b). While Europe is advanced when it comes to designating marine protected areas, the vast majority of marine habitats and species targeted need to be better protected by the network to achieve good

conservation status. Developing management plans and implementing tailored management practices remain key challenges to effectively protect Europe's seas. Another challenge is the lack of data on species and habitats when it comes to marine protection; significant data gaps remain and thus assessment of conservation status and the effectiveness of measures proves difficult. There is still work to be done to complete the network, especially in terms of offshore waters and certain regional seas, as well as advanced monitoring practices and management measures to ensure that the protected area is not only growing in area and number of sites but also improving its actual capacity to deliver effective conservation of habitats and species. Seabird species are protected by sites classified as Special Protection Areas under the Birds Directive, but site designation is moving too slowly to safeguard species properly against significant disturbance.

Note: ^(*) Anadromous fish species hatch in freshwater, thereafter migrate to salt water for most of their life, and then returns to freshwater to spawn.

Sources: EC (2019b, 2020e).

Table 5.1 Largest and smallest Natura 2000 sites

TERRESTRIAL	MARINE
Five largest sites	Five largest sites
Vindelfjällen, Sweden, SPA/SAC, 5 547 km ²	Mers Celtiques — Talus du golfe de Gascogne, France, SPA, 71 861 km ²
Delta Dunării și Complexul Razim — Sinoie, SPA, Romania, 5 083 km ²	Mers Celtiques — Talus du golfe de Gascogne, France, SCI, 62 320 km ²
Delta Dunării, Romania, SCI, 4 536 km ²	Southern North Sea, United Kingdom, SCI, 36 951 km ²
Kaldoaivin Erämaa, Finland, SPA/SAC, 3 514 km ²	Banco Gorringe, Portugal, SCI, 22 928 km ²
Sologne, France, SAC, 3 462 km ²	Espacio marino del oriente y sur de Lanzarote-Fuerteventura, Spain, SCI, 14 328 km ²
Smallest site	Smallest site
Fledermausquartiere im Walpersberg bei Großbeutersdorf, Germany, SAC, 100 m ²	Leixão da Gaiyota, Portugal, SPA, 1 600 m ²



DASHBOARD
Natura 2000 VIEWER



DASHBOARD
with detailed information by Member State:
Natura 2000 barometer

5.2 Land use in Natura 2000 sites


The following analysis is based on a comparison of the area of the Natura 2000 network (Natura 2000 database 2018) with the Corine Land Cover (CLC) inventory from the Copernicus Land Monitoring Service. The terrestrial part of the Natura 2000 network is largely covered by forests and transitional woodland shrub as well as natural grasslands and heathlands (see Figure 5.4), making up over 60 % of the network. Moreover, about 41 % of all wetlands and water bodies (terrestrial and coastal combined) are included in the network.

The land cover composition of the Natura 2000 network area differs from that of the area outside the network. Coastal and marine wetlands and water bodies make up the smallest area in the EU, and they encompass many species and habitats that are particularly vulnerable to human activities such as drainage, which is part of agricultural activities, or hydropower

installations. Other major land cover classes with better coverage by the Natura 2000 network include natural grasslands and heathlands (41 %) and inland wetlands and water bodies (35 %).

Changes in land use between 2012 and 2018 differ inside and outside the Natura 2000 network, with less change overall within the network than outside (see Figure 5.4). The biggest changes **outside** the network relate to artificial surfaces, which increased by 2 571 km², and pastures and mosaic farmland, which decreased by 2 054 km²; this represents a huge loss of this land cover class that includes many natural and semi-natural habitats. The dominant changes **within** the network, which are of a much smaller order than changes outside the network, occurred in grassland ecosystems, which decreased by 291 km², and forests, which decreased by 214 km². The one exception is coastal and marine wetlands, which both grew by 8 km² between 2012 and 2018 inside and outside the network.

Figure 5.4 Land cover and land cover changes inside and outside the Natura 2000 network (in km²)



Inside Natura 2000 (km ²)	202	69	8	-214	-72	-291	-127
Outside Natura 2000 (km ²)	-128	2.571	8	-1.021	218	-684	-2.054

Sources: EEA (2020b); Corine Land Cover data for 2012 and 2018.

Within the network, arable land and permanent crops have increased, while grasslands and forests have decreased. Table 5.2 shows a matrix of land cover changes between 2012 and 2018. Pastures and mosaic farmland (with approximately 18 %) and inland wetlands and water bodies (with approximately 10 %) have been extensively transformed into arable land and permanent crops both inside and outside the network. Recent research has shown, however, that high nature value (HNV) farmland inside Natura 2000 sites is less likely to be converted into artificial surfaces than such farmland outside the network and is more likely to maintain its pattern of mosaic farming (Anderson and Mammides, 2020). The dominant change across land cover classes was conversion into arable land and permanent crops, while natural grasslands and heathlands (with approximately 7 %) have changed mainly into forest ecosystems.

Some of these changes can be linked to fires, which have significant effects on forest ecosystems and on natural grasslands and heathlands. The available data on wildfires show a high variability between years and regions in the EU but indicate an increasing risk, particularly in the Mediterranean countries. While a number of factors play a role in the frequency and level of impact of wildfires, climate change is critical and is projected to pose increasing risks in the years to come with potentially devastating effects on ecosystem functioning and biodiversity (de Rigo et al., 2017). In November 2017, the annual cumulative burnt area of Portugal, Spain and Italy alone exceeded 0.8 million hectares (de Rigo et al. 2017), the largest impact of forest fires in Europe. The FireLife project in Hungary demonstrates how fires can be prevented in some regions through targeted communication campaigns and by promoting biomass management among forest workers (see Box 5.2).

Table 5.2 Land cover changes/flows inside the Natura 2000 network (2012-2018)

Change matrix inside Natura 2000 (% of total changes)		2012						
		Arable land and permanent crops	Artificial surfaces	Coastal/marine wetlands and water bodies	Forests and transitional woodland shrub	Inland wetlands and water bodies	Natural grasslands and heathlands	Pastures and mosaic farmland
2018	Arable land and permanent crops		0.28	0.17	5.60	10.23	5.11	18.38
	Artificial surfaces	3.11		0.05	2.62	0.30	1.21	3.02
	Coastal/marine wetlands and water bodies	0.47	0.18		0.09	0.15	0.01	0.02
	Forests and transitional woodland shrub	3.89	0.72			0.37	7.09	5.07
	Inland wetlands and water bodies	3.05	1.22		1.59		0.95	2.54
	Natural grasslands and heathlands	0.88	0.25		4.08	0.06		0.54
	Pastures and mosaic farmland	7.84	0.73	0.20	4.27	0.77	2.88	

Sources: Corine Land Cover data for 2012 and 2018.

Box 5.2 Forest fires in Hungary — nature at risk

Photo: May 2012, after a 6-day forest fire in Bugac, Hungary © FIRELIFE

Over the last few decades, forest fires in Hungary have increased greatly. Fires have started because of climate extremes, a decrease in precipitation, an increase in mean annual temperature and a series of winters without snowfall. As a result, the period of fire risk has extended. Not only has the frequency of fires increased but also their intensity and the speed at which they spread. This was especially true during the extremely hot summers of recent years. These intense forest fires burn larger areas, as they are more difficult to extinguish. Fires are having an increasingly negative impact on vegetation, as well as on the structure of the forest.

The FIRELIFE project (LIFE13 INF/HU/000827) contributed to forest fire prevention in Hungary by carrying out a targeted communication campaign and by promoting biomass management among forest workers. Raising awareness of this climate change impact is crucial to increasing resilience. In 2012, the year before the launch of the project, 2 500 forest fires affected around 14 000 ha of forest in the country. Forest fires are mostly initiated by people, so the project was set up to show that effective communication of prevention measures can reduce the number and impact of fires. Highlighting appropriate regulation and targeted subsidies can also improve biomass and land use conditions to slow down and limit the spread of fires. As a result of the project's work, the number of forest fires had fallen to one third of the 2012 figure by the time it finished early in 2020.

Sources: EC (2019c); FIRELIFE (2020).

Key messages

- Forests and grasslands make up over 60 % of the terrestrial area of the Natura 2000 network.
- The biggest changes outside the network include an increase in artificial surfaces of about 2 600 km² and a decline in pastures and mosaic farmland of more than 2 000 km².
- While grasslands and forests have decreased within the network, these decreases have been smaller than the decrease in the same habitats outside the network.
- The dominant change across land cover classes within the Natura 2000 network was conversion of pastures and mosaic farmland into arable land and permanent crops.



DASHBOARD

Copernicus Land Monitoring Service Corine

Land Cover: land cover and change statistics

5.3 Coverage of species and habitats by Natura 2000


The habitats and species protected under the Birds and Habitats Directives are covered by the Natura 2000 network to varying degrees. Protected areas in general tend to be less effective in protecting wide-ranging and relatively common habitats and species and species that are highly mobile or have dispersed distributions. These are therefore covered to a smaller extent by the Natura 2000 network than rare and place-bound habitats and species. Variability in coverage also relates to the different approaches taken by Member States to designating and managing Natura 2000 areas and to targeting different species and habitats groups. When reporting, Member States were asked to indicate the habitat area or species populations covered by the Natura 2000 network. The completeness of these data varies among Member States and habitats/species groups.

To evaluate the importance of the Natura 2000 network in terms of coverage of habitats or species, Member States provide an estimation of the global habitat area or population size occurring within the network. In this section, the degree of coverage is illustrated by **violin plots**. The graphics in Figures 5.5 and 5.6 show a continuous distribution between 0 % and 100 %. The thickness of the tube corresponds to the number of assessments reported for the degree of coverage. The white point is the median indicating the 'middle' value in the list of numbers and the black line indicates the area containing 50 % of the data points (25th to 75th quantile).

Although coverage varies among different habitat groups (and single habitats), the overall mean coverage of habitats is relatively high (64 %). Habitats in general have slightly lower coverage by the Natura 2000 network than species (70 %). For sclerophyllous scrub habitats, the median and mean values are the highest of all the habitat groups. Forest Annex I habitats are the least covered at a median of 56 %. For forest habitats, there are many cases with medium coverage and only a smaller proportion with very high coverage.

It is worth noting that more freshwater habitats and grasslands have lower network coverage than is the case for the other habitat types. Freshwater and grassland habitat groups contain several relatively common and widespread habitats, which typically occur in intensively used landscapes with a lower density of protected areas. However, they are important for several species covered by both directives and should be more strongly promoted


Figure 5.5 Percentage coverage by the Natura 2000 network of Annex I terrestrial habitats



Note: The number of reports are indicated in parentheses. The total number of reports is 2 970.


Source: Article 17 Member States' reports and assessments.

Figure 5.6 Percentage coverage by the Natura 2000 network of Annex I marine habitats




Note: The number of reports are indicated in parentheses. The total number of reports is 2 970.

Source: Article 17 Member States' reports and assessments.

Figure 5.7 Percentage coverage by the Natura 2000 network of Annex II terrestrial species


Note: The number of reports are indicated in parentheses. The total number of reports is 3 214.

Source: Article 17 Member States' reports and assessments.

Figure 5.8 Percentage coverage by Natura 2000 of Annex II marine species

Note: The number of reports are indicated in parentheses. The total number of reports is 3 214.

Source: Article 17 Member States' reports and assessments.

Figure 5.9 Percentage coverage of birds by SPAs at the EU level in the period 2013-2018

Note: The number of reports are indicated in parentheses. The total number of reports is 2 825.

Source: Article 12 Member States' reports and assessments.

as key features to improve the ecological coherence of Natura 2000. Examples of these habitats include lowland hay meadows (6510), natural lakes (3150) and the comparatively less common and widespread plain to mountain rivers (3260, see Boxes 4.5 and 4.6). Sea caves (8330) receive mostly very high medium coverage; marine coastal habitats are slightly less well covered by the Natura 2000 network relative to terrestrial coastal habitats (see Figure 5.6).

Natura 2000 sites cover the core habitats of 1 035 Annex II taxa (see Figure 5.7). The best covered groups are vascular plants with a median of 90 %. A much lower coverage is reported for other species groups, especially fish with a median of only 49 %. Lower coverage for some species groups is perhaps due to their high mobility, with the network sometimes only including key areas of their habitat (e.g. for spawning) instead of all stretches of their migratory routes. Some species enjoy regionally full coverage under the Natura 2000 network, such as several vascular plants (especially endemic species occurring in few localities) or mammals such as Schreiber's Bent-winged Bat (*Miniopterus schreibersii*) in the Mediterranean region of France. This is also one of the species that underwent a genuine improvement within this reporting period (see Section 3.4.3). Marine species have very low coverage by the Natura 2000 network relative to most terrestrial species, with a median for marine mammals of 43 % coverage and for marine turtles of around 4 % (see Figure 5.8). This is understandable given the high mobility of these species and the fact that Natura 2000 sites mostly include breeding sites for these species. The specific challenge for these species lies in identifying and designating resting and foraging areas and their migration routes, which has not progressed far given the lack of studies and difficulty in outlining areas where they are significantly present.

The number of assessments for which Natura 2000 site coverage is unknown is much higher for species than for habitats. An unknown coverage is reported for over half of reptile, amphibian and mammal species, which denotes a lack of appropriate inventories and monitoring.

Member States designate SPAs for Annex I bird species, as well as for regularly occurring migratory species not listed in Annex I, known as 'SPA trigger species'. Coverage for birds is especially lacking for marine birds: on average, SPAs cover 16 % of the breeding distribution of seabird species and only 1.4 % of their non-breeding distribution (Ramírez et al., 2017). Coverage is best for passage birds with a median of 83 % (see Figure 5.9). However, the majority of the broad-front migrant species are not covered by the SPA network. This is likely to be explained by the location of important coastal and inland wetlands


and other roosting sites where many passage birds congregate within the Natura 2000 network. Breeding birds have a median coverage of only 48 %. This can possibly be attributed to the fact that their breeding distributions are much larger and nesting sites of several species are much more dispersed. Thus, it would be difficult to include the entire breeding area of a species within the Natura 2000 network.

Key messages

- Scrub, dunes, bogs and rocky habitats are better covered by the Natura 2000 network than forests or grasslands.
- Natura 2000 network coverage is generally higher for non-bird species than for habitats.
- Vascular plants are the best covered species group, while fish species are the least well covered.
- Terrestrial reptiles are well covered by the network on average, but the network falls short in protecting many reptile species.
- Coverage for birds is generally lower than for non-bird species, especially for breeding birds and marine birds, but there are many data gaps.




Figure 5.10 Percentage of Annex I habitats with $\geq 75\%$ and $< 75\%$ of their area covered by Natura 2000 with good conservation status



Note: The number of assessments per group is indicated in parentheses. The total number of assessments is 2 970.

Source: Article 17 Member States' reports and assessments.

Figure 5.11 Percentage of Annex II non-bird species with $\geq 75\%$ and $< 75\%$ of their area covered by Natura 2000 with good conservation status



Note: The number of assessments per group is indicated in parentheses. The total number of assessments is 3 527.

Source: Article 17 Member States' reports and assessments.

5.4 Effectiveness of the Natura 2000 network

The Natura 2000 network was established with the purpose of creating a coherent European ecological network of SACs. According to Article 3.1 of the Habitats Directive, the network 'shall enable the natural habitat types and the species' habitats concerned to be maintained or, where appropriate, restored at a favourable conservation status in their natural range'. The data currently reported do not allow a direct comparison of the conservation status of species and habitats inside and outside the Natura 2000 network to measure its effectiveness. Therefore, this section looks into other potential parameters, such as the link between level of coverage of species (Annex II of the Habitats Directive) and Annex I habitats and the proportion of favourable status assessments, and, more importantly, at the reported changes in status but also improvements in trends between the current (2013-2018) and last (2007-2012) reporting period. For this, the habitats and species assessments have been divided into three groups based on percentages covered by Natura 2000: $< 35\%$, $35-75\%$, $> 75\%$. These classes were chosen using a statistical technique (*k*-method) that is designed to divide data into homogeneous groups. The first analysis focuses on the comparison of high coverage for different groups, whereas the analysis of improvement looks at the differences between the different degrees of coverage. For the assessment of bird population trends, a comparison was made between SPA species and non-SPA species. The analyses are based on Member States' data and assessments.

Across Europe, **habitats** with a high coverage by the network have a significantly higher proportion of good assessments than habitats with a lower coverage. Higher Natura 2000 coverage is correlated with positive impacts on conservation status for most of the habitat groups, such as heath and scrub, forests, bogs, mires and fens, and coastal and dune habitats (Figure 5.10). Other habitat groups, such as freshwater, grasslands, sclerophyllous scrub and rocky habitats, show the opposite correlation: more of those with less than 75 % coverage are reported to have good conservation status than habitats with a higher level of coverage.

Non-bird species with high Natura 2000 coverage were assessed as having good conservation status significantly more often than those with lower coverage. Of the species with over 75 % coverage by Natura 2000 sites, 35 % were assessed as having good conservation status (Figure 5.11). In comparison, only 28 % of the species covered to a lesser extent by the


network were assessed as having a good conservation status. At the species group level, a clear pattern for good assessments emerges when seen in relation to the Natura 2000 coverage (i.e. over 75 % coverage). The data for reptiles, amphibians, fish, mammals and non-vascular plants support this conclusion, with greater spatial coverage significantly correlated with a higher proportion of good conservation status compared with groups with less Natura 2000 coverage. However, the data for arthropods, molluscs, other invertebrates and vascular plants do not support this correlation.

Interpretation of these results remains difficult because the status of these species before the establishment of the network is not known. Therefore, as was done for the measures in Section 4.2.3, the following analysis uses **improvements** within this period to further investigate the effectiveness of Natura 2000. These improvements were identified by comparing status and trends between the current and last reporting period (for more detail, see Sections 2.4 and 3.4). For Natura 2000, the focus is on Annex I habitats, Annex II species (Habitats Directive) and Annex I and SPA trigger species (Birds Directive).

The results show that **habitats** with more than 75 % coverage by Natura 2000 sites have a higher proportion of improvement (approximately 7 %) than those with less than 35 % coverage (5 %) (see Figure 5.12). At the same time, a higher coverage also resulted in less deterioration. However, the improvements remain rather small; most of the assessments did not change and predominantly have unfavourable status.

Bogs, mires and fens as well as dunes and forest habitats show the same results at the habitat group level, with higher coverage resulting in more improvement. With a reported improvement of over 8 %, the habitat group of bogs, mires and fens benefited particularly from higher levels of Natura 2000 coverage compared with low coverage (improvement of 1 %). At the biogeographical level, the Alpine region exhibits quite low degrees of improvement — despite high coverage — although the share of assessments reporting good status already is the highest among all regions (over 40 %). The habitats reported on under Article 17 that experienced the most significant improvements overall are widely protected by the Natura 2000 network, such as the dry Atlantic coastal heaths with *Erica vagans* (4040) and coastal dunes with *Juniperus* species (2250). Overall, the Natura 2000 network's effectiveness in improving habitats is highly variable and difficult to generalise, as additional indicators, such as the date of accession to the EU, would need be considered (EEA, 2020a).


Figure 5.12 Changes in conservation status and trends for Annex I habitats within different Natura 2000 coverage classes



Note: 'Improvement' corresponds to unfavourable assessments that improved or became good, 'Remained favourable' to assessments that kept their good status, 'No change' to unfavourable assessments that did not improve or deteriorate, 'Deterioration' to unfavourable assessments that further deteriorated or changed from favourable to unfavourable, and 'Unknown' to assessments with unknown status. The number of assessments per group is indicated in parentheses. The total number of assessments is 2 970.

Source: Article 17 Member States' reports and assessments.

Figure 5.13 Changes in conservation status and trends for Annex II non-bird species within different Natura 2000 coverage classes




Note: 'Improvement' corresponds to unfavourable assessments that improved or became good, 'Remained favourable' to assessments that kept their good status, 'No change' to unfavourable assessments that did not improve or deteriorate, 'Deterioration' to unfavourable assessments that further deteriorated or changed from favourable to unfavourable, and 'Unknown' to assessments with unknown status. The number of assessments per group is indicated in parentheses. The total number of assessments is 3 865.

Source: Article 17 Member States' reports and assessments.

For **non-bird species**, the results are not as clear as for habitats. Across European species, those with a Natura 2000 coverage of between 35 % and 75 % show the highest degree of improvement (approximately 9 %) compared with the other coverage classes (see Figure 5.13). However, the figure also shows that the share of already good assessments is lowest in this class, resulting in the smallest proportion of combined good status and improvement compared with the other groups. The species with more than 75 % coverage have the lowest proportion of deteriorating changes and also the most unknown assessments.

For **birds**, the non-SPA taxa are more frequently reported with ongoing decreasing trends or no change (Figure 5.14). However, the comparison is impaired by a significantly higher proportion of unknown trends within the SPA taxa.

Figure 5.14 Changes in population trends for Annex I and SPA trigger bird taxa compared with those for species not triggering the SPA classification



Note: The analysis shows the short term-population trends. The 'stable' trend category is split in two distinct categories: 'Stabilised', corresponding to populations with stable or fluctuating short-term but decreasing long-term trends, and 'No change', which includes populations with stable or fluctuating short-term trends but which are not decreasing in the long term. Category 'Increased' corresponds to increasing trends, 'Decreased' to decreasing trends and 'Unknown' to unknown population trends. The graph combines information for breeding, wintering and passage birds. The number of assessments per group is indicated in parentheses. The total number of assessments is 8 094.

Source: Article 12 Member States' reports and assessments.

One reason for the more inconclusive results for species could be that site protection is not always the best means of achieving a better conservation status for species with bigger home ranges or dispersed distribution or that generally have more complex requirements than habitats. Zehetmair et al. (2015) conducted their study on Natura 2000 beech forests as a protective habitat for forest-dwelling bats and found that the current management of Natura 2000 beech forests is almost identical to that of non-Natura 2000 commercial forests. Management, beyond the act of designating protected areas, is thus crucial to effective species protection under Natura 2000 (see Section 4.2). The effectiveness of Natura 2000 sites is not only subject to designation, as different factors can slow down or halt positive effects (EEA, 2020a). At a Natura 2000 site aiming to conserve cetaceans and turtles in Andalusia and Murcia, for example, a lack of political will and support was found to undermine the conservation efforts of the stakeholders involved in the site (Tucker et al., 2019).

Overall, it remains difficult to draw definitive conclusions about the effectiveness of the Natura 2000 network. This is especially because of the monitoring approach taken by several Member States, whereby the conservation status of habitats and species is mainly monitored and assessed in Natura 2000. Thus, there is a lack of data enabling comparison between the conservation status of and trends in species and habitats inside and outside the Natura 2000 network. While a holistic approach, using ecological data and data on pressures and threats, offers a more comprehensive evaluation of conservation status, an aggregation of data submitted

by Member States can mean that many positive developments at the local, regional or national scale are lost (Sotirov, 2017). A representative study of 9 602 Natura 2000 sites for birds indicated that the abundance of non-target species increased as the proportion of landscape covered by Natura 2000 increased (Pellissier et al., 2019). In addition, Koschova et al. (2018) looked at 252 European breeding bird species, of which 79 are listed under Annex I of the Birds Directive, and found significant positive effects in Member States on their trends, which was not the case in non-Member States. Yet the network's effectiveness for birds could be enhanced by taking account of richness patterns when designing SPAs for birds (Davis et al., 2020). Overall, there is not sufficient qualitative information on the design and implementation of the conservation objectives and measures within and outside the Natura 2000 network, which is crucial to achieving effective protection and monitoring.

Key messages

- The effectiveness of the Natura 2000 network is determined by the sites' coverage, management and ability to reduce pressures.
- Species and habitats are more likely to have a good conservation status if they are well covered by the Natura 2000 network.
- Habitats with a level of high coverage (> 75 %) by Natura 2000 sites show a significantly higher proportion of improvement and less deterioration than habitats with lower levels of coverage.
- Measuring the effectiveness of Natura 2000 network and its measures remains restricted by the limited implementation of comprehensive and appropriate monitoring inside and outside the network.
- Despite signs of the network's positive contribution, its potential has not been fully unlocked and the implementation gap is still significant.



5.5 Actions needed to strengthen the Natura 2000 network

Although the Natura 2000 network has been positively correlated with improvements in the status of the habitats and species it protects, there is significant room to move beyond surface area targets for protected areas and to concentrate on increasing their effectiveness. A comprehensive literature review by Davis et al. (2020) conducted in tandem with the *State of nature in the EU* report identified **four key overarching influencing factors** affecting the ecological effectiveness of the Natura 2000 network: (1) selection of Natura 2000 sites at Member State level; (2) management and monitoring of Natura 2000 sites; (3) the policy planning and implementation process; and (4) global and local challenges. These factors are briefly outlined below, drawing on the Member States' reports and EEA (2020a), and complemented with recommendations for future action.

5.5.1 Selection of Natura 2000 sites at Member State level

The selection of Natura 2000 sites can directly affect the effectiveness of the network as a whole. Inefficient site selection has been linked to politically motivated selection and giving low priority to conservation objectives compared with economic objectives. Unclear conservation objectives for the network also reduce its effectiveness, necessitating increased coherency between **conservation objectives and conservation measures**, not least by ensuring spatially coherent objective setting at the national scale accompanied by appropriate monitoring schemes with measurable indicators. Additional factors include a bias towards areas more removed from human activity, a delayed focus on marine protected areas and the inefficient protection of soil diversity. Incoherent planning and site selection approaches between and within Member States has led to insufficient functional connectivity and spatial connectedness between neighbouring countries and habitats and gaps in coherence within Member States. This highlights the need to increase connections between **protected areas and the level of protection** beyond the site. The current **'functional' area network** should be expanded as part of Natura 2000 and beyond to ensure coherence, connectivity, resilience, species/habitat benefits and the delivery of ecosystem services (including carbon storage as a

contribution to climate change mitigation). In the long term, this may include attaching stricter protections to Natura 2000 designations or more rigorously integrating environmental impact assessments.

5.5.2 Management and monitoring of Natura 2000 sites

Insufficient development and implementation of management plans for species and habitats, designed to increase the ecological effectiveness and cost-effectiveness of conservation measures, have had a negative impact on the achievement of conservation goals. Effective management and monitoring is further challenged by inadequate personnel, administrative and financial resources and by including diverse land categories with different ownership statuses, types of land use and human activity levels, and varying amounts of data availability. There is thus a significant need to increase the effectiveness of **management and monitoring practices**, in part via increased stakeholder participation. **Guidelines and models for effective public participation** should be developed and implemented, including encouraging volunteer-driven citizen science and outlining how to incorporate the data generated in existing monitoring systems⁽³⁷⁾. Guaranteeing that responsible authorities fully implement **Natura 2000 legal requirements** would ensure that important criteria for what is considered effective management, especially in terms of management planning, are met. Authorities should further **prioritise an ecological focus** in implementation and monitoring, and increase efforts to **harmonise monitoring** inside and outside the Natura 2000 network across Member States. Conservation and restoration objectives for Natura 2000 sites should be improved by defining **specific, measurable, comprehensive and realistic objectives** that can be monitored and evaluated and that also take into account the co-benefits of ecosystem services (e.g. carbon storage) delivered by certain habitats. Specific guidance from the European Commission would be useful to support Member States to clarify and implement their monitoring obligations, both in relation to species and habitats and in relation to Natura 2000 areas. Dedicated funding for monitoring (e.g. from the LIFE+ programme) might be necessary to achieve a step change in this respect. Incoherent management approaches between marine and terrestrial Natura 2000 sites and lack of coordination with other protected area networks were raised as another barrier to effectiveness.

⁽³⁷⁾ See, for example, the standards of the IUCN Green List of Protected and Conserved Areas: <https://www.iucn.org/regions/europe/our-work/natura-2000-europes-protected-areas-network/iucn-green-list>

Box 5.3 Closer look: Good practice examples of monitoring, knowledge and policy**Improved governance and capacity building**

Examples of enhanced capacity building can be found in all Member States since the beginning of the LIFE programme. Large carnivore conservation in Romania is among the beneficiaries of capacity building, underpinned by three LIFE projects. These projects supported the establishment of an independent network of experts and managers, who were able to advise the Romanian Ministry of Environment on large carnivore issues (Vrancea, LIFE02 NAT/RO/008576, Carnivores Vrancea II, LIFE05 NAT/RO/000170, and URSUSLIFE, LIFE08 NAT/RO/000500). The first project started with a team of three people, which grew to over 25 people working on large carnivore conservation issues in 2013 within URSUSLIFE or other parallel projects. The expert network created and reinforced over the years has proven itself especially useful for implementing monitoring activities. Since the initial three projects were concluded, two other LIFE projects have taken over: LIFE FOR BEAR (LIFE13 NAT/RO/001154) and WOLFLIFE (LIFE13 NAT/RO/000205). Based on the initial LIFE projects, the operators have the necessary expertise and capacity to lead large nature conservation projects. For instance, WOLFLIFE is coordinated by the Vrancea County Environmental Protection Inspectorate, which led the LIFE02 project with the Asociația Pentru Conservarea Diversității Biologice (APCDB) as an associated beneficiary. The APCDB was a local non-governmental organisation at the time of the first project in 2002. In 2013, it was still managed by the same core team of experts, who had managed six other projects on large carnivore conservation, funded by different sources (from National Geographic to EEA grants — more than EUR 2 million investment in total), drawing on the initial capacity-building impact of the LIFE programme. In Romania, the wolf has a favourable conservation status, which has remained stable over the years.



Photo: Citizen science © Dimitris Poursanidis, Environment & Me / EEA

Volunteer-driven citizen science and knowledge exchange

During the project LIFE Euro Bird Portal (LIFE15 PRE/ES/000002), the partners of the European Bird Census Council developed a web portal showing the distribution of 105 bird species across Europe. The Euro Bird Portal (EBP) is able to produce weekly animated maps that display data from January 2010 up to the current week at a resolution of 30 km × 30 km ^(a). The bird observation data are collected on a daily basis from 28 European countries and submitted automatically to a central repository that was created during the LIFE project. The main outcome is the production of daily maps and graphs showing near to real-time information. Altogether, over 40 million map combinations are possible. The EBP viewer and the central database now cover all EU countries (except Malta) as well as Turkey, Norway and Switzerland and have been updated with more than 320 million new inputs since the LIFE project started. This enormous effort is possible because of the participation of more than 120 000 bird watchers across the EU and highlights an efficient and far-reaching collaboration between 82 European entities. It is the largest citizen science initiative at EU level and is the only big data project in Europe dealing with biodiversity data. It shows how the work of many entities, scientists and bird watchers can be gathered and combined to deliver relevant outcomes at the European level. The final project outputs are highly relevant for research, nature conservation and education as well as policy enforcement and development. The EBP offers enormous possibilities for the future, although it will largely depend on external funding. Future collaborations with relevant organisation and institutions at EU level will be crucial to make the best use of the data produced.

Note: ^(a) <https://eurobirdportal.org/ebp/en>

5.5.3 Policy planning and implementation process

The ability of policymakers and planners to inform and evaluate effective planning is often jeopardised by a lack of knowledge of the ecological requirements and pressures affecting habitats and species, the lack of standardised monitoring across Natura 2000 sites, the absence of smart objectives and measures, and poor financial planning and lack of resources. A lack of reliable data or insufficient communication of existing data to policymakers and planners create bottlenecks in the use of information from gap analyses and monitoring of ecological processes, which should be used as the basis for decision-making. The resulting low awareness of the diverse benefits produced by the Natura 2000 network is often compounded by a long-standing conflict between economic or political interests and conservation goals. There is thus an urgent need to increase coherence between **biodiversity policy** and other policy areas, such as in the fields of agriculture and economic and rural development, and create a more integrated approach to address potential conflicts and trade-offs between various interests while fostering synergies. Concrete recommendations include strengthening requirements for **environmental impact assessment** for EU policies and increasing the focus of responsible authorities on **potential synergies**, such as green infrastructure and ecosystem-based disaster protection, climate mitigation and adaptation. Greater **political support** to enforce commitments made under other environmental or land use policies (e.g. Water Framework Directive and the greening measures of the common agricultural policy) would also make a considerable difference in reducing high-impact pressures and threats such as nitrogen pollution and hydrological change. The targets set in the new EU Biodiversity Strategy for 2030 respond to most of these challenges but require much stronger implementation than those in the EU Biodiversity Strategy to 2020 and significant efforts to take a cross-sectoral approach (see also Chapter 7).

5.5.4 Global and local challenges

Research on the effect of climate change on the Natura 2000 network identify it as a serious challenge in terms of both habitat and species protection. Although the impact on network connectivity is predicted to be rather small, it is anticipated that species and habitat shifts in range and population declines due to climate impacts will reduce the network's effectiveness. A particular concern is that protected sites may become unsuitable for targeted species. Adopting a long-term perspective, it is thus critical to increase the **adaptive capacity** of and **functional connectivity** between protected sites. This entails creating functional connections enabling species to migrate over long distances in pursuit of habitats that have shifted as a consequence of climate change. Management decisions should consider an **expansion of the network** to increase coverage and include more sites to help species and habitats adapt to climate change, increase the network's contribution to **carbon storage** and provide a range of other ecosystem services and benefits. Finally, more consideration should be given to **non-conservation activities** within the network and activities outside protected areas. Furthermore, although site-level management is sufficient in many countries, conservation objectives are sometimes not being met due to pressures from outside the sites (e.g. nitrogen pollution or changes in hydrological regimes). The effective implementation of other directives is therefore critical, as well as an increased policy focus on establishing effective **buffer zones and semi-natural habitats** outside protected areas and ensuring adequate levels of protection inside sites. Moreover, national and local measures need to be directed not only at habitat management but also at reducing cross-cutting pressures and threats such as air and water pollution. This requires investment in integrated approaches⁽³⁸⁾ and sharing learning and experiences between Member States.

⁽³⁸⁾ Examples of such approaches include the catchment sensitive farming air quality strategy in the United Kingdom and the integrated approach to nitrogen (PAS) in the Netherlands.

5.5.5 Addressing knowledge and communication gaps

Across all of these key areas, there is the cross-cutting recommendation to address **knowledge and communication gaps** in science, policy, and practice. This entails putting more emphasis on the effective dissemination of the data reported by Member States to encourage their broader uptake and application. Research can help to fill gaps in the data reported and provide valuable insights and detailed case studies on selected species and habitats. On this basis, **sound scientific knowledge** should play a greater role in decisions about planning, site selection, management and policy, and insights from implementation should be more fundamentally integrated into science. Furthermore, there is a need to facilitate **targeted knowledge sharing** on the effectiveness of management practices and the exchange of best practice at site level.


Key messages

The key actions needed to strengthen the Natura 2000 network entail:

- Increasing the adaptive capacity of and functional connectivity between protected sites and the level of protection beyond the site;
- Increasing the effectiveness of management and monitoring practices;
- Addressing knowledge and communication gaps in science, policy and practice; and
- Streamlining biodiversity protection across key sectoral policies.



Figure 6.1 Summary of progress towards EU 2020 biodiversity strategy targets



6 Meeting EU 2020 Biodiversity Strategy targets

In an effort to halt the loss of biodiversity and the degradation of ecosystems in the EU, the European Commission adopted a biodiversity strategy in 2011 that covered the period up to 2020. The strategy includes six targets to be reached by 2020, two of which explicitly mention species and habitat conservation status. Target 1 strives to achieve a proper implementation of the nature directives, not least through improvements in the status of all species and habitats covered. Target 3 aims to optimise agriculture and forestry's benefits for biodiversity.


This chapter outlines the progress made towards targets 1 and 3 of the EU Biodiversity Strategy to 2020. Target 1 focuses on the amount of secure/favourable or improving assessments for birds (Birds Directive) and habitats and species of Community interest (Habitats Directive). Target 3 provides more detailed information for agriculture and forest ecosystems, outlining the current status, contributing factors and necessary actions to reverse undesirable trends. These analyses stem from the biogeographical assessments of habitat types and species listed in the Habitats Directive and the population status assessments of bird species from the Birds Directive, using methodologies approved by the Member States and the European Commission ⁽³⁹⁾.

6.1 Measuring progress towards target 1

Target 1 of the EU Biodiversity Strategy to 2020 aims to achieve a full implementation of the Habitats and Birds Directives for European habitats and species, maintaining them at and restoring them to favourable conservation status by 2020. Specifically, it aims to reach:

- 100 % more habitat assessments showing a favourable or improved ⁽⁴⁰⁾ conservation status;
- 50 % more species assessments under the Habitats Directive showing a favourable or improved conservation status; and

Figure 6.2 Progress towards target 1 (percentage of assessments)



Note: Each bar represents the percentage of assessments showing a favourable/secure or improving status.


Source: Article 12 EU assessments and Article 17 Member States' reports and EU assessments.

- 50 % more species assessments under the Birds Directive showing a secure or improved population status.

The overall progress made across species and habitats towards target 1 is relatively small compared with the 2010 baselines, i.e. **the target was not met for any of the groups** (see Figure 6.2). More specifically, the 2020 target of reaching favourable and improving assessments for 34 % of habitat assessments was not met (12 % short of the target). In comparison, non-bird species assessed as favourable or improving have increased almost in line with their 2020 target, falling only 2 % short of the 35 % target when considered collectively. The picture for birds is more diverse, with an increase in the number of secure and improving taxa between 2010 and 2015, followed by a decrease between 2015 and 2020 (-3 %). Overall, a gap of over 20 % remains to achieve target 1. Although the decline could be due in part to better and more comprehensive data and to a slight change in the assessment methodology, e.g. more sensitive thresholds for stable,

⁽³⁹⁾ Doc NADEG 18-05-06 (Measuring progress under Target 1 of the EU 2020 biodiversity strategy): <https://circabc.europa.eu/w/browse/72cd273c-a270-4b8c-8604-fcb57549839c>.

⁽⁴⁰⁾ The term 'improved' includes improvement in both status and trend.

Figure 6.3 Trends in unfavourable assessments not contributing to target 1 (%)

Note: For habitats and non-bird species, 'Deteriorating' includes unfavourable assessments (poor and bad status) that show a negative trend between 2010 and 2020 and 'No change' indicates unfavourable assessments that have not improved or deteriorated; for bird species, 'Deteriorating' corresponds to long-term trends where short-term trends are not stable or increasing and 'No change' includes stable/fluctuating trends.

Source: Article 12 EU assessments and Article 17 Member States' reports and EU assessments.

decreasing and increasing trends, there is evidence that many species have undergone genuine deterioration of their population trends in both the short and long terms. For non-bird species, it should be noted that there was a decrease in the number of unknown assessments, which may have contributed to more favourable and improved assessments.

Over 41 % of habitat assessments remained stable (i.e. no improvement or deterioration in their unfavourable status) and 32 % deteriorated between 2010 and 2020. For species, 31 % of the assessments did not change and 26 % deteriorated. Still, over 73 % of the habitat and 57 % of the species assessments remain unfavourable (see Figure 6.3). For birds, 23 % of the assessments indicate a deterioration (declining short-term trend) and 1 % have a stable or fluctuating short-term trend and a long-term trend that is not declining.

Target 1 has not been achieved in large part because of a failure to sufficiently address the main drivers of biodiversity loss, including intensive agricultural practices, land abandonment, urban sprawl, grey infrastructure development, pollution and human activities at sea (see details in Chapter 4 on pressures and threats). Stronger mainstreaming of biodiversity — making it part of everyday practice — with other sectors and coordination of policies is needed, including agriculture, marine, forestry, energy and water legislation (EFH, 2019). The nature directives' effectiveness has also been impeded by delays in establishing the Natura 2000 network and in implementing necessary management actions (e.g. conservation measures, species protection, restoration, monitoring, financing) (EC, 2016; EFH, 2019). According to the fitness check (EC, 2016), further challenges include limited and poorly targeted funding (including the continuation of perverse subsidies), knowledge gaps, insufficient stakeholder engagement and human resource constraints.

Key messages

- Target 1 of the EU Biodiversity Strategy to 2020 has not been reached.
- A 12 % gap remains to achieve the target for habitats, a 20 % gap remains for bird species and only a 2 % gap remains for non-bird species.
- High shares of assessments are further deteriorating: 32 % for habitats, 31 % for non-bird species and 23 % for bird species.

6.2 Measuring progress towards target 3

Target 3 of the EU Biodiversity Strategy focuses on two of the most significant economic activities having an impact on biodiversity, i.e. agriculture (target 3a) and forestry (target 3b). Combined, these sectors comprise more than two thirds of the entire EU terrestrial area. For more information on the specific pressures, see Chapter 4: Pressures and responses.

6.2.1 Agriculture

Target 3a of the EU Biodiversity Strategy to 2020 focuses on agriculture, which is one of the main sectors generating pressures on European habitats. The target aims to 'maximise areas under agriculture across grasslands, arable land and permanent crops that are covered by biodiversity-related measures under the Common Agricultural Policy', targeting agricultural habitats that make up 28 % of the 1.3 million km² terrestrial area reported for Annex I habitats. The assessments at the EU level (based on Member States' reporting on 233 habitats) highlight that:

- The conservation status of Annex I agricultural habitats is still assessed as bad (45 %) and poor (38 %); other Annex I habitats, for comparison, are assessed as 31 % bad and 45 % poor.
- Only 8 % of agricultural habitats are assessed as improving, whereas 45 % are assessed as deteriorating.

The negative effects of the pressures linked to agriculture become even more evident when comparing habitats that are partially (over 212 300 km²) or fully (over 169 200 km²) dependent on agricultural activities.

67 % of agricultural habitats that are fully dependent on adequate agricultural management and 37 % of those that are partially dependent were assessed as 'bad' (see Figure 6.4). Similarly, deteriorating trends are found far more frequently in habitats that are fully dependent on agricultural management measures than in those that are only partially dependent. Improving trends are infrequent, reported in only 6 % of fully dependent and 9 % of partially dependent agricultural habitats. This compares with 32 % of other non-agricultural habitats deteriorating and over 9 % improving (see Figure 6.5). Finally, the condition of agricultural habitats that are fully dependent on

Figure 6.4 Conservation status of agricultural habitats (partially and fully dependent on agricultural management)





Figure 6.5 Trend in conservation status of agricultural habitats




Source: Article 17 Member States' reports and EU assessments.

management had fewer 'good' and more 'not good' cases than those that are partially dependent on management. Overall, the percentage of good cases for the agricultural habitats was lower than that of other habitats (12 % versus 16 %); the not good cases were similar for both agricultural and other habitats.


Grasslands make up 16 % of the total area of Annex I habitats, with over 265 700 km² and 52 Annex I habitats. A large share of grasslands are also categorised as agricultural habitats. Almost half of the grasslands assessed have a bad conservation status and over a third were assessed as poor. Only 7 % of grasslands assessed showed an improving trend, while nearly 51 % of grassland trends were classified as deteriorating.

Figure 6.6 Conservation status of grasslands by type




Source: Article 17 Member States' reports and assessments.

Figure 6.7 Trend in conservation status of grasslands by type



Source: Article 17 Member States' reports and assessments.

Figure 6.8 Trends in status of birds and non-bird species in agricultural habitats and grasslands



Source: Article 12 and Article 17 EU-level assessments.

Grasslands that are fully dependent on the implementation of management measures (over 116 600 km²) are more frequently reported as having a bad conservation status (68 %) than grasslands that are partially dependent on their management (over 166 400 km²) and natural grasslands (over 166 100 km²), with 44 % and 45 %, respectively. The differences between the conservation status assessments of natural grasslands and those grasslands that are partially dependent on agricultural activities are minimal (see Figure 6.6). Similar results were observed for trends in conservation status. Grasslands that are fully dependent on management are less frequently reported as improving (4 %) than natural grasslands and grasslands that are partially dependent on management (both with 8 % improving). However, the percentage of unknown assessments was far higher for natural grasslands and grasslands that are partially dependent on management (see Figure 6.7). For grassland species, the results are mixed (see Figure 6.8). Approximately 18 % of agricultural bird species are improving, but almost 54 % are deteriorating. In contrast, only 5 % of non-bird grassland species show improvements, but also have a lower share deteriorating (29 %).

Grasslands are also key habitats for many pollinators. As shown above, neither their status nor their trends are improving. The importance of grassland and other habitats for pollinators is detailed in Box 6.2.

Box 6.1 Habitats depending on agricultural management for their viability

- Agricultural habitats fully dependent on agricultural management refers to semi-natural habitats established under regular (usually low-intensity) agricultural management. The species composition is a product of the site conditions and the type and intensity of management. Stopping management practices or changing the management intensity will result in (usually irreversible) changes in the habitat structure and species composition and trigger a change to another habitat type.
- Agricultural habitats partially dependent on agricultural management refer to habitats that profit from agricultural management measures because they either maintain the habitat type or enlarge/maintain the habitat distribution, often by preventing or reducing secondary succession to another habitat type.

Source: Adapted from Halada et al. (2011).

Box 6.2 Closer look: Grasslands and pollinators

The European Commission recognises the importance of certain habitats for pollinators within the EU pollinator initiative (EC, 2018). The initiative aims to address the decline of pollinators in the EU and contribute to global conservation efforts and sets out measures under three priorities:

1. improving knowledge of the decline in pollinators, its causes and consequences;
2. tackling the causes of the decline; and
3. raising awareness, engaging society-at-large and promoting collaboration.

Recent work by the EEA and European Topic Centre on Biological Diversity consortium identified Annex I habitats that are particularly important for pollinators (Kudrnovsky et al., 2020). As a first step, plant species classified as important for pollinators were assigned to their respective habitats. Based on the number of animal-pollinated plant species, the importance of each Annex I habitat was derived. With a mean of 111 plant species classified as being important for pollinators, grasslands are the most important habitat group for pollinators. The second most important habitat group is sclerophyllous scrub with a mean of 77 species. The three most relevant habitats following these groups are heath and scrub, bogs, mires and fens and forests. Single habitats with more than 400 plant species assigned to them include semi-natural dry grasslands (see also the European Commission's habitats action plan for dry calcareous grasslands (6210); EC, 2019d) or alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (91E0).

Furthermore, the analysis focuses on the five most important habitat types for pollinators in each group. When comparing the top five habitat types with the whole group, it can be seen that these habitats mostly have a poorer conservation status than the average of the whole group (Figure 6.9; see Figure 4.2 for the five most relevant habitat groups). This is especially true for the two most important habitat groups for pollinators, grasslands and sclerophyllous scrub, but also applies to bogs, mires and fens as well as forests. The same applies for the conservation status trend, which is worse for the top five habitat types within grasslands, sclerophyllous scrub and forests (Figure 6.10). For the purpose of comparison, the overall results for conservation status and trends per group can be found in Sections 3.2.1 and 3.3.1.

Figure 6.9 Proportion of conservation status in the five most relevant habitat groups





Figure 6.10 Proportion of conservation status trends in the five most relevant habitat groups



Source: Article 17 Member States' reports and assessments.

The main reasons for the poor status of grasslands important for pollinators are grassland abandonment, natural succession, intensive grazing and the application of fertilisers. Furthermore, alluvial forests that are rich in species important for pollinators (e.g. based on their many early spring species in the herb layer which are pollinated) often have bad conservation status. There are various reasons for this; the main impact for pollinators is, however, changes in hydrology and the connected conversion of those forests into other habitats.

Species linked to agricultural habitats show similar results to grassland species regarding their assessments of conservation status and trends (see Figure 6.8). Almost one third of species were assessed as having good conservation status, while around 15 % had bad status (e.g. the Tartarian Breadplant and *Tripolium sorrentinoi*, many butterflies and amphibians) and 44 % had poor conservation status. Only around 5 % of the species linked to agricultural or grassland habitats showed improving trends (see Figure 6.9), such as a few bat species (*Miniopterus schreibersii*, *Myotis* spp.), plants such as the Marsh Angelica, the endangered *Carduus myriacanthus*, the vulnerable *Carlina onopordifolia* and the Eurasian Toothed Grasshopper. Trends for non-bird species showed that around one quarter of species are deteriorating and one third are stable. Trends in conservation status are unknown for almost one third of non-bird species, hindering firm conclusions being drawn. Status assessments of bird species show an exceptionally high rate of not secure populations at 75 % for species present on farmland (e.g. larks, shrikes and buntings). Short-term population trends for farmland birds reveal that 54 % are deteriorating, 21 % are stable and 18 % are improving; these trends echo those of the common farmland bird index and the grassland butterflies index (see Box 6.3).

Compared to the mid-term results in 2015, assessments of agricultural habitats show an overall deterioration in conservation status: good status decreased from 14 % to 12 % and bad status increased from 39 % to 45 %. Assessments with improving conservation status trends tripled from over 3 % to 10 %, while simultaneously those with deteriorating trends display a 34 % to 39 % increase. Agricultural species (other than birds) show similar, but slightly better results: good status assessments are more frequent than in 2015 (from 27 % to 30 %), but poor and bad assessments also increased, particularly the later ones from 14 % to 16 %. Improving trends increased slightly (from 4 % to over 5 %), as did the deteriorating trends from 25 % to over 29 %.

As shown by the results presented, farmland biodiversity continues to decline. While targeted

agri-environmental and climate measures funded through the common agricultural policy can have positive impacts on sustainable farming systems and the conservation of semi-natural farmland habitats and species, their impacts are limited at European scale because of insufficient application and coverage (14.6 % of utilised agricultural area in 2017) (Pe'er et al., 2017; EC, 2019e). This favouring of low-impact measures rather than greening and cross-compliance options by Member States inhibits the potential to deliver biodiversity benefits (ECA, 2020). Another weakness is the inadequate designation of environmentally sensitive permanent grasslands, leaving the majority of grasslands vulnerable to ploughing and further jeopardising biodiversity (Pe'er et al., 2017; EC, 2019e). Moreover, there are no reliable indicators for measuring the results and impacts of direct payment schemes and rural development programmes in relation to biodiversity (ECA, 2020). An in-depth analysis on ongoing agricultural pressures is provided in Section 4.1.1.

6.2.2 Forests

Target 3b of the EU Biodiversity Strategy 2020 aims to increase the contribution of forestry to maintaining and enhancing forest biodiversity, identifying a number of actions to achieve measurable improvements. Forest habitats make up almost one third of the area of Annex I habitats with over 500 000 km², approximately 60 % of which are temperate forests and 25 % are boreal and Mediterranean (overlap between the groups is possible). Assessments revealed that:


- Close to one third of the European forest assessments showed bad conservation status (31 %).
- Over half of the assessments showed poor conservation status (54 %).
- Approximately 14 % of the assessments showed good conservation status.
- On average all other Annex I habitats were assessed as 38 % bad, 41 % poor and 15 % good.

Box 6.3 Abundance and distribution of common bird and butterfly species


Birds and butterflies are sensitive to environmental change and their population sizes can reflect changes in ecosystems as well as in other animal and plant populations. Trends in bird and butterfly populations can thus be excellent barometers of the health of the environment. The status of birds and butterflies has been the subject of long-term monitoring in Europe, much of it via voluntary efforts. Both species groups have a strong resonance with the interested public and are good examples of how the power of citizen science can be harnessed by effective targeting.

The long-term trends (over 25 years) revealed by monitoring schemes for common birds — in particular farmland birds — show significant declines, with no signs of recovery (Figure 6.11). Between 1990 and 2016, there was a decrease of 9 % in the index of common birds across the 26 EU Member States with bird population monitoring schemes. The common forest bird index decreased by 3 % in the EU. The decline in common farmland bird numbers was much more pronounced, 32 % at the EU level.

The index of grassland butterflies has declined significantly in the 15 EU countries where butterfly monitoring schemes exist. In 2017, the index was 39 % below its 1990 values (Figure 6.12).


Figure 6.11 EU common birds population index from 1990 to 2016

Source: EEA (2020b).

Figure 6.12 Population index of grassland butterflies 1990-2017


Source: EEA (2020b).

Figure 6.13 Conservation status of forests by region




Source: Article 17 Member States' reports and EU assessments.

Figure 6.14 Trend in conservation status of forests by region



Source: Article 17 Member States' reports and EU assessments.

Figure 6.15 Trends in status of forest non-bird and bird species



Source: Article 12 and Article 17 Member States' reports and EU assessments.

While these numbers do not show much progress, the trends in conservation status are more favourable with over 54 % of all forest habitats assessed showing stable or improving trends. However, about one third of the forest assessments are deteriorating (27 %) and 19 % are unknown. In comparison, 41 % of Annex I habitats other than forests were assessed as deteriorating and 37 % as either stable or improving.

Boreal forests had the highest amount of bad assessments (56 %) (see Figure 6.13) and the highest percentage of habitats with deteriorating trends in conservation status (43 %) (see Figure 6.14). Other forest groups had noticeably better results.

Close to one third of forest bird species were assessed as having an improving trend (34 %), while 37 % have a stable trend and 17 % a deteriorating trend (see Figure 6.15). For forest non-bird species, almost one third of assessments have an unknown conservation trend (31 %), 27 % have a deteriorating trend and only 6 % are improving. Bird species in forest habitats were assessed as having a substantial percentage of not secure populations (79 %), with only 5 % assessed as secure. This indicates an even higher negative population status than farmland bird species. In terms of trends, however, forest bird species show more promise than farmland species. Over one third of forest species are increasing in the short term (34 %) and 37 % are stable, while only 17 % were assessed as decreasing (compared with 54 % of farmland birds assessed as decreasing; see Box 6.3).

Compared to the mid-term results in 2015, assessments of forest habitats show a deterioration in conservation status: good status decreased from 16 % to 14 % and assessments revealing a bad status increased from 27 % to 31 %. Meanwhile, assessments with improving conservation status trends increased by four, from over 3 % to 13 %, and those with deteriorating trends decreased from 29 % to 24 %. The percentage of assessments of forest species (other than birds) in good status is higher than in 2015 (from 27 % to 30 %), but poor and bad assessments also increased in total from 56 % to 59 %. Improving trends also increased slightly (from 4 % to 6 %), as did the deteriorating trends from 21 % to over 27 %.

In conclusion, the conservation status of forest habitats and species covered by EU nature legislation shows no significant signs of improvement. This is in part because of the large variance in thresholds across Member States regarding the eligibility of forest management plans or equivalent instruments for funding (ranging from requiring plans for all forests to only requiring plans for forests larger than 100 ha) (Langhout, 2019). Yet, further forestry management plans are being developed and could serve as a step forward in ongoing conservation efforts. An in-depth analysis on ongoing forestry pressures is provided in Section 4.1.3.

Key messages

- No real progress was made towards reaching target 3 for agriculture (target 3a) or for forestry (target 3b).
- More than 46 % of Annex I assessments of agricultural habitats and 31 % of assessments of forest habitats show a bad conservation status.
- The status and trends for habitats that are fully dependent on agricultural management are worse than for habitats that are partly dependent and, for example, natural grasslands.
- Over 90 % of boreal forest habitat assessments show an unfavourable conservation status and worse trends than temperate and Mediterranean forests.
- Agricultural/farmland species have worse trends than forest species, with farmland bird taxa and forest bird taxa showing high rates of deteriorating trends (54 % and 17 %, respectively).
- The poor status of and trends in agricultural habitats and species is not surprising, taking into account the pressures they undergo (see Section 4.1).



7 Conclusion and outlook

Despite significant efforts across Member States, biodiversity continues to decline and faces deteriorating trends in the greater part of Europe. The majority of protected habitats and species have a poor or bad status as a result of ongoing pressures from changes in land and sea use, overexploitation (e.g. relating to agriculture, fisheries and hunting) and unsustainable management practices (e.g. agricultural or forestry management). These drivers are compounded by the modification of water regimes, pollution, invasive alien species and the effects of climate change.

Yet there are also positive developments being reported, and a number of species and habitats have shown improved status and trends. Forests, mammals and birds (e.g. several birds of prey) are among the top beneficiaries of the conservation measures applied. The Natura 2000 network, which is a cornerstone of the nature directives, often provides an important framework for implementing such measures, and positive effects have been reported both within the network and globally (with slightly higher positive impacts inside the network than globally). This correlation between effective management measures and the successful protection of species and habitats highlights the need for wider application of measures to increase the scope of the impact.

These improvements were, however, insufficient to achieve the aims of the EU Biodiversity Strategy to 2020. None of the groups studied met their targets, and habitats and birds lag particularly far behind. These groups saw less than half of the improvements needed to achieve their targets.

The new EU Biodiversity Strategy for 2030 aims to address these gaps and ensure that ecosystems are healthy, resilient to climate change, rich in biodiversity and deliver the range of services essential for citizens' prosperity and well-being. Ambitious targets address the main drivers of biodiversity loss and aim to reduce key pressures on nature and ecosystem services in the EU. The strategy focuses on restoring ecosystems rather than on merely averting degradation and biodiversity loss and responds to key

gaps and pressures identified in the state of nature assessments, for example:

- **Agricultural pressures:** increase organic farming to more than 25 %; reduce the overall use of and risk from pesticides by 50 % by 2030; provide space for wild animals, plants, pollinators and natural pest regulators; and recover at least 10 % of agricultural area as high-diversity landscape features.
- **Need for increased restoration:** an ambitious EU nature restoration plan will be developed by 2021 to improve protection of intact habitats and restore degraded areas, e.g. by more effectively protecting marine habitats and restoring at least 25 000 km of rivers to free-flowing rivers by 2030.
- **Exploitation of forest resources and extensive management:** propose a dedicated EU forest strategy in 2021, including a roadmap for planting 3 billion trees by 2030 and the strict protection of all remaining EU primary and old-growth forests.
- **Insufficiency of the current Natura 2000 network:** protect at least 30 % of the land and 30 % of the sea in the EU (including Natura 2000 and nationally designated areas), whereby one third of protected areas will be strictly protected; effectively manage and monitor all protected areas, defining clear conservation objectives and measures; and improve habitat connectivity for a coherent and resilient Trans-European Nature†Network.

Through these ambitious objectives, the new strategy offers a great opportunity to halt or reverse biodiversity decline. Nevertheless, several pressures outlined in the state of nature assessments are not directly addressed by quantitative goals (e.g. urbanisation and tourism development-related pressures, leading to fragmentation and coastal degradation as well as legal hunting, which particularly affects birds).

Additional effort is also needed within the nature directives to improve monitoring capacities within

Member States so that they can support the targets outlined. Although the state of nature assessments show improved data availability, many data gaps ('unknowns') persist, especially for marine species. Furthermore, new indicators and data are necessary to be able to duly evaluate the role of the Natura 2000 network in achieving the objectives of the nature directives and the targets of the EU Biodiversity Strategy for 2030. Emerging challenges such as climate change have also been raised by Member States as a growing threat to habitats and species (not least because of the increased spread of invasive alien species). Expected changes in abundance and distribution (e.g. a potential northwards migration) should thus also be addressed in future reporting under the nature directives and taken into account within the context of the new strategy.

Finally, for the EU Biodiversity Strategy for 2030 to be effective, implementation of measures has to be significantly increased compared with the EU Biodiversity Strategy to 2020. Tremendous efforts are needed to put the cross-sectoral approach into practice, such as adapting the common agricultural fisheries policies to be tools for protecting biodiversity. Additional requirements for effectiveness include setting standards (e.g. for the ecological quality of new designated areas and restoration areas) and ensuring sufficient financial and human resources to put in place a robust governance and policy framework for nature conservation, achieve the 2030 targets, and support the post-2020 global biodiversity framework of the United Nations Convention on Biological Diversity.



Abbreviations

AEWA	Agreement on the Conservation of African-Eurasian Migratory Waterbirds
CAP	Common agricultural policy
CLC	Corine Land Cover
EEA	European Environment Agency
EIP-Agri	European Innovation Partnership 'Agricultural productivity and sustainability'
EBP	Euro Bird Portal
ETC/BD	European Topic Centre on Biological Diversity
EU	European Union
EU-28	The 28 Member States of the EU (1 July 2013 to 31 January 2020)
EU-27	EU-28 minus UK (after 31 January 2020)
EUNIS	European University Information Systems organisation
HNV	High nature value
IAS	Invasive alien species
IPBES	Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
IUCN	International Union for Conservation of Nature
MS	Member State
MSAP	Multi-species action plan
PECPMS	Pan-European Common Bird Monitoring Scheme
SAC	Special Area of Conservation
SAP	Species action plan
SDGs	Sustainable Development Goals
SCI	Site of Community Importance
SPA	Special Protection Area

References

- Alonso, I., et al., 2012, 'Carbon storage by habitat: Review of the evidence of the impacts of management decisions and condition of carbon stores and sources', Natural England Research Report NERR043, Natural England, York, UK.
- Anderson, E., and Mammides, C. 2020, 'Changes in land-cover within high nature value farmlands inside and outside Natura 2000 sites in Europe: a preliminary assessment', *Ambio*, pp. 1-14.
- Arnett, E. B., et al., 2008, 'Patterns of bat fatalities at wind energy facilities in North America', *Journal of Wildlife Management* 72(1), pp. 61-78.
- BirdLife International, 2009, 'A landscape approach to conservation can help tackle climate change' (<http://datazone.birdlife.org/sowb/casestudy/a-landscape-approach-to-conservation-can-help-tackle-climate-change>) accessed 20 April 2020.
- BirdLife International, 2011, 'Seeds of success. How agri-environment can yield results for nature and farming' (<https://www.birdlife.org/sites/default/files/attachments/Seeds%20of%20success.pdf>) accessed 5 May 2020.
- BirdLife International, 2012, 'More research is needed into the impact of wave and tidal energy devices on seabirds' (<http://datazone.birdlife.org/more-research-is-needed-into-the-impact-of-wave-and-tidal-energy-devices-on-seabirds>) accessed 30 April 2020.
- BfN — Bund für Naturschutz, 2020, 'Impacts on marine species and habitats' (<https://www.bfn.de/en/activities/marine-nature-conservation/pressures-on-the-marine-environment/offshore-wind-power/impacts-on-marine-species-and-habitats.html>) accessed 30 April 2020.
- BLUEREEF, 2013, 'Best practice for restoration of stone reefs in Denmark (codes of conduct)', LIFE06 NAT/DK000159, Final Technical Report, Annex 9, June 2013 (https://ec.europa.eu/environment/life/project/Projects/index.cfm?fuseaction=home.showFile&rep=file&fil=BLUEREEF_Best_Practices_EN.pdf) accessed 10 August 2020.
- CABI, 2020, 'Invasive Species Compendium', CAB International (<https://www.cabi.org/isc/>) accessed 18 April 2020.
- Davis et al., 2020, 'Literature Review: The ecological effectiveness of the Natura 2000 Network', European Topic Centre for Biological Diversity.
- de Rigo, D., et al., 2017, 'Forest fire danger extremes in Europe under climate change: Variability and uncertainty', Publications Office of the European Union, Luxembourg.
- Directorate-General for Environment, 2017, 'Reporting under Article 17 of the Habitats Directive: Explanatory notes and guidelines for the period 2013-2018', European Commission, Brussels.
- EC, 2016, 'Fitness check of the EU Nature Legislation (Birds and Habitats Directives)', Commission Staff Working Document (SWD(2016) 472 final) (https://ec.europa.eu/info/sites/info/files/swd-2016-472-final_en.pdf) accessed 14 August 2020.
- EC, 2018, Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions — EU Pollinators Initiative (COM(2018) 395) (<https://eur-lex.europa.eu/legal-content/EN/TXT/?qid=1528213737113&uri=CELEX:52018DC0395>) accessed 14 August 2020.
- EC, 2019a, Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions — The European Green Deal (COM(2019) 640 final of 11 December 2019) (https://ec.europa.eu/info/sites/info/files/european-green-deal-communication_en.pdf) accessed 14 August 2020.
- EC, 2019b, 'The EU Nature Directives: Protecting Europe's marine biodiversity', European Commission (<https://ec.europa.eu/environment/nature/natura2000/marine/docs/Marine%20brochure%20WEB.pdf>) accessed 15 March 2020.

- EC, 2019c, 'Stopping forest fires starts early in Hungary', European Commission, 7 August (<https://ec.europa.eu/easme/en/news/stopping-forest-fires-starts-early-hungary>) accessed 29 July 2020.
- EC, 2019d, 'First EU action plan for restoring a key habitat for pollinators', European Commission, 18 November (https://ec.europa.eu/info/news/first-eu-action-plan-restoring-key-habitat-wild-pollinators-2019-nov-18_en) accessed 29 July 2020.
- EC, 2019e, 'Evaluation of the impact of the CAP on habitats, landscapes, biodiversity', Publications Office of the European Union, Luxembourg (https://ec.europa.eu/info/sites/info/files/food-farming-fisheries/key_policies/documents/ext-eval-biodiversity-final-report_2020_en.pdf) accessed 14 August 2020.
- EC, 2020a, 'The Atlantic region', European Commission (https://ec.europa.eu/environment/nature/natura2000/biogeog_regions/atlantic/index_en.htm) accessed 15 March 2020.
- EC, 2020b, COM(2020) 380 final https://ec.europa.eu/info/files/communication-eu-biodiversity-strategy-2030-bringing-nature-back-our-lives_en
- EC, 2020c, https://ec.europa.eu/food/sites/food/files/safety/docs/f2f_action-plan_2020_strategy-info_en.pdf
- EC, 2020d, COM(2020) 380 final https://ec.europa.eu/environment/nature/biodiversity/strategy/index_en.htm
- EC, 2020e, 'Natura 2000 in the marine environment' European Commission (https://ec.europa.eu/environment/nature/natura2000/marine/index_en.htm) accessed 15 March 2020.
- EC, 2020f, 'How will we be affected?' (https://ec.europa.eu/clima/policies/adaptation/how_en) accessed 20 May 2020.
- ECA, 2020, 'Biodiversity on farmland: CAP contribution has not halted the decline', European Court of Auditors. European Union, 2020.
- EEA, 2016, 'European forest ecosystems — State and trends', EEA Report No 5/2016, European Environment Agency.
- EEA, 2017, 'Climate change, impacts and vulnerability in Europe 2016' EEA Report No 1/2017, European Environment Agency.
- EEA, 2020a, 'Preliminary review of management effectiveness in the EU's Natura 2000 network of protected areas', European Environment Agency (link to be included end September).
- EEA, 2020b, 'Abundance and distribution of selected European species (SEBI 001)', European Environment Agency (<https://www.eea.europa.eu/data-and-maps/indicators/abundance-and-distribution-of-selected-species-8/assessment-1>) accessed 29 July 2020.
- EEA, 2020c, 'Natura 2000 data — the European network of protected sites' (<https://www.eea.europa.eu/data-and-maps/data/natura-11>) accessed 29 July 2020.
- EFH, 2019, 'The implementation of the EU 2020 Biodiversity Strategy and recommendations for the post 2020 Biodiversity Strategy', European Habitats Forum (https://www.iucn.org/sites/dev/files/content/documents/eu_2020_biodiversity_strategy_and_recommendations_for_post_2020.pdf) accessed 14 August 2020.
- EU, 1992, Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora, OJ L 206, 22.7.1992, pp. 7-50.
- EU, 2008, Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive) (https://ec.europa.eu/environment/marine/eu-coast-and-marine-policy/marine-strategy-framework-directive/index_en.htm) accessed 29 July 2020.
- EU, 2009, Directive 2009/147/EC of the European Parliament and of the Council of 30 November 2009 on the conservation of wild birds, OJ L 20, 26.1.2018, pp. 7-25.
- EU, 2014, Regulation (EU) No 1143/2014 of the European Parliament and of the Council of 22 October 2014 on the prevention and management of the introduction and spread of invasive alien species, OJ L 317, 4.11.2014, pp. 35-55.
- EU, 2018, Directive (EU) 2018/2001 on the promotion of the use of energy from renewable sources.
- Eurostat, 2020a, 'Farms and farmland in the European Union — statistics' (https://ec.europa.eu/eurostat/statistics-explained/index.php/Farms_and_farmland_in_the_European_Union_-_statistics) accessed 20 May 2020.

- Eurostat, 2020b, 'What do we produce in the EU?' (<https://ec.europa.eu/eurostat/cache/infographs/energy/bloc-2b.html>) accessed 30 April 2020.
- FERN, 2020, 'European Commission faces major hurdle to protect and restore forests' (<https://www.fern.org/news-resources/european-commission-faces-major-hurdle-to-protect-and-restore-forests-2148/>) accessed 16 June 2020.
- Ferreras, P. and MacDonald, D. W., 1999, 'The impact of American mink *Mustela vison* on water birds in the upper Thames' *Journal of Applied Ecology* 36(5), pp. 701-708.
- FireLife, 2020, 'Hungarian forest fire prevention and training program' (https://ec.europa.eu/environment/life/project/Projects/index.cfm?fuseaction=search.dspPage&n_proj_id=5062) accessed 29 July 2020.
- Forest Europe, 2015, 'State of Europe's forests, 2015 report' (<http://foresteurope.org/state-europes-forests-2015-report/>) accessed 5 June 2020.
- Furness, R. W., et al., 2013, 'Assessing vulnerability of seabird populations to offshore wind farms', *Journal of Environmental Management* 119C, pp. 56-66 (<https://doi.org/10.1016/j.jenvman.2013.01.025>).
- Garthe, S., et al., 2017, 'Possible impacts of offshore wind farms on seabirds: a pilot study in Northern Gannets in the southern North Sea', *Journal of Ornithology* 158, pp. 345-349.
- Goldstein, A., et al., 2020, 'Protecting irrecoverable carbon in Earth's ecosystems', *Nature Climate Change* 10, pp. 287-295 (<https://doi.org/10.1038/s41558-020-0738-8>).
- Gove, B., et al., 2013, 'Wind farms and birds: an updated analysis of the effects of wind farms on birds, and best practice guidance on integrated planning and impact assessment', Bern Convention Bureau Meeting, Strasbourg, 26 August.
- Halada, L., et al., 2011, 'Which habitats of European importance depend on agricultural practices?', *Biodiversity and Conservation* 20(11), pp. 2365-2378.
- Hettelingh J-P., et al., 2017, 'European critical loads: database, biodiversity and ecosystems at risk', CCE Final Report 2017, Coordination Centre for Effects, RIVM Report 2017-0155, Bilthoven, Netherlands.
- Hirschfeld, A., et al., 2019, 'Bird hunting in Europe: an analysis of bag figures and the potential impact on the conservation of threatened species', *British Birds*, pp. 153-166.
- IPBES (2019): 'Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services', S. Díaz, J. Settele, E. S. Brondízio E.S., H. T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A. Purvis, J. Razzaque, B. Reyers, R. Roy Chowdhury, Y. J. Shin, I. J. Visseren-Hamakers, K. J. Willis, and C. N. Zayas (eds.). IPBES secretariat, Bonn, Germany. 56 pages. (<https://doi.org/10.5281/zenodo.3553579>) accessed 14 August 2020.
- Keller et al., 2020, 'European Breeding Bird Atlas 2: Distribution, Abundance and Change', European Bird Census Council & Lynx Ediciones, Barcelona.
- Keränen, I., et al., 2020, 'Boreal Baltic coastal meadows in Finland', Good practice example from Finland (sent by e-mail), Finnish Environment Institute, Ministry of the Environment, Centre for Economic Development, Transport and the Environment, Southwest Finland, Parks & Wildlife Finland.
- Kudrnovsky, H., et al., 2020, 'Report for a list of habitats important for pollinators', Final report for European Environment Agency, European Topic Centre for Biological Diversity.
- Langhout, W., 2019, 'The EU Biodiversity Strategy. Progress report 2011-2018', BirdLife International.
- Mineau, P., and Whiteside, M., 2013, 'Pesticide acute toxicity is a better correlate of US grassland bird declines than agricultural intensification', *PLoS ONE* 8(2).
- NEEMO, forthcoming, 'The impact of LIFE: Benefits to nature, biodiversity and society', Report to the European Commission.
- Olson, D.H. and Saenz, D., 2013, 'Climate change and amphibians', US Department of Agriculture, Forest Service, Climate Change Resource Centre (www.fs.usda.gov/ccrc/topics/wildlife/amphibians/) accessed 10 June 2020.
- Paillet, Y., et al., 2010, 'Biodiversity differences between managed and unmanaged forests: meta-analysis of species richness in Europe', *Conservation Biology* 24(1), pp. 101-112.

- Pe'er, G., et al., 2017, 'Is the CAP fit for purpose? An evidence-based fitness check assessment', German Centre for Integrative Biodiversity Research, Halle-Jena-Leipzig.
- Pellissier, V., et al., 2019, 'Effects of Natura 2000 on nontarget bird and butterfly species based on citizen science data', *Conservation Biology* 34(3), pp. 666-676 (<https://doi.org/10.1111/cobi.13434>).
- Polish Society for the Protection of Birds, 2020, 'Aquatic warbler' (<https://otop.org.pl/our-projects/we-protect/aquatic-warbler/?lang=en>) accessed 29 July 2020.
- Ramírez, I., et al., 2017, 'How well is the EU protecting its seabirds? Progress in implementing the Birds Directive at sea', *Marine Policy* 81, pp. 179-184 (<http://dx.doi.org/10.1016/j.marpol.2017.03.034>).
- Roekaerts, M., et al., 2020, 'Draft summary reporting under Resolution No 8 (2012) First Observations', Convention on the Conservation of European Wildlife and Natural Habitats.
- Rollins, K. E., et al., 2012, 'A forensic investigation into the aetiology of bat mortality at a wind farm: barotrauma or traumatic injury?', *Veterinary Pathology* 49(2), pp. 362-371.
- Röschel et al., 2020, 'State of nature in the EU — Methodological paper. Methodologies under the Nature Directives reporting 2013-2018 and analysis for the State of nature 2020', ETC/BD Technical paper 2/2020 (<https://www.eionet.europa.eu/etcs/etc-bd/products/etc-bd-reports/etc-bd-technical-paper-2-2020-state-of-nature-in-the-eu-methodological-paper-methodologies-under-the-nature-directives-reporting-2013-2018-and-analysis-for-the-state-of-nature-2000>) accessed 29 July 2020.
- Sánchez-Bayo, F. and Wyckhuys, K. A., 2019, 'Worldwide decline of the entomofauna: a review of its drivers', *Biological Conservation* 232, pp. 8-27 (<https://doi.org/10.1016/j.biocon.2019.01.020>).
- Searchinger, T. D., et al., 2018, 'Europe's renewable energy directive poised to harm global forests', *Nature Communications* 9, p. 3741 (<https://doi.org/10.1038/s41467-018-06175-4>).
- Sotoriv, M., 2017, 'Natura 2000 and forests — Assessing the state of implementation and effectiveness', European Forest Institute.
- Spitzen-van der Sluijs, A., et al., 2016, 'Expanding distribution of lethal amphibian fungus *Batrachochytrium salamandrivorans* in Europe', *Emerging Infectious Diseases* 22(7), p. 1286.
- Tucker, G., et al., 2019, 'Study on identifying the drivers of successful implementation of the Birds and Habitats Directives', Report to the European Commission, DG Environment on Contract ENV.F.1/FRA/2014/0063, Institute for European Environmental Policy, Brussels.
- Vilén, T. et al., 2015, 'Increasing carbon sinks in European forests: effects of afforestation and changes in mean growing stock volume', *Forestry* 89(1), p. 8290.
- Warwick-Evans, V., et al., 2017, 'Predicting the impacts of wind farms on seabird: an individual-based model', *Journal of Applied Ecology* 55(2), pp. 503-515 (<https://doi.org/10.1111/1365-2664.12996>).
- Zehetmaira, T., et al., 2015, 'Poor effectiveness of Natura 2000 beech forests in protecting forest-dwelling bats', *Journal for Nature Conservation* 23, pp. 53-60.

European Environment Agency

State of nature in the EU

Results from reporting under the nature directives 2013-2018

2020 — 142 pp. — 21.0 x 29.7 cm

ISBN 978-92-9480-259-0

doi:10.2800/088178

Getting in touch with the EU

In person

All over the European Union there are hundreds of Europe Direct information centres. You can find the address of the centre nearest you at: https://europa.eu/european-union/contact_en

On the phone or by email

Europe Direct is a service that answers your questions about the European Union. You can contact this service:

- by freephone: 00 800 6 7 8 9 10 11 (certain operators may charge for these calls),
- at the following standard number: +32 22999696 or
- by email via: https://europa.eu/european-union/contact_en

Finding information about the EU

Online

Information about the European Union in all the official languages of the EU is available on the Europa website at: https://europa.eu/european-union/index_en

EU publications

You can download or order free and priced EU publications at: <https://publications.europa.eu/en/publications>.

Multiple copies of free publications may be obtained by contacting Europe Direct or your local information centre (see https://europa.eu/european-union/contact_en).

European Environment Agency
Kongens Nytorv 6
1050 Copenhagen K
Denmark

Tel.: +45 33 36 71 00
Web: eea.europa.eu
Enquiries: eea.europa.eu/enquiries



Publications Office
of the European Union

